



VACCINES REQUIRED FOR CHILD CARE/PRESCHOOL ATTENDANCE

July 1, 2018 – June 30, 2019

	Hepatitis B	DTaP (Diphtheria, Tetanus, Pertussis)	Hib (<i>Haemophilus influenzae</i> type B)	Polio	PCV (Pneumococcal Conjugate)	MMR (Measles, Mumps, Rubella)	Varicella (Chickenpox)
By 3 Months (on or before last day of mo 2)	2 doses May get Dose 1 at birth and Dose 2 as early as 1 month of age	1 dose	1 dose	1 dose	1 dose	Not given before 12 months of age	Not given before 12 months of age
By 5 Months (on or before last day of mo 4)	2 doses	2 doses	2 doses	2 doses May get Dose 2 as early as 4 months of age	2 doses		
By 7 Months (on or before last day of mo 6)	2 doses	3 doses May get Dose 3 as early as 6 months of age	3 doses	2 doses	3 doses		
By 16 Months (on or before last day of mo 15)	2 doses	3 doses	4 doses	2 doses	4 doses*	1 dose May get Dose 1 as early as 12 months of age	1 dose May get Dose 1 as early as 12 months of age OR Healthcare provider verified disease
By 19 Months (on or before last day of mo 18)	3 doses	4 doses May get Dose 4 as early as 12 months as long as 6 months separate Dose 3 and Dose 4	4 doses	3 doses	4 doses*	1 dose	1 dose OR Healthcare provider verified disease
By 7 Years (on or before last day of year 6) or by Kindergarten Entry	3 doses	5 doses	Not given after 5 years of age unless child has medical condition	4 doses	Not given after 5 years of age unless child has medical condition	2 doses	2 doses OR Healthcare provider verified disease

*Some children may get 5 total doses. A single supplemental dose of PCV13 is recommended, but not required, for all children aged 14–59 months who got 4 doses of PCV7.

- School-aged children (K-12) in before and after-school programs must meet the immunization requirements for their grade in school.
- Find information on other vaccines recommended, but not required, for child care/preschool attendance: www.immunize.org/cdc/schedules/
- Review the Individual Vaccine Requirements Summary for more detailed information: www.doh.wa.gov/Immunization/schoolandchildcare/VaccineRequirements.aspx

Minimum Age & Interval for Valid Vaccine Doses

Vaccine	Dose #	Minimum Age	Minimum Interval Between Doses	Notes
Hepatitis B (HepB)	Dose 1	Birth	4 weeks between Dose 1 & 2	<ul style="list-style-type: none"> The final dose in the series should be given at least 24 weeks of age.
	Dose 2	4 weeks	8 weeks between Dose 2 & 3	
	Dose 3	24 weeks	16 weeks between Dose 1 & 3	
Diphtheria, Tetanus, and Pertussis (DTaP)	Dose 1	6 weeks	4 weeks between Dose 1 & 2	<ul style="list-style-type: none"> Typical vaccine schedule: 2, 4, 6, and 15-18 months of age. Recommended: 6 months between Dose 3 and Dose 4, but at least 4 months minimum interval acceptable.
	Dose 2	10 weeks	4 weeks between Dose 2 & 3	
	Dose 3	14 weeks	6 months between Dose 3 & 4	
	Dose 4	12 months	6 months between Dose 4 & 5	
	Dose 5	4 years	--	
<i>Haemophilus influenzae</i> type B (Hib)	Dose 1	6 weeks	4 weeks between Dose 1 & 2	<ul style="list-style-type: none"> If all 3 doses of PedvaxHIB given, only need 3 doses total. Dose 3 must be ≥ 12 months of age. Only one dose required if the dose given on or after 15 months of age. Review the Individual Vaccine Requirements Summary for minimum doses required: www.doh.wa.gov/immunization/schoolandchildcare/VaccineRequirements.aspx
	Dose 2	10 weeks	4 weeks between Dose 2 & 3	
	Dose 3	14 weeks	8 weeks between Dose 3 & 4	
	Dose 4	12 months	--	
Pneumococcal Conjugate (PCV7 or PCV13)	Dose 1	6 weeks	4 weeks between Dose 1 & 2	<ul style="list-style-type: none"> A single supplemental dose of PCV13 recommended for all children 14–59 months of age who got 4 doses of PCV7. Only one dose required if the dose given on or after 24 months of age. Review the Individual Vaccine Requirements Summary for minimum doses required: www.doh.wa.gov/immunization/schoolandchildcare/VaccineRequirements.aspx
	Dose 2	10 weeks	4 weeks between Dose 2 & 3	
	Dose 3	14 weeks	8 weeks between Dose 3 & 4	
	Dose 4	12 months	--	
Polio (IPV or OPV)	Dose 1	6 weeks	4 weeks between Dose 1 & 2	<ul style="list-style-type: none"> Three doses acceptable if child got Dose 3 on or after the 4th birthday.
	Dose 2	10 weeks	4 weeks between Dose 2 & 3	
	Dose 3	14 weeks	6 months between Dose 3 & 4	
	Dose 4	4 years	--	
Measles, Mumps, and Rubella (MMR or MMRV)	Dose 1	12 months	4 weeks between Dose 1 & 2	<ul style="list-style-type: none"> MMRV (MMR + varicella) may be used in place of separate MMR and varicella vaccines. Must get the same day as VAR OR at least 28 days apart. 4-day grace DOES apply between doses of the same live vaccine such as MMR/MMR or MMRV/MMRV. The 4 day grace period DOES NOT apply between Dose 1 and Dose 2 of different live vaccines, such as between MMR and Varicella or between MMR and live flu vaccine.
	Dose 2	13 months	--	
Varicella (chickenpox) (VAR)	Dose 1	12 months	3 months between Dose 1 & 2 (12 months through 12 years) 4 weeks between Dose 1 & 2 (13 years and older)	<ul style="list-style-type: none"> Recommended: 3 months between varicella doses, but at least 28 days minimum interval acceptable. Minimum age of 13 months acceptable. Must get the same day as MMR OR at least 28 days apart. 4-day grace DOES apply between doses of the same live vaccine; DOES NOT apply between doses of different live vaccines.
	Dose 2	15 months	--	

If you have a disability and need this document in another format, please call 1-800-525-0127 (TDD/TTY call 711).

DOH 348-053 December 2017