

2

The Chemical Context of Life



▲ **Figure 2.1** Who tends this “garden”?

variety of plant life. Visitors traveling near the Amazon’s headwaters in Peru are therefore surprised to come across tracts of forest like that seen in the foreground of the photo in **Figure 2.1**. This patch is almost completely dominated by a single plant species—a small flowering tree called *Duroia hirsuta*. Travelers may wonder if the plot of land is planted and maintained by local people, but the indigenous people are as mystified as the visitors. They call these stands of *Duroia* trees “devil’s gardens,” from a legend attributing them to an evil forest spirit.

Seeking a scientific explanation, a research team at Stanford University recently solved the “devil’s garden” mystery. **Figure 2.2** describes their main experiment. The researchers showed that the “farmers” who create and maintain these gardens are actually ants that live in the hollow stems of the *Duroia* trees. The ants do not plant the *Duroia* trees, but they prevent other plant species from growing in the garden by injecting intruders with a poisonous chemical. In this way, the ants create space for the growth of the *Duroia* trees that serve as their home. With the ability to maintain and expand its habitat, a single colony of devil’s garden ants can live for hundreds of years.

The chemical used by the ants to weed their garden turns out to be formic acid. This substance is produced by many species of ants and in fact got its name from the Latin word for ant, *formica*. For many ant species, the formic acid probably serves as a disinfectant that protects the ants against microbial parasites. The devil’s garden ant is the first ant species found to use formic acid as an herbicide, an important addition to the list of functions mediated by chemicals in the insect world. Scientists have long known that chemicals play a major role in insect communication, attraction of mates, and defense against predators.

Research on devil’s gardens is only one example of the relevance of chemistry to the study of life. Unlike a list of college courses, nature is not neatly packaged into the individual natural sciences—biology, chemistry, physics, and so forth. Biologists specialize in the study of life, but organisms and their environments are natural systems to which the concepts of chemistry and physics apply. Biology is a multidisciplinary science.

This unit of chapters introduces some basic concepts of chemistry that apply to the study of life. We will make many connections to the themes introduced in Chapter 1. One of these themes is the organization of life into a hierarchy of structural levels, with additional properties emerging at each successive level. In this unit, we will see how emergent properties are apparent at the lowest levels of biological organization—such as the ordering of atoms into molecules and the interactions of those molecules within cells. Somewhere in the transition from molecules to cells, we will cross the blurry boundary between nonlife and life. This chapter focuses on the chemical components that make up all matter.

KEY CONCEPTS

- 2.1** Matter consists of chemical elements in pure form and in combinations called compounds
- 2.2** An element’s properties depend on the structure of its atoms
- 2.3** The formation and function of molecules depend on chemical bonding between atoms
- 2.4** Chemical reactions make and break chemical bonds

OVERVIEW

A Chemical Connection to Biology

The Amazon rain forest in South America is a showcase for the diversity of life on Earth. Colorful birds, insects, and other animals live in a densely-packed environment of trees, shrubs, vines, and wildflowers, and an excursion along a waterway or a forest path typically reveals a lush


▼ Figure 2.2

INQUIRY

What creates “devil’s gardens” in the rain forest?


EXPERIMENT Working under Deborah Gordon and with Michael Greene, graduate student Megan Frederickson sought the cause of “devil’s gardens,” stands of a single species of tree, *Duroia hirsuta*. One hypothesis was that ants living in these trees, *Myrmelachista schumanni*, produce a poisonous chemical that kills trees of other species; another was that the *Duroia* trees themselves kill competing trees, perhaps by means of a chemical.

To test these hypotheses, Frederickson did field experiments in Peru. Two saplings of a local nonhost tree species, *Cedrela odorata*, were planted inside each of ten devil’s gardens. At the base of one sapling, a sticky insect barrier was applied; the other was unprotected. Two more *Cedrela* saplings, with and without barriers, were planted about 50 meters outside each garden.



The researchers observed ant activity on the *Cedrela* leaves and measured areas of dead leaf tissue after one day. They also chemically analyzed contents of the ants’ poison glands.

RESULTS The ants made injections from the tips of their abdomens into leaves of unprotected saplings in their gardens (see photo). Within one day, these leaves developed dead areas (see graph). The protected saplings were uninjured, as were the saplings planted outside the gardens. Formic acid was the only chemical detected in the poison glands of the ants.



CONCLUSION Ants of the species *Myrmelachista schumanni* kill non-host trees by injecting the leaves with formic acid, thus creating hospitable habitats (devil’s gardens) for the ant colony.

SOURCE M. E. Frederickson, M. J. Greene, and D. M. Gordon, “Devil’s gardens” bedevilled by ants, *Nature* 437:495–496 (2005).

INQUIRY IN ACTION Read and analyze the original paper in *Inquiry in Action: Interpreting Scientific Papers*.

WHAT IF? What would be the results if the unprotected saplings’ inability to grow in the devil’s gardens was caused by a chemical released by the *Duroia* trees rather than by the ants?

CONCEPT 2.1


Matter consists of chemical elements in pure form and in combinations called compounds

Organisms are composed of **matter**, which is defined as anything that takes up space and has mass.* Matter exists in many diverse forms. Rocks, metals, oils, gases, and humans are just a few examples of what seems an endless assortment of matter.

Elements and Compounds

Matter is made up of elements. An **element** is a substance that cannot be broken down to other substances by chemical reactions. Today, chemists recognize 92 elements occurring in nature; gold, copper, carbon, and oxygen are examples. Each element has a symbol, usually the first letter or two of its name. Some symbols are derived from Latin or German; for instance, the symbol for sodium is Na, from the Latin word *natrium*.

A **compound** is a substance consisting of two or more different elements combined in a fixed ratio. Table salt, for example, is sodium chloride (NaCl), a compound composed of the elements sodium (Na) and chlorine (Cl) in a 1:1 ratio. Pure sodium is a metal, and pure chlorine is a poisonous gas. When chemically combined, however, sodium and chlorine form an edible compound. Water (H₂O), another compound, consists of the elements hydrogen (H) and oxygen (O) in a 2:1 ratio. These are simple examples of organized matter having emergent properties: A compound has characteristics different from those of its elements (**Figure 2.3**).



▲ **Figure 2.3** The emergent properties of a compound. The metal sodium combines with the poisonous gas chlorine, forming the edible compound sodium chloride, or table salt.

*Sometimes we substitute the term weight for mass, although the two are not identical. Mass is the amount of matter in an object, whereas the weight of an object is how strongly that mass is pulled by gravity. The weight of an astronaut walking on the moon is approximately 1/6 the astronaut’s weight on Earth, but his or her mass is the same. However, as long as we are earthbound, the weight of an object is a measure of its mass; in everyday language, therefore, we tend to use the terms interchangeably.

The Elements of Life

Of the 92 natural elements, about 20–25% are **essential elements** that an organism needs to live a healthy life and reproduce. The essential elements are similar among organisms, but there is some variation—for example, humans need 25 elements, but plants need only 17.

Just four elements—oxygen (O), carbon (C), hydrogen (H), and nitrogen (N)—make up 96% of living matter. Calcium (Ca), phosphorus (P), potassium (K), sulfur (S), and a few other elements account for most of the remaining 4% of an organism's mass. **Trace elements** are required by an organism in only minute quantities. Some trace elements, such as iron (Fe), are needed by all forms of life; others are required only by certain species. For example, in vertebrates (animals with backbones), the element iodine (I) is an essential ingredient of a hormone produced by the thyroid gland. A daily intake of only 0.15 milligram (mg) of iodine is adequate for normal activity of the human thyroid. An iodine deficiency in the diet causes the thyroid gland to grow to abnormal size, a condition called goiter. Where it is available, eating seafood or iodized salt reduces the incidence of goiter. All the elements needed by the human body are listed in **Table 2.1**.

Some naturally occurring elements are toxic to organisms. In humans, for instance, the element arsenic has been linked to numerous diseases and can be lethal. In some areas of the world, arsenic occurs naturally and can make its way into the groundwater. As a result of using water from drilled wells in southern Asia, millions of people have been inadvertently exposed to arsenic-laden water. Efforts are under way to reduce arsenic levels in their water supply.

Table 2.1 Elements in the Human Body

Element	Symbol	Percentage of Body Mass (including water)	
Oxygen	O	65.0%	} 96.3%
Carbon	C	18.5%	
Hydrogen	H	9.5%	
Nitrogen	N	3.3%	
Calcium	Ca	1.5%	} 3.7%
Phosphorus	P	1.0%	
Potassium	K	0.4%	
Sulfur	S	0.3%	
Sodium	Na	0.2%	
Chlorine	Cl	0.2%	
Magnesium	Mg	0.1%	
Trace elements (less than 0.01% of mass): Boron (B), chromium (Cr), cobalt (Co), copper (Cu), fluorine (F), iodine (I), iron (Fe), manganese (Mn), molybdenum (Mo), selenium (Se), silicon (Si), tin (Sn), vanadium (V), zinc (Zn)			



▲ Figure 2.4 Serpentine plant community. The plants in the large photo are growing on serpentine soil, which contains elements that are usually toxic to plants. The insets show a close-up of serpentine rock and one of the plants, a Tiburon Mariposa lily.

Case Study: Evolution of Tolerance to Toxic Elements

EVOLUTION Some species have become adapted to environments containing elements that are usually toxic. A compelling example is found in serpentine plant communities. Serpentine is a jade-like mineral that contains toxic elements such as chromium, nickel, and cobalt. Although most plants cannot survive in soil that forms from serpentine rock, a small number of plant species have adaptations that allow them to do so (**Figure 2.4**). Presumably, variants of ancestral, nonserpentine species arose that could survive in serpentine soils, and subsequent natural selection resulted in the distinctive array of species we see in these areas today.

CONCEPT CHECK 2.1

- 1. MAKE CONNECTIONS** Review the discussion of emergent properties in Chapter 1 (p. 3). Explain how table salt has emergent properties.
- 2.** Is a trace element an essential element? Explain.
- 3.** In humans, iron is a trace element required for the proper functioning of hemoglobin, the molecule that carries oxygen in red blood cells. What might be the effects of an iron deficiency?
- 4. MAKE CONNECTIONS** Review the discussion of natural selection in Chapter 1 (pp. 14–16) and explain how natural selection might have played a role in the evolution of species that are tolerant of serpentine soils.

For suggested answers, see Appendix A.

CONCEPT 2.2


An element's properties depend on the structure of its atoms

Each element consists of a certain type of atom that is different from the atoms of any other element. An **atom** is the smallest unit of matter that still retains the properties of an element. Atoms are so small that it would take about a million of them to stretch across the period printed at the end of this sentence. We symbolize atoms with the same abbreviation used for the element that is made up of those atoms. For example, the symbol C stands for both the element carbon and a single carbon atom.

Subatomic Particles

Although the atom is the smallest unit having the properties of an element, these tiny bits of matter are composed of even smaller parts, called *subatomic particles*. Physicists have split the atom into more than a hundred types of particles, but only three kinds of particles are relevant here: **neutrons**, **protons**, and **electrons**. Protons and electrons are electrically charged. Each proton has one unit of positive charge, and each electron has one unit of negative charge. A neutron, as its name implies, is electrically neutral.

Protons and neutrons are packed together tightly in a dense core, or **atomic nucleus**, at the center of an atom; protons give the nucleus a positive charge. The electrons form a sort of cloud of negative charge around the nucleus, and it is the attraction between opposite charges that keeps the electrons in the vicinity of the nucleus. **Figure 2.5** shows two commonly used models of the structure of the helium atom as an example.



▲ **Figure 2.5** Simplified models of a helium (He) atom. The helium nucleus consists of 2 neutrons (brown) and 2 protons (pink). Two electrons (yellow) exist outside the nucleus. These models are not to scale; they greatly overestimate the size of the nucleus in relation to the electron cloud.

The neutron and proton are almost identical in mass, each about 1.7×10^{-24} gram (g). Grams and other conventional units are not very useful for describing the mass of objects so minuscule. Thus, for atoms and subatomic particles (and for molecules, too), we use a unit of measurement called the **dalton**, in honor of John Dalton, the British scientist who helped develop atomic theory around 1800. (The dalton is the same as the *atomic mass unit*, or *amu*, a unit you may have encountered elsewhere.) Neutrons and protons have masses close to 1 dalton. Because the mass of an electron is only about 1/2,000 that of a neutron or proton, we can ignore electrons when computing the total mass of an atom.

Atomic Number and Atomic Mass

Atoms of the various elements differ in their number of subatomic particles. All atoms of a particular element have the same number of protons in their nuclei. This number of protons, which is unique to that element, is called the **atomic number** and is written as a subscript to the left of the symbol for the element. The abbreviation ${}^2\text{He}$, for example, tells us that an atom of the element helium has 2 protons in its nucleus. Unless otherwise indicated, an atom is neutral in electrical charge, which means that its protons must be balanced by an equal number of electrons. Therefore, the atomic number tells us the number of protons and also the number of electrons in an electrically neutral atom.

We can deduce the number of neutrons from a second quantity, the **mass number**, which is the sum of protons plus neutrons in the nucleus of an atom. The mass number is written as a superscript to the left of an element's symbol. For example, we can use this shorthand to write an atom of helium as ${}^4_2\text{He}$. Because the atomic number indicates how many protons there are, we can determine the number of neutrons by subtracting the atomic number from the mass number: The helium atom, ${}^4_2\text{He}$, has 2 neutrons. For sodium (Na):



$$\text{Mass number} = \text{number of protons} + \text{neutrons} \\ = 23 \text{ for sodium}$$

$$\text{Atomic number} = \text{number of protons} \\ = \text{number of electrons in a neutral atom} \\ = 11 \text{ for sodium}$$

$$\text{Number of neutrons} = \text{mass number} - \text{atomic number} \\ = 23 - 11 = 12 \text{ for sodium}$$

The simplest atom is hydrogen, ${}^1_1\text{H}$, which has no neutrons; it consists of a single proton with a single electron.

As mentioned earlier, the contribution of electrons to mass is negligible. Therefore, almost all of an atom's mass is concentrated in its nucleus. Because neutrons and protons each have a mass very close to 1 dalton, the mass number is an approximation of the total mass of an atom, called its **atomic mass**. So we might say that the atomic mass of sodium (${}^{23}_{11}\text{Na}$) is 23 daltons, although more precisely it is 22.9898 daltons.





◀ **Figure 2.7 A PET scan, a medical use for radioactive isotopes.** PET, an acronym for positron-emission tomography, detects locations of intense chemical activity in the body. The bright yellow spot marks an area with an elevated level of radioactively labeled glucose, which in turn indicates high metabolic activity, a hallmark of cancerous tissue.

The Energy Levels of Electrons



The simplified models of the atom in Figure 2.5 greatly exaggerate the size of the nucleus relative to the volume of the whole atom. If an atom of helium were the size of a typical football stadium, the nucleus would be the size of a pencil eraser in the center of the field. Moreover, the electrons would be like two tiny gnats buzzing around the stadium. Atoms are mostly empty space.

When two atoms approach each other during a chemical reaction, their nuclei do not come close enough to interact. Of the three kinds of subatomic particles we have discussed, only electrons are directly involved in the chemical reactions between atoms.

An atom's electrons vary in the amount of energy they possess. **Energy** is defined as the capacity to cause change—for instance, by doing work. **Potential energy** is the energy that matter possesses because of its location or structure. For example, water in a reservoir on a hill has potential energy because of its altitude. When the gates of the reservoir's dam are opened and the water runs downhill, the energy can be used to do work, such as turning generators. Because energy has been expended, the water has less energy at the bottom of the hill than it did in the reservoir. Matter has a natural tendency to move to the lowest possible state of potential energy; in this example, the water runs downhill. To restore the potential energy of a reservoir, work must be done to elevate the water against gravity.

The electrons of an atom have potential energy because of how they are arranged in relation to the nucleus. The negatively charged electrons are attracted to the positively charged nucleus. It takes work to move a given electron farther away from the nucleus, so the more distant an electron is from the nucleus, the greater its potential energy. Unlike the continuous flow of water downhill, changes in the potential energy of electrons can occur only in steps of fixed amounts. An electron having a certain amount of energy is something like a ball on a staircase (**Figure 2.8a**). The ball can have different amounts of potential energy, depending on which step it is

(a) A ball bouncing down a flight of stairs provides an analogy for energy levels of electrons, because the ball can come to rest only on each step, not between steps.



(b) An electron can move from one shell to another only if the energy it gains or loses is exactly equal to the difference in energy between the energy levels of the two shells. Arrows in this model indicate some of the stepwise changes in potential energy that are possible.

▲ **Figure 2.8 Energy levels of an atom's electrons.** Electrons exist only at fixed levels of potential energy called electron shells.

on, but it cannot spend much time between the steps. Similarly, an electron's potential energy is determined by its energy level. An electron cannot exist between energy levels.

An electron's energy level is correlated with its average distance from the nucleus. Electrons are found in different **electron shells**, each with a characteristic average distance and energy level. In diagrams, shells can be represented by concentric circles (**Figure 2.8b**). The first shell is closest to the nucleus, and electrons in this shell have the lowest potential energy. Electrons in the second shell have more energy, and electrons in the third shell even more energy. An electron can change the shell it occupies, but only by absorbing or losing an amount of energy equal to the difference in potential energy between its position in the old shell and that in the new shell. When an electron absorbs energy, it moves to a shell farther out from the nucleus. For example, light energy can excite an electron to a higher energy level. (Indeed, this is the first step taken when plants harness the energy of sunlight for photosynthesis, the process that produces food from carbon dioxide and water.) When an electron loses energy, it "falls back" to a shell closer to the nucleus, and the lost energy is usually released to the environment as heat. For example, sunlight excites electrons in the surface of a car to higher energy levels. When the electrons fall back to their original levels, the car's surface heats up. This thermal energy can be transferred to the air or to your hand if you touch the car.



















Electron Distribution and Chemical Properties

The chemical behavior of an atom is determined by the distribution of electrons in the atom's electron shells. Beginning with hydrogen, the simplest atom, we can imagine building the atoms of the other elements by adding 1 proton and 1 electron at a time (along with an appropriate number of neutrons). **Figure 2.9**, an abbreviated version of what is called the *periodic table of the elements*, shows this distribution of electrons for the first 18 elements, from hydrogen (${}_1\text{H}$) to argon (${}_{18}\text{Ar}$). The elements are arranged in three rows, or periods, corresponding to the number of electron shells in their atoms. The left-to-right sequence of elements in each row corresponds to the sequential addition of electrons and protons. (See Appendix B for the complete periodic table.)

Hydrogen's 1 electron and helium's 2 electrons are located in the first shell. Electrons, like all matter, tend to exist in the lowest available state of potential energy. In an atom, this state is in the first shell. However, the first shell can hold no more than 2 electrons; thus, hydrogen and helium are the only elements in the first row of the table. An atom with more than 2 electrons must use higher shells because the first shell

is full. The next element, lithium, has 3 electrons. Two of these electrons fill the first shell, while the third electron occupies the second shell. The second shell holds a maximum of 8 electrons. Neon, at the end of the second row, has 8 electrons in the second shell, giving it a total of 10 electrons.

The chemical behavior of an atom depends mostly on the number of electrons in its *outermost* shell. We call those outer electrons **valence electrons** and the outermost electron shell the **valence shell**. In the case of lithium, there is only 1 valence electron, and the second shell is the valence shell. Atoms with the same number of electrons in their valence shells exhibit similar chemical behavior. For example, fluorine (F) and chlorine (Cl) both have 7 valence electrons, and both form compounds when combined with the element sodium (see Figure 2.3). An atom with a completed valence shell is unreactive; that is, it will not interact readily with other atoms. At the far right of the periodic table are helium, neon, and argon, the only three elements shown in Figure 2.9 that have full valence shells. These elements are said to be *inert*, meaning chemically unreactive. All the other atoms in Figure 2.9 are chemically reactive because they have incomplete valence shells.

First shell	Hydrogen ${}_1\text{H}$		<div><div>2</div><div>Atomic number</div><div>He</div><div>4.00</div><div>Atomic mass</div><div>Element symbol</div></div> <div><div>Helium</div><div>${}_2\text{He}$</div><div>Electron distribution diagram</div></div>				Helium ${}_2\text{He}$	
								
Second shell	Lithium ${}_3\text{Li}$	Beryllium ${}_4\text{Be}$	Boron ${}_5\text{B}$	Carbon ${}_6\text{C}$	Nitrogen ${}_7\text{N}$	Oxygen ${}_8\text{O}$	Fluorine ${}_9\text{F}$	Neon ${}_{10}\text{Ne}$
								
Third shell	Sodium ${}_{11}\text{Na}$	Magnesium ${}_{12}\text{Mg}$	Aluminum ${}_{13}\text{Al}$	Silicon ${}_{14}\text{Si}$	Phosphorus ${}_{15}\text{P}$	Sulfur ${}_{16}\text{S}$	Chlorine ${}_{17}\text{Cl}$	Argon ${}_{18}\text{Ar}$
								

▲ Figure 2.9 Electron distribution diagrams for the first 18 elements in the periodic table. In a standard periodic table (see Appendix B), information for each element is presented as shown for helium in the inset. In the diagrams in this table, electrons are represented as yellow dots and electron

shells as concentric circles. These diagrams are a convenient way to picture the distribution of an atom's electrons among its electron shells, but these simplified models do not accurately represent the shape of the atom or the location of its electrons. The elements are arranged in rows, each representing the filling of an

electron shell. As electrons are added, they occupy the lowest available shell.

? What is the atomic number of magnesium? How many protons and electrons does it have? How many electron shells? How many valence electrons?

Electron Orbitals

In the early 1900s, the electron shells of an atom were visualized as concentric paths of electrons orbiting the nucleus, somewhat like planets orbiting the sun. It is still convenient to use two-dimensional concentric-circle diagrams, as in Figure 2.9, to symbolize three-dimensional electron

shells. However, you need to remember that each concentric circle represents only the *average* distance between an electron in that shell and the nucleus. Accordingly, the concentric-circle diagrams do not give a real picture of an atom. In reality, we can never know the exact location of an electron. What we can do instead is describe the space in which an electron spends most of its time. The three-dimensional space where an electron is found 90% of the time is called an **orbital**.

Each electron shell contains electrons at a particular energy level, distributed among a specific number of orbitals of distinctive shapes and orientations. **Figure 2.10** shows the orbitals of neon as an example, with its electron distribution diagram for reference. You can think of an orbital as a component of an electron shell. The first electron shell has only one spherical *s* orbital (called *1s*), but the second shell has four orbitals: one large spherical *s* orbital (called *2s*) and three dumbbell-shaped *p* orbitals (called *2p* orbitals). (The third shell and other higher electron shells also have *s* and *p* orbitals, as well as orbitals of more complex shapes.)


No more than 2 electrons can occupy a single orbital. The first electron shell can therefore accommodate up to 2 electrons in its *s* orbital. The lone electron of a hydrogen atom occupies the *1s* orbital, as do the 2 electrons of a helium atom. The four orbitals of the second electron shell can hold up to 8 electrons, 2 in each orbital. Electrons in each of the four orbitals have nearly the same energy, but they move in different volumes of space.

The reactivity of atoms arises from the presence of unpaired electrons in one or more orbitals of their valence shells. As you will see in the next section, atoms interact in a way that completes their valence shells. When they do so, it is the *unpaired* electrons that are involved.

CONCEPT CHECK 2.2

1. A lithium atom has 3 protons and 4 neutrons. What is its atomic mass in daltons?
2. A nitrogen atom has 7 protons, and the most common isotope of nitrogen has 7 neutrons. A radioactive isotope of nitrogen has 8 neutrons. Write the atomic number and mass number of this radioactive nitrogen as a chemical symbol with a subscript and superscript.
3. How many electrons does fluorine have? How many electron shells? Name the orbitals that are occupied. How many electrons are needed to fill the valence shell?
4. **WHAT IF?** In Figure 2.9, if two or more elements are in the same row, what do they have in common? If two or more elements are in the same column, what do they have in common?

For suggested answers, see Appendix A.



▲ **Figure 2.10** Electron orbitals.


CONCEPT 2.3

The formation and function of molecules depend on chemical bonding between atoms

Now that we have looked at the structure of atoms, we can move up the hierarchy of organization and see how atoms combine to form molecules and ionic compounds. Atoms with incomplete valence shells can interact with certain other atoms in such a way that each partner completes its valence shell: The atoms either share or transfer valence electrons. These interactions usually result in atoms staying close together, held by attractions called **chemical bonds**. The strongest kinds of chemical bonds are covalent bonds and ionic bonds.

Covalent Bonds

A **covalent bond** is the sharing of a pair of valence electrons by two atoms. For example, let's consider what happens when two hydrogen atoms approach each other. Recall that hydrogen has 1 valence electron in the first shell, but the shell's capacity is 2 electrons. When the two hydrogen atoms come close enough for their 1s orbitals to overlap, they can share their electrons (**Figure 2.11**). Each hydrogen atom is now associated with 2 electrons in what amounts



▲ **Figure 2.11** Formation of a covalent bond.

to a completed valence shell. Two or more atoms held together by covalent bonds constitute a **molecule**, in this case a hydrogen molecule.

Figure 2.12a shows several ways of representing a hydrogen molecule. Its *molecular formula*, H_2 , simply indicates that the molecule consists of two atoms of hydrogen. Electron sharing can be depicted by an electron distribution diagram or by a *Lewis dot structure*, in which element symbols are surrounded by dots that represent the valence electrons ($H:H$). We can also use a *structural formula*, $H-H$, where the line represents a **single bond**, a pair of shared electrons. A space-filling model comes closest to representing the actual shape of the molecule.

Oxygen has 6 electrons in its second electron shell and therefore needs 2 more electrons to complete its valence shell. Two oxygen atoms form a molecule by sharing *two* pairs of valence electrons (**Figure 2.12b**). The atoms are thus joined by a **double bond** ($O=O$).

Name and Molecular Formula	Electron Distribution Diagram	Lewis Dot Structure and Structural Formula	Space-Filling Model
(a) Hydrogen (H_2). Two hydrogen atoms share one pair of electrons, forming a single bond.		$H:H$ $H-H$	
(b) Oxygen (O_2). Two oxygen atoms share two pairs of electrons, forming a double bond.		$\ddot{O}::\ddot{O}$ $O=O$	
(c) Water (H_2O). Two hydrogen atoms and one oxygen atom are joined by single bonds, forming a molecule of water.		$\begin{array}{c} \ddot{O}:H \\ \\ H \end{array}$ $O-H$	
(d) Methane (CH_4). Four hydrogen atoms can satisfy the valence of one carbon atom, forming methane.		$\begin{array}{c} H \\ \vdots \\ H:C:H \\ \vdots \\ H \end{array}$ $\begin{array}{c} H \\ \\ H-C-H \\ \\ H \end{array}$	

▲ **Figure 2.12** Covalent bonding in four molecules. The number of electrons required to complete an atom's valence shell generally determines how many covalent bonds that atom will form. This figure shows several ways of indicating covalent bonds.


Each atom that can share valence electrons has a bonding capacity corresponding to the number of covalent bonds the atom can form. When the bonds form, they give the atom a full complement of electrons in the valence shell. The bonding capacity of oxygen, for example, is 2. This bonding capacity is called the atom's **valence** and usually equals the number of unpaired electrons required to complete the atom's outermost (valence) shell. See if you can determine the valences of hydrogen, oxygen, nitrogen, and carbon by studying the electron distribution diagrams in Figure 2.9. You can see that the valence of hydrogen is 1; oxygen, 2; nitrogen, 3; and carbon, 4. However, the situation is more complicated for elements in the third row of the periodic table. Phosphorus, for example, can have a valence of 3, as we would predict from the presence of 3 unpaired electrons in its valence shell. In some molecules that are biologically important, however, phosphorus can form three single bonds and one double bond. Therefore, it can also have a valence of 5.

The molecules H_2 and O_2 are pure elements rather than compounds because a compound is a combination of two or more *different* elements. Water, with the molecular formula H_2O , is a compound. Two atoms of hydrogen are needed to satisfy the valence of one oxygen atom. **Figure 2.12c** shows the structure of a water molecule. Water is so important to life that Chapter 3 is devoted entirely to its structure and behavior.

Methane, the main component of natural gas, is a compound with the molecular formula CH_4 . It takes four hydrogen atoms, each with a valence of 1, to complement one atom of carbon, with its valence of 4 (**Figure 2.12d**). We will look at many other compounds of carbon in Chapter 4.

Atoms in a molecule attract shared electrons to varying degrees, depending on the element. The attraction of a particular atom for the electrons of a covalent bond is called its **electronegativity**. The more electronegative an atom is, the more strongly it pulls shared electrons toward itself. In a covalent bond between two atoms of the same element, the electrons are shared equally because the two atoms have the same electronegativity—the tug-of-war is at a standoff. Such a bond is called a **nonpolar covalent bond**. For example, the single bond of H_2 is nonpolar, as is the double bond of O_2 . However, when one atom is bonded to a more electronegative atom, the electrons of the bond are not shared equally. This type of bond is called a **polar covalent bond**. Such bonds vary in their polarity, depending on the relative electronegativity of the two atoms. For example, the bonds between the oxygen and hydrogen atoms of a water molecule are quite polar (**Figure 2.13**).

Because oxygen (O) is more electronegative than hydrogen (H), shared electrons are pulled more toward oxygen.




▲ **Figure 2.13** Polar covalent bonds in a water molecule.

Oxygen is one of the most electronegative of all the elements, attracting shared electrons much more strongly than hydrogen does. In a covalent bond between oxygen and hydrogen, the electrons spend more time near the oxygen nucleus than they do near the hydrogen nucleus. Because electrons have a negative charge and are pulled toward oxygen in a water molecule, the oxygen atom has a partial negative charge (indicated by the Greek letter δ with a minus sign, δ^- , or “delta minus”), and each hydrogen atom has a partial positive charge (δ^+ , or “delta plus”). In contrast, the individual bonds of methane (CH_4) are much less polar because the electronegativities of carbon and hydrogen are similar.


Ionic Bonds

In some cases, two atoms are so unequal in their attraction for valence electrons that the more electronegative atom strips an electron completely away from its partner. This is what happens when an atom of sodium ($_{11}\text{Na}$) encounters an atom of chlorine ($_{17}\text{Cl}$) (**Figure 2.14**). A sodium atom has a total of 11 electrons, with its single valence electron in the third electron shell. A chlorine atom has a total of 17 electrons,

1 The lone valence electron of a sodium atom is transferred to join the 7 valence electrons of a chlorine atom.



2 Each resulting ion has a completed valence shell. An ionic bond can form between the oppositely charged ions.



▲ **Figure 2.14** Electron transfer and ionic bonding. The attraction between oppositely charged atoms, or ions, is an ionic bond. An ionic bond can form between any two oppositely charged ions, even if they have not been formed by transfer of an electron from one to the other.

with 7 electrons in its valence shell. When these two atoms meet, the lone valence electron of sodium is transferred to the chlorine atom, and both atoms end up with their valence shells complete. (Because sodium no longer has an electron in the third shell, the second shell is now the valence shell.)

The electron transfer between the two atoms moves one unit of negative charge from sodium to chlorine. Sodium, now with 11 protons but only 10 electrons, has a net electrical charge of $1+$. A charged atom (or molecule) is called an **ion**. When the charge is positive, the ion is specifically called a **cation**; the sodium atom has become a cation. Conversely, the chlorine atom, having gained an extra electron, now has 17 protons and 18 electrons, giving it a net electrical charge of $1-$. It has become a chloride ion—an **anion**, or negatively charged ion. Because of their opposite charges, cations and anions attract each other; this attraction is called an **ionic bond**. The transfer of an electron is not the formation of a bond; rather, it allows a bond to form because it results in two ions of opposite charge. Any two ions of opposite charge can form an ionic bond. The ions do not need to have acquired their charge by an electron transfer with each other.

Compounds formed by ionic bonds are called **ionic compounds**, or **salts**. We know the ionic compound sodium chloride (NaCl) as table salt (**Figure 2.15**). Salts are often found in nature as crystals of various sizes and shapes. Each salt crystal is an aggregate of vast numbers of cations and anions bonded by their electrical attraction and arranged in a three-dimensional lattice. Unlike a covalent compound, which consists of molecules having a definite size and number of atoms, an ionic compound does not consist of molecules. The formula for an ionic compound, such as NaCl , indicates only the ratio of elements in a crystal of the salt. “ NaCl ” by itself is not a molecule.

Not all salts have equal numbers of cations and anions. For example, the ionic compound magnesium chloride (MgCl_2) has two chloride ions for each magnesium ion. Magnesium ($_{12}\text{Mg}$) must lose 2 outer electrons if the atom is to have a

complete valence shell, so it tends to become a cation with a net charge of $2+$ (Mg^{2+}). One magnesium cation can therefore form ionic bonds with two chloride anions.

The term *ion* also applies to entire molecules that are electrically charged. In the salt ammonium chloride (NH_4Cl), for instance, the anion is a single chloride ion (Cl^-), but the cation is ammonium (NH_4^+), a nitrogen atom with four covalently bonded hydrogen atoms. The whole ammonium ion has an electrical charge of $1+$ because it is 1 electron short.

Environment affects the strength of ionic bonds. In a dry salt crystal, the bonds are so strong that it takes a hammer and chisel to break enough of them to crack the crystal in two. If the same salt crystal is dissolved in water, however, the ionic bonds are much weaker because each ion is partially shielded by its interactions with water molecules. Most drugs are manufactured as salts because they are quite stable when dry but can dissociate (come apart) easily in water. In the next chapter, you will learn how water dissolves salts.

Weak Chemical Bonds

In organisms, most of the strongest chemical bonds are covalent bonds, which link atoms to form a cell's molecules. But weaker bonding within and between molecules is also indispensable in the cell, contributing greatly to the emergent properties of life. Many large biological molecules are held in their functional form by weak bonds. In addition, when two molecules in the cell make contact, they may adhere temporarily by weak bonds. The reversibility of weak bonding can be an advantage: Two molecules can come together, respond to one another in some way, and then separate.


Several types of weak chemical bonds are important in organisms. One is the ionic bond as it exists between ions dissociated in water, which we just discussed. Hydrogen bonds and van der Waals interactions are also crucial to life.

Hydrogen Bonds


Among the various kinds of weak chemical bonds, hydrogen bonds are so important in the chemistry of life that they deserve special attention. The partial positive charge on a hydrogen atom that is covalently bonded to an electronegative atom allows the hydrogen to be attracted to a different electronegative atom nearby. This noncovalent attraction between a hydrogen and an electronegative atom is called a **hydrogen bond**. In living cells, the electronegative partners are usually oxygen or nitrogen atoms. Refer to **Figure 2.16** to examine the simple case of hydrogen bonding between water (H_2O) and ammonia (NH_3).

Van der Waals Interactions

Even a molecule with nonpolar covalent bonds may have positively and negatively charged regions. Electrons are not always symmetrically distributed in such a molecule; at any



▲ **Figure 2.15 A sodium chloride (NaCl) crystal.** The sodium ions (Na^+) and chloride ions (Cl^-) are held together by ionic bonds. The formula NaCl tells us that the ratio of Na^+ to Cl^- is $1:1$.



▲ **Figure 2.16 A hydrogen bond.**


DRAW IT Draw five water molecules using structural formulas and indicating partial charges, and show how they can make hydrogen bonds with each other.

instant, they may accumulate by chance in one part of the molecule or another. The results are ever-changing regions of positive and negative charge that enable all atoms and molecules to stick to one another. These **van der Waals interactions** are individually weak and occur only when atoms and molecules are very close together. When many such interactions occur simultaneously, however, they can be powerful: Van der Waals interactions are the reason a gecko lizard (right) can walk straight up a wall! Each gecko toe has hundreds of thousands of tiny hairs, with multiple projections at each hair's tip that increase surface area. Apparently, the van der Waals interactions between the hair tip molecules and the molecules of the wall's surface are so numerous that despite their individual weakness, together they can support the gecko's body weight.

Van der Waals interactions, hydrogen bonds, ionic bonds in water, and other weak bonds may form not only between molecules but also between parts of a large molecule, such as a protein. The cumulative effect of weak bonds is to reinforce the three-dimensional shape of the molecule. You will learn more about the very important biological roles of weak bonds in Chapter 5.

Molecular Shape and Function

A molecule has a characteristic size and shape. The precise shape of a molecule is usually very important to its function in the living cell.



(a) **Hybridization of orbitals.** The single s and three p orbitals of a valence shell involved in covalent bonding combine to form four teardrop-shaped hybrid orbitals. These orbitals extend to the four corners of an imaginary tetrahedron (outlined in pink).

Space-Filling Model	Ball-and-Stick Model	Hybrid-Orbital Model (with ball-and-stick model superimposed)
Water (H₂O)		
Methane (CH₄)		

(b) **Molecular-shape models.** Three models representing molecular shape are shown for water and methane. The positions of the hybrid orbitals determine the shapes of the molecules.

▲ **Figure 2.17 Molecular shapes due to hybrid orbitals.**

A molecule consisting of two atoms, such as H_2 or O_2 , is always linear, but most molecules with more than two atoms have more complicated shapes. These shapes are determined by the positions of the atoms' orbitals. When an atom forms covalent bonds, the orbitals in its valence shell undergo rearrangement. For atoms with valence electrons in both s and p orbitals (review Figure 2.10), the single s and three p orbitals form four new hybrid orbitals shaped like identical teardrops extending from the region of the atomic nucleus (**Figure 2.17a**). If we connect the larger ends of the teardrops with lines, we have the outline of a geometric shape called a tetrahedron, a pyramid with a triangular base.

For the water molecule (H_2O), two of the hybrid orbitals in the oxygen atom's valence shell are shared with hydrogen atoms (**Figure 2.17b**). The result is a molecule shaped roughly like a V, with its two covalent bonds spread apart at an angle of 104.5° .

The methane molecule (CH_4) has the shape of a completed tetrahedron because all four hybrid orbitals of the carbon atom are shared with hydrogen atoms (see Figure 2.17b). The carbon nucleus is at the center, with its four covalent bonds radiating to hydrogen nuclei at the corners of the tetrahedron. Larger molecules containing multiple carbon atoms, including many of the molecules that make up living matter, have more complex overall shapes. However, the tetrahedral shape of a carbon atom bonded to four other atoms is often a repeating motif within such molecules.

Molecular shape is crucial in biology because it determines how biological molecules recognize and respond to one another with specificity. Biological molecules often bind temporarily to each other by forming weak bonds, but this can happen only if their shapes are complementary. We can see this specificity in the effects of opiates, drugs derived from

opium. Opiates, such as morphine and heroin, relieve pain and alter mood by weakly binding to specific receptor molecules on the surfaces of brain cells. Why would brain cells carry receptors for opiates, compounds that are not made by our bodies? The discovery of endorphins in 1975 answered this question. Endorphins are signaling molecules made by the pituitary gland that bind to the receptors, relieving pain and producing euphoria during times of stress, such as intense exercise. It turns out that opiates have shapes similar to endorphins and mimic them by binding to endorphin receptors in the brain. That is why opiates (such as morphine) and endorphins have similar effects (Figure 2.18). The role of molecular shape in brain chemistry illustrates the relationship between structure and function, one of biology's unifying themes.

CONCEPT CHECK 2.3


1. Why does the structure $\text{H}-\text{C}=\text{C}-\text{H}$ fail to make sense chemically?
2. What holds the atoms together in a crystal of magnesium chloride (MgCl_2)?
3. **WHAT IF?** If you were a pharmaceutical researcher, why would you want to learn the three-dimensional shapes of naturally occurring signaling molecules?

For suggested answers, see Appendix A.


CONCEPT 2.4

Chemical reactions make and break chemical bonds


The making and breaking of chemical bonds, leading to changes in the composition of matter, are called **chemical reactions**. An example is the reaction between hydrogen and oxygen molecules that forms water:




This reaction breaks the covalent bonds of H_2 and O_2 and forms the new bonds of H_2O . When we write a chemical reaction, we use an arrow to indicate the conversion of the starting materials, called the **reactants**, to the **products**. The coefficients indicate the number of molecules involved; for example, the coefficient 2 in front of the H_2 means that




Natural endorphin



Morphine




(a) Structures of endorphin and morphine. The boxed portion of the endorphin molecule (left) binds to receptor molecules on target cells in the brain. The boxed portion of the morphine molecule (right) is a close match.



(b) Binding to endorphin receptors. Both endorphin and morphine can bind to endorphin receptors on the surface of a brain cell.

▲ **Figure 2.18 A molecular mimic.** Morphine affects pain perception and emotional state by mimicking the brain's natural endorphins.




▲ Figure 2.19 Photosynthesis: a solar-powered rearrangement of matter. *Elodea*, a freshwater plant, produces sugar by rearranging the atoms of carbon dioxide and water in the chemical process known as photosynthesis, which is powered by sunlight. Much of the sugar is then converted to other food molecules. Oxygen gas (O_2) is a by-product of photosynthesis; notice the bubbles of oxygen escaping from the leaves in the photo.

? Explain how this photo relates to the reactants and products in the equation for photosynthesis given in the text. (You will learn more about photosynthesis in Chapter 10.)


the reaction starts with two molecules of hydrogen. Notice that all atoms of the reactants must be accounted for in the products. Matter is conserved in a chemical reaction: Reactions cannot create or destroy matter but can only rearrange it.

Photosynthesis, which takes place within the cells of green plant tissues, is a particularly important example of how chemical reactions rearrange matter. Humans and other animals ultimately depend on photosynthesis for food and oxygen, and this process is at the foundation of almost all ecosystems. The following chemical shorthand summarizes the process of photosynthesis:



The raw materials of photosynthesis are carbon dioxide (CO_2), which is taken from the air, and water (H_2O), which is absorbed from the soil. Within the plant cells, sunlight powers the conversion of these ingredients to a sugar called glucose ($\text{C}_6\text{H}_{12}\text{O}_6$) and oxygen molecules (O_2), a by-product that the plant releases into the surroundings (Figure 2.19). Although photosynthesis is actually a sequence of many chemical reactions, we still end up with the same number and types of atoms that we had when we started. Matter has simply been rearranged, with an input of energy provided by sunlight.

All chemical reactions are reversible, with the products of the forward reaction becoming the reactants for the reverse reaction. For example, hydrogen and nitrogen molecules can combine to form ammonia, but ammonia can also decompose to regenerate hydrogen and nitrogen:



The two opposite-headed arrows indicate that the reaction is reversible.

One of the factors affecting the rate of a reaction is the concentration of reactants. The greater the concentration of reactant molecules, the more frequently they collide with one another and have an opportunity to react and form products. The same holds true for products. As products accumulate, collisions resulting in the reverse reaction become more frequent. Eventually, the forward and reverse reactions occur at the same rate, and the relative concentrations of products and reactants stop changing. The point at which the reactions offset one another exactly is called **chemical equilibrium**. This is a dynamic equilibrium; reactions are still going on, but with no net effect on the concentrations of reactants and products. Equilibrium does *not* mean that the reactants and products are equal in concentration, but only that their concentrations have stabilized at a particular ratio. The reaction involving ammonia reaches equilibrium when ammonia decomposes as rapidly as it forms. In some chemical reactions, the equilibrium point may lie so far to the right that these reactions go essentially to completion; that is, virtually all the reactants are converted to products.

We will return to the subject of chemical reactions after more detailed study of the various types of molecules that are important to life. In the next chapter, we focus on water, the substance in which all the chemical processes of organisms occur.

CONCEPT CHECK 2.4

- 1. MAKE CONNECTIONS** Consider the reaction between hydrogen and oxygen that forms water, shown with ball-and-stick models on page 42. Study Figure 2.12 and draw the Lewis dot structures representing this reaction.
- 2.** Which type of chemical reaction occurs faster at equilibrium, the formation of products from reactants or reactants from products?
- 3. WHAT IF?** Write an equation that uses the products of photosynthesis as reactants and the reactants of photosynthesis as products. Add energy as another product. This new equation describes a process that occurs in your cells. Describe this equation in words. How does this equation relate to breathing?

For suggested answers, see Appendix A.

2 CHAPTER REVIEW

SUMMARY OF KEY CONCEPTS

CONCEPT 2.1

Matter consists of chemical elements in pure form and in combinations called compounds (pp. 31–32)


- **Elements** cannot be broken down chemically to other substances. A **compound** contains two or more different elements in a fixed ratio. Oxygen, carbon, hydrogen, and nitrogen make up approximately 96% of living matter.

? In what way does the need for iodine or iron in your diet differ from your need for calcium or phosphorus?

CONCEPT 2.2

An element's properties depend on the structure of its atoms (pp. 33–37)

- An **atom**, the smallest unit of an element, has the following components:



- An electrically neutral atom has equal numbers of electrons and protons; the number of protons determines the **atomic number**. The **atomic mass** is measured in **daltons** and is roughly equal to the sum of protons plus neutrons. **Isotopes** of an element differ from each other in neutron number and therefore mass. Unstable isotopes give off particles and energy as radioactivity.
- In an atom, electrons occupy specific **electron shells**; the electrons in a shell have a characteristic energy level. Electron distribution in shells determines the chemical behavior of an atom. An atom that has an incomplete outer shell, the **valence shell**, is reactive.
- Electrons exist in **orbitals**, three-dimensional spaces with specific shapes that are components of electron shells.




DRAW IT Draw the electron distribution diagrams for neon ($_{10}\text{Ne}$) and argon ($_{18}\text{Ar}$). Use these diagrams to explain why these elements are chemically unreactive.


CONCEPT 2.3

The formation and function of molecules depend on chemical bonding between atoms (pp. 38–42)

- **Chemical bonds** form when atoms interact and complete their valence shells. **Covalent bonds** form when pairs of electrons are shared.



- **Molecules** consist of two or more covalently bonded atoms. The attraction of an atom for the electrons of a covalent bond is its **electronegativity**. If both atoms are the same, they have the same electronegativity and share a **nonpolar covalent bond**. Electrons of a **polar covalent bond** are pulled closer to the more electronegative atom.
- An **ion** forms when an atom or molecule gains or loses an electron and becomes charged. An **ionic bond** is the attraction between two oppositely charged ions.



- Weak bonds reinforce the shapes of large molecules and help molecules adhere to each other. A **hydrogen bond** is an attraction between a hydrogen atom carrying a partial positive charge (δ^+) and an electronegative atom (δ^-). **Van der Waals interactions** occur between transiently positive and negative regions of molecules.
- A molecule's shape is determined by the positions of its atoms' valence orbitals. Covalent bonds result in hybrid orbitals, which are responsible for the shapes of H_2O , CH_4 , and many more complex biological molecules. Shape is usually the basis for the recognition of one biological molecule by another.

? In terms of electron sharing between atoms, compare nonpolar covalent bonds, polar covalent bonds, and the formation of ions.

CONCEPT 2.4

Chemical reactions make and break chemical bonds (pp. 42–43)

- **Chemical reactions** change **reactants** into **products** while conserving matter. All chemical reactions are theoretically reversible. **Chemical equilibrium** is reached when the forward and reverse reaction rates are equal.

? What would happen to the concentration of products if more reactants were added to a reaction that was in chemical equilibrium? How would this addition affect the equilibrium?

TEST YOUR UNDERSTANDING

LEVEL 1: KNOWLEDGE/COMPREHENSION

- In the term *trace element*, the adjective *trace* means that
 - the element is required in very small amounts.
 - the element can be used as a label to trace atoms through an organism's metabolism.
 - the element is very rare on Earth.
 - the element enhances health but is not essential for the organism's long-term survival.
 - the element passes rapidly through the organism.

- Compared with ^{31}P , the radioactive isotope ^{32}P has
 - a different atomic number.
 - a different charge.
 - one more proton.
 - one more electron.
 - one more neutron.
- The reactivity of an atom arises from
 - the average distance of the outermost electron shell from the nucleus.
 - the existence of unpaired electrons in the valence shell.
 - the sum of the potential energies of all the electron shells.
 - the potential energy of the valence shell.
 - the energy difference between the s and p orbitals.
- Which statement is true of all atoms that are anions?
 - The atom has more electrons than protons.
 - The atom has more protons than electrons.
 - The atom has fewer protons than does a neutral atom of the same element.
 - The atom has more neutrons than protons.
 - The net charge is $1-$.
- Which of the following statements correctly describes any chemical reaction that has reached equilibrium?
 - The concentrations of products and reactants are equal.
 - The reaction is now irreversible.
 - Both forward and reverse reactions have halted.
 - The rates of the forward and reverse reactions are equal.
 - No reactants remain.

LEVEL 2: APPLICATION/ANALYSIS

- We can represent atoms by listing the number of protons, neutrons, and electrons—for example, $2p^+$, $2n^0$, $2e^-$ for helium. Which of the following represents the ^{18}O isotope of oxygen?
 - $6p^+$, $8n^0$, $6e^-$
 - $8p^+$, $10n^0$, $8e^-$
 - $9p^+$, $9n^0$, $9e^-$
 - $7p^+$, $2n^0$, $9e^-$
 - $10p^+$, $8n^0$, $9e^-$
- The atomic number of sulfur is 16. Sulfur combines with hydrogen by covalent bonding to form a compound, hydrogen sulfide. Based on the number of valence electrons in a sulfur atom, predict the molecular formula of the compound.
 - HS
 - HS₂
 - H₂S
 - H₃S₂
 - H₄S
- What coefficients must be placed in the following blanks so that all atoms are accounted for in the products?

$$\text{C}_6\text{H}_{12}\text{O}_6 \rightarrow \text{C}_2\text{H}_6\text{O} + \text{CO}_2$$
 - 1; 2
 - 3; 1
 - 1; 3
 - 1; 1
 - 2; 2
- DRAW IT** Draw Lewis dot structures for each hypothetical molecule shown below, using the correct number of valence electrons for each atom. Determine which molecule makes sense because each atom has a complete valence shell and each bond has the correct number of electrons. Explain what makes the other molecules nonsensical, considering the number of bonds each type of atom can make.
 - O=C—H
 - $$\begin{array}{c} \text{H} \quad \text{H} \\ | \quad | \\ \text{H}-\text{C}-\text{H}-\text{C}=\text{O} \\ | \\ \text{H} \end{array}$$
 - $$\begin{array}{c} \text{H} \quad \text{H} \\ | \quad | \\ \text{H}-\text{O}-\text{C}-\text{C}=\text{O} \\ | \\ \text{H} \end{array}$$
 - $$\begin{array}{c} \text{O} \\ | \\ \text{H}-\text{N}=\text{H} \end{array}$$

LEVEL 3: SYNTHESIS/EVALUATION

10. EVOLUTION CONNECTION

The percentages of naturally occurring elements making up the human body (see Table 2.1) are similar to the percentages of these elements found in other organisms. How could you account for this similarity among organisms?

11. SCIENTIFIC INQUIRY

Female silkworm moths (*Bombyx mori*) attract males by emitting chemical signals that spread through the air. A male hundreds of meters away can detect these molecules and fly toward their source. The sensory organs responsible for this behavior are the comblike antennae visible in the photograph shown here. Each filament of an antenna is equipped with thousands of receptor cells that detect the sex attractant. Based on what you learned in this chapter, propose a hypothesis to account for the ability of the male moth to detect a specific molecule in the presence of many other molecules in the air. What predictions does your hypothesis make? Design an experiment to test one of these predictions.



12. WRITE ABOUT A THEME

Emergent Properties While waiting at an airport, Neil Campbell once overheard this claim: “It’s paranoid and ignorant to worry about industry or agriculture contaminating the environment with their chemical wastes. After all, this stuff is just made of the same atoms that were already present in our environment.” Drawing on your knowledge of electron distribution, bonding, and the theme of emergent properties (pp. 3–5), write a short essay (100–150 words) countering this argument.

For selected answers, see Appendix A.



www.masteringbiology.com

1. MasteringBiology® Assignments

Tutorials The Anatomy of Atoms • Atomic Number and Mass Number

Activities Structure of the Atomic Nucleus • Electron Arrangement • Covalent Bonds • Nonpolar and Polar Molecules • Ionic Bonds • Hydrogen Bonds

Questions Student Misconceptions • Reading Quiz • Multiple Choice • End-of-Chapter

2. eText

Read your book online, search, take notes, highlight text, and more.

3. The Study Area

Practice Tests • Cumulative Test • **BioFlix** 3-D Animations • MP3 Tutor Sessions • Videos • Activities • Investigations • Lab Media • Audio Glossary • Word Study Tools • Art