## **Tube Feeding Authorization for School**

School District/School	Fax Number	
This form provides health care provider and pare the prescribing health care provider and the par provided. Any alteration of the form invalidates t	ent/legal guardian are required to complete	
<b>Note:</b> Physician's orders are required for <b>all medica</b> portion of the form and return it to the school or have	•	
The following section is to be completed by the The student named in this document is under m the following treatment that is necessary to be g the prescribed treatment may be administered i	ny medical supervision for the diagnosis de given during school hours for the child's he	scribed below. I have prescribed
Diagnosis for which tube feeding will be required in	school:	
Type of appliance placed: Peg/ Long G-Tube L	_ow Profile G-Tube Button GJ-Tube J-Tube	e Other, describe:
Type of formula:		Venting instructions:
Time(s) of tube feeding syringe water flush:		Amount of water flush:
Time(s) of formula feedings and amount: (If homema	ade blended formula, please specify minimum a	nd maximum per feed)
Tube feeding method: G-Tube J-Tube Gravit Mechanical pump – Type of pump	ty bag bolus Syringe bolus Pump-assisted Rate of	

Date of Birth

Date

Frequency:

Is student allowed oral feedings?	Yes	No If yes, Type:

## Side effects to be reported:

Student's Name

## **Stoma Preservation Plan**

Prompt attention is important if a gastrointestinal feeding tube becomes dislodged. The tract can narrow or close in less then one (1) hour if it is not preserved. Do not use the G-tube or Foley catheter for feedings or medications until placement is verified by the parent. If stoma < 6 weeks old, placement must be verified by provider.

School nurse or trained personnel will preserve the stoma:

- Using a G-tube: use new or dislodged balloon G-tube (Mic-Key) if available and undamaged. Deflate balloon, lubricate shaft with
  water-soluble lubricant if available (such as Surgilube) and insert into gastrostomy site. DO NOT INFLATE THE BALLOON. Secure in
  place with medical tape.
- Using a Foley Catheter: Use Foley catheter of the same diameter (French) or one size smaller than patient's dislodged Gtube. Lubricate the shaft with water-soluble lubricant if available and insert approximately 2-4 inches into gastrostomy site. DO NOT INFLATE THE BALLOON. Secure in place with medical tape.

 Provider's Name (Print):
 Phone Number:

 Provider's Signature:
 Date:

 Fax Number:
 Date:

## Parent/Legal Guardian Consent

The following section is to be completed by the parent/legal guardian: I authorize this procedure to be performed by the school nurse or the nurse's delegatee as directed above. I agree to provide the needed supplies for the procedure and understand that new forms must be completed annually or with any changes in the student's health status. By signing this document, I give permission for the nurse or nurse designee to administer this procedure as prescribed and give my permission for this Health Care Provider to share information about this procedure with the Registered Nurse or nurse designee. The undersigned parent(s) or guardian(s) hereby agree(s) to exempt and release the school district and its directors, officers, employees, volunteers and agents, from any and all liability, claims, demands or actions whatsoever arising out of any damage, loss or injury that my child or I/we might sustain or which they now have or may hereafter have arising out of the administration of this procedure. Should the G-Tube become dislodged, I will be immediately contacted by school personnel.				
Parent/Guardian Name:		Relationship:		
Home phone:	Business phone:	Emergency phone:		
Parent/Guardian Signature:		Date:		

Revised 1/2018