COLORADO ASTHMA CARE PLAN AND MEDICATION ORDER FOR SCHOOL AND CHILD CARE SETTINGS*

PARENT/GUARDIAN COMPLETE, SIGN AND DATE:			
Child Name:			Birthdate:
School:			
Parent/Guardian Name:			Phone:
I approve this care plan and give permission for school personnel to share this information, follow this plan, administer medication and care for my child/youth, and if necessary, contact our health care provider. I assume responsibility for providing the school/program prescribed, non-expired medication and supplies (such as a spacer), and to comply with board policies, if applicable. I am aware 911 may be called if a quick relief inhaler is not at school and my child/youth is experiencing symptoms.			
Parent/Guardian Signature			Date
HEALTH CARE PROVIDER COMPLETE ALL ITEMS, SIGN AND DATE:			
QUICK RELIEF MEDICATION: ☐ Albuterol ☐ Other:			
Controller medication used at home:			
TRIGGERS: ☐ Weather ☐ Illness ☐ Exercise ☐ Smoke ☐ Dust ☐ Pollen ☐ Poor Air Quality ☐ Other: ☐ Life threatening allergy specify:			
QUICK RELIEF INHALER ADMINISTRATION: With assistance or self-carry.			
Student needs supervision or assistance to use inhaler. Student will not self-carry inhaler.			
☐ Student understands proper use of asthma medications, and in my opinion, can self-carry and use his/her inhaler at			
school independently with approval from school nurse and completion of contract.			
IF YOU SEE THIS: DO THIS:			
GREEN ZONE: No Symptoms Pretreat	 No current symptoms Strenuous activity planned PRETREATMENT FOR STRENUOUS ACTIVITY, please choose ONE: □ Not required OR □ Student/Parent request OR □ Routinely Give QUICK RELIEF MED 10-15 minutes before activity: □ 2 puffs □ 4 puffs 		Parent request OR
	p	Repeat in 4 hours, if needed for additional physical activity. If child is currently experiencing symptoms, follow YELLOW or RED ZONE.	
YELLOW ZONE: Mild symptoms	Trouble breathing	1. Give QUICK RELIEF MED: □ 2 puffs □ 4 puffs	
	Wheezing	2. Stay with child/youth and maintain sitting position.	
	Frequent cough	3. REPEAT QUICK RELIEF MED if not improving in 15 minutes: ☐ 2 puffs ☐ 4 puffs	
	• Chest tightness	If symptoms do not improve or worsen, follow RED ZONE.	
	 Not able to do activities 	 Child/youth may go back to normal activities, once symptoms are relieved. Notify parents/guardians and school nurse. 	
5. Notify parents/guardians and school hurse.			
RED ZONE: EMERGENCY Severe Symptoms	 Coughs constantly Struggles to breathe Trouble talking (only speaks 3-5 words) Give QUICK RELIEF MED: □ 2 puffs □ 4 puffs Refer to the anaphylaxis care plan if the student has a life threatening allergy. there is no anaphylaxis care plan follow emergency guidelines for anaphylaxis Call 911 and inform EMS the reason for the call. 		
	• Skin of chest and/or neck	 3. REPEAT QUICK RELIEF MED if not improving: 2 puffs 4 puffs Can repeat every 5-15 minutes until EMS arrives. 4. Stay with child/youth. Remain calm, encouraging slower, deeper breaths. 	
	pull in with breathing		
Se	 Lips/fingernails gray/blue 		
5. Notify parents/guardians and school nurse.			
Health Care Provider Signature Print Provider Name Date			
Fax Phone		one	Email
School Nurse/CCHC Signature Date Self-carry contract on file. Anaphylaxis plan on file for life threatening allergy to:			

^{*}Including reactive airways, exercise-induced bronchospasm, twitchy airways.



Revised: February 2021