


Northwest Independent School District
Epinephrine Health Care Plan



EMERGENCY HEALTH CARE PLAN		CAMPUS	YEAR
Student's Name:		Grade:	Student's Weight:
Birthdate:	Allergic to:	Asthma: <input type="checkbox"/> Yes <input type="checkbox"/> No	

EMERGENCY TREATMENT / MILD Symptoms

If student experiences a few hives; itchy skin; swelling at site of insect sting OR if an ingestion (or sting) is *suspected* watch for progression to SEVERE symptoms.

Treatment:

- Send student to health clinic **ACCOMPANIED BY AN ADULT**
- Give _____ of _____ by mouth if ordered.
Amount Antihistamine
- Contact the PARENT or emergency contact person.
- Student should rest. MONITOR closely for improvement or worsening of symptoms until parent arrives. Prepare to give Epinephrine if needed.

EMERGENCY TREATMENT / SEVERE Allergic Reaction – CALL 911

SYMPTOMS: Mild symptoms may become SEVERE. Signs of a life threatening reaction may include:

- | | |
|----------|--|
| MOUTH | itching swelling of lips and/or tongue, metallic taste in mouth |
| THROAT | itching, tightness/closure, hoarseness |
| SKIN | itching, hives, redness, swelling (more than a localized reaction) |
| GUT | nausea, vomiting, diarrhea, cramps |
| LUNG | shortness of breath, cough, wheeze, trouble swallowing, nasal congestion/sneezing |
| HEART | weak pulse, dizziness, passing out, low blood pressure |
| BEHAVIOR | anxiety, difficulty talking/slurred speech, headache, confusion, "feeling of doom" |

TREATMENT:

****911 should always be called when an EPINEPHRINE AUTO-INJECTOR is given**

- Give EpiPen® Auto-injector immediately for severe symptoms or more than one mild symptom
- Call 911 (or local emergency response team) immediately. EpiPen® only lasts 20-30 minutes
- Contact parents or emergency contact person
- When parents are unavailable, school personnel should accompany the child to the hospital

Directions for use of EpiPen®:

- Remove from case - Pull off BLUE safety cap
- Place ORANGE tip against upper outer thigh
- Press hard into outer thigh, until it clicks
- HOLD in place 10 seconds, then remove
- Discard EpiPen® in "sharps container" or give to the emergency medical responder
(Do not return to holder)

Directions for use of Auvi-Q®

- Pull device out from case
- Follow VOICE PROMPTS for directions
- Press hard into outer thigh, until it clicks
- HOLD in place 5 seconds, then remove
- Discard Auvi-Q® in "sharps container", or give to the emergency medical responder
(Do not return to holder)

PHYSICIAN INSTRUCTIONS:

- | | | |
|-----|----|--|
| YES | NO | Student understands proper use and may carry their EpiPen®. |
| YES | NO | Epinephrine Auto-Injector in clinic (0.15 MG ____ 0.3 MG ____) – must be provided by parent |
| YES | NO | Antihistamine (Benadryl) in clinic - must be provided by parent |

Additional Instructions: _____

PHYSICIAN NAME:		SIGNATURE:	
Phone Number:	Fax Number :	Date:	


PARENT: I give consent with my signature for this information to be shared with Staff and the Healthcare provider.

PARENT/GUARDIAN NAME:		SIGNATURE:	
Phone Number(s):		Date:	


- Copy provided to Parent Copy sent to Healthcare Provider

EpiPen® (epinephrine) Auto-Injector Directions

- First, remove the EpiPen® (epinephrine) Auto-Injector from the plastic carrying case
- Pull off the blue safety release cap



- Hold orange tip near outer thigh (always apply to thigh)



- Swing and firmly push orange tip against outer thigh. Hold on thigh for approximately 10 seconds.


Remove EpiPen® (epinephrine) Auto-Injector and massage the area for 10 more seconds.



EpiPen®, EpiPen 2-Pak®, and EpiPen Jr 2-Pak® are registered trademarks of Mylan Inc. licensed exclusively to its wholly-owned subsidiary, Mylan Specialty L.P.


Auvi-Q™ (epinephrine injection, USP) Directions

Remove the outer case of Auvi-Q. This will automatically activate the voice instructions.



Pull off RED safety guard.

Place black end against outer thigh, then press firmly and hold for 5 seconds.



Auvi-Q™
epinephrine injection, USP
0.15 mg/0.3 mg auto-injectors

© 2002-2013 sanofi-aventis U.S. LLC. All rights reserved.

Adrenaclick® 0.3 mg and Adrenaclick® 0.15 mg Directions



Remove GREY caps labeled "1" and "2."



Place RED rounded tip against outer thigh, press down hard until needle penetrates. Hold for 10 seconds, then remove.