



FERTILIZER WITH CONFRONT* 3 & DIMENSION

19-0-6 WITH MESA

GUARANTEED ANALYSIS

Total Nitrogen (N) 19%
11.2% Ammoniacal Nitrogen
4.0% Water Insoluble Nitrogen*
0.8% Urea Nitrogen
3.0% Other Water Soluble Nitrogen*

Soluble Potash (K2O) 6%
Sulfur (S) 14.5%
14.5% Combined Sulfur (S)

Derived from: Ammonium Sulfate, Methylene Ureas, Urea, Muriate of Potash.
*7.0% Slowly Available Nitrogen from Methylene Urea

Chlorine (Cl) not more than 6%
Mesa®, PennMulch® and ProScape® are registered trademarks of Lebanon Seaboard Corporation.

Table with 2 columns: Spreader and Settings (3.5 lbs., 4 lbs., 4.5 lbs., 5 lbs.). Includes rows for LebanonTurf, Andersons AccuPro, Earthway Rotary, Lely, and Lesco.

For the postemergence control of annual and perennial broadleaf weeds, and preemergence control of annual grassy weeds and labeled broadleaf weeds, in established ornamental turf, lawns, golf courses, roadsides and cemeteries.

ACTIVE INGREDIENTS:

Table listing active ingredients: Dimethylamine salt of 2,4-D (0.819%), Triethylamine salt of triclopyr (0.303%), Methylheptyl ester of fluroxypyr (0.177%), Dithiopyr (0.145%).

OTHER INGREDIENTS: 98.556%
TOTAL: 100.000%

Table of Acid Equivalents: 2,4-dichlorophenoxyacetic acid (0.680% 13.60 lbs./ton), triclopyr (0.217% 4.34 lbs./ton), fluroxypyr (0.123% 2.46 lbs./ton), dithiopyr (0.145% 2.9 lbs./ton).

EPA Reg No. 961-406 (L) (D)
EPA Est. No. 961-PA-01: 961-IL-01
*Trademark of Dow AgroSciences LLC. Superscript used is first letter of lot number.

KEEP OUT OF REACH OF CHILDREN CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

READ THE ENTIRE LABEL BEFORE USING THIS PRODUCT. USE ONLY ACCORDING TO LABEL INSTRUCTIONS.

Before buying this product, read the Warranty Section elsewhere on this label.

If terms are not acceptable, return the unopened container to the place of purchase at once.

In New York State: This product may be applied only by commercial applicators at no more than 241 pounds (0.35 lb. of active ingredient of dithiopyr) per acre (5.5 lbs. per 1,000 sq.ft.) per year. Use of this product in Nassau and Suffolk Counties is prohibited.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if swallowed, inhaled or absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Avoid breathing dust. Prolonged or frequently repeated skin contact may cause allergic reaction in some individuals.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

All loaders, applicators, and other handlers must wear:

- Long-sleeved shirt and long pants,
Socks plus shoes.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS
Users should: Wash hands after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove clothing/PPE immediately if pesticide gets inside. Wash the outside of gloves before removing.

FIRST AID
IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
IF ON SKIN OR CLOTHING: Rinse skin immediately with plenty of water for 15-20 minutes.
IF SWALLOWED: Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor.
IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.

ENVIRONMENTAL HAZARDS
This pesticide may be toxic to fish and aquatic invertebrates. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. NOT FOR USE ON TURF BEING GROWN FOR SALE OR OTHER COMMERCIAL USE AS SOD, OR FOR COMMERCIAL SEED PRODUCTION OR FOR RESEARCH PURPOSES.

GENERAL INFORMATION

Lebanon ProScape® Fertilizer with Confront 3® & Dimension provides broad-spectrum postemergence control of actively growing annual and perennial broadleaf weeds, preemergence or early postemergence control of certain annual grassy weeds, and preemergence control or suppression of certain broadleaf weeds in established turf, including residential lawns, golf courses, sports fields, roadsides, cemeteries and around commercial buildings and other commercial and residential turf areas.

USE PRECAUTIONS AND RESTRICTIONS

Be sure that use of this product conforms to all applicable regulations. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Do not use on grass species such as bentgrass or St. Augustinegrass unless injury can be tolerated. Do not use this product on seedling buffalograss in the spring of establishment until the turfgrass is fully green and has established new roots.

Lebanon ProScape® Fertilizer with Confront® 3 & Dimension is recommended for use on the following established turfgrass species:

Table with 3 columns: Common Name, Scientific Name, and Established Cool/Warm Season Turf Grasses. Lists species like Bahiagrass, Bermudagrass, Buffalograss, etc.

(1) Use Lebanon ProScape® Fertilizer with Confront 3® & Dimension on this species only if some injury can be tolerated.

Established Lawns: Apply when broadleaf weeds are actively growing and prior to germination or tillering of annual grasses. Lebanon ProScape® Fertilizer with Confront® 3 & Dimension may be applied as a single application, as a split application or as a sequential application in the spring, summer or fall as described in the following use programs.

HOW TO APPLY

For optimum broadleaf weed control, apply to recently mowed turf when foliage is moist. When possible, delay mowing or irrigation for several days after treatment. The dithiopyr preemergent component in this product is not effective until activated by rainfall or irrigation.

APPLICATION RATES

Apply 152 to 218 lbs. per acre (3.5 to 5 lbs. per 1,000 sq. ft.) of Lebanon ProScape® Fertilizer with Confront 3 & Dimension to control actively growing broadleaf weeds and prevent germination of labeled annual grasses and broadleaf weeds growing in established turf.

Transition: DE, KS, KY, MD, MO, NJ, VA, southeastern PA, southeastern areas of IL, IN, OH, and coastal areas of CT & RI
South: AL, AR, GA, LA, MS, NC, SC, and TX
Coastal South: HI, FL, and southern coastal areas of AL, GA, LA, MS, NC, SC, and TX
West: AZ, CA, and NV

Supplemental applications of lawn fertilizer are needed to achieve a minimum 1 pound nitrogen per 1,000 sq. ft. POSTEMERGENCE & PREEMERGENCE PROGRAMS WITH APPLICATION RATES & WEEDS CONTROLLED

Table with 5 columns: Application Rate (152 LB./ACRE, 3.5 LBS./1,000Q. FT.), Fluroxypyr AE/Acre (0.187), Triclopyr AE/Acre (0.330), 2,4-D AE/Acre (1.034), Dithiopyr AI/Acre (0.2204).

Table listing weed species controlled by the product, such as Bedstraw, Bittercress, Black medic, etc.

Table with 5 columns: Application Rate (174 LB./ACRE, 4.0 LBS./1,000 SQ. FT.), Fluroxypyr AE/Acre (0.214), Triclopyr AE/Acre (0.378), 2,4-D AE/Acre (1.183), Dithiopyr AI/Acre (0.252).

Increases postemergent control to include the additional weeds listed below and provides 4-6 months of preemergent control on low cut turf** where weed control has been conducted during the previous year. Also provides early postemergent control of crabgrass up to the 3 leaf or first tiller stage on high cut turf*.

Table with 5 columns: Application Rate (196 LB./ACRE, 4.5 LBS./1,000 SQ. FT.), Fluroxypyr AE/Acre (0.241), Triclopyr AE/Acre (0.425), 2,4-D AE/Acre (1.333), Dithiopyr AI/Acre (0.284).

Improves postemergent control and provides 3-5 months of preemergent control on high cut turf* and low cut turf** where weed control has been conducted during the previous year.

Table with 5 columns: Application Rate (218 LB./ACRE, 5.0 LBS./1,000 SQ. FT.), Fluroxypyr AE/Acre (0.268), Triclopyr AE/Acre (0.473), 2,4-D AE/Acre (1.482), Dithiopyr AI/Acre (0.316).

Improves postemergent control and provides 4-6 months of preemergent control on low cut turf** where weed control has been conducted during the previous year in the coastal south and west. For coastal south split applications of 5.0 + 5.0 lbs are recommended.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. PESTICIDE STORAGE: Store this product only in its original container in a cool, dry, locked place out of the reach of children and animals. PESTICIDE DISPOSAL: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

WARRANTY

IMPORTANT: READ BEFORE USE

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once. By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

For technical assistance or more information about our products visit: www.LebanonTurf.com

Manufactured by: Lebanon Seaboard Corporation
1600 E. Cumberland St. • Lebanon, PA 17042
www.LebanonTurf.com

rev: 1019JC

Rev: KW-L/21

Net Weight: 40 LBS. (18.14 kg)