

School Guidelines for Transmittable Illnesses/Rashes.

****For any and all illnesses** (including common colds) – must be fever and vomit free for 24 hours prior to returning to school.

Strep Throat

- Must be fever free for 24 hours without the use of medication, have taken antibiotics at least 12 hours, and the child is feeling better.

Flu

- Must be fever and vomit free for 24 hours without the use of medication and the child is feeling better.

RSV

- Must be fever/vomit free for 24 hours without the use of medications and child is feeling better. It is recommended to remain home for 3-8 days (contagious period).

Covid-19

- Must be fever and vomit free for 24 hours without the use of medications and the child is feeling better. Recommend CDC guidelines (5 days at home from start of symptoms, then 5 days with a mask).

Mono

- Must be fever free for 24 hours without the use of medications and the child is feeling better. However, can be contagious for months.

Lice

- May return to school after one treatment with an over-the-counter lice-killing product and there are no longer any live lice seen on scalp. Will check scalp for periodically until 10 days after the first treatment.

Ringworm

- Rash on the body – May return after antifungal treatment is started and area can be covered.
- Rash on the scalp – May return after oral antifungal medication has been started and must have a doctor's excuse to return.

Impetigo

- May return 24 hours after treatment has started.

Hand, Foot, and Mouth

- Must be fever free for 24 hours without the use of medications and blisters are no longer open or have crusted over.

Chicken Pox

- Must stay home for at least 5 days and until all the blisters have crusted over.