Place Child's Picture Here

Food Allergy Action Plan

Student's Name:_____ **D.O.B:**_____ **Teacher:** _____

ALLERGY TO: _____

Asthmatic Yes* No *Higher risk for severe reaction

◆ STEP 1: TREATMENT ◆

Symptoms:		**Give Checked Medication**:**
▪	If a food allergen has been ingested, but *no symptoms*:	☐ Epinephrine ☐ Antihistamine
▪ Mouth	Itching, tingling, or swelling of lips, tongue, mouth	☐ Epinephrine ☐ Antihistamine
▪ Skin	Hives, itchy rash, swelling of the face or extremities	☐ Epinephrine ☐ Antihistamine
▪ Gut	Nausea, abdominal cramps, vomiting, diarrhea	☐ Epinephrine ☐ Antihistamine
▪ Throat†	Tightening of throat, hoarseness, hacking cough	☐ Epinephrine ☐ Antihistamine
▪ Lung†	Shortness of breath, repetitive coughing, wheezing	☐ Epinephrine ☐ Antihistamine
▪ Heart†	Thready pulse, low blood pressure, fainting, pale, blueness	☐ Epinephrine ☐ Antihistamine
▪ Other†	_____	☐ Epinephrine ☐ Antihistamine
▪	If reaction is progressing (several of the above areas affected), give:	☐ Epinephrine ☐ Antihistamine

The severity of symptoms can quickly change. †Potentially life-threatening.

DOSAGE

Epinephrine: inject intramuscularly (circle one) EpiPen® EpiPen® Jr. Twinject™ 0.3 mg Twinject™ 0.15 mg *(see reverse side for instructions)*

Antihistamine: give_____
medication/dose/route

Other: give_____
medication/dose/route

◆ STEP 2: EMERGENCY CALLS ◆

1. Call 911 (or Rescue Squad: _____). State that an allergic reaction has been treated, and additional epinephrine may be needed.

2. Dr. _____ at _____

3. Emergency contacts:

Name/Relationship	Phone Number(s)	
a. _____	1.)_____	2.) _____
b. _____	1.)_____	2.)_____
c. _____	1.)_____	2.) _____

**EVEN IF PARENT/GUARDIAN CANNOT BE REACHED,
DO NOT HESITATE TO MEDICATE OR TAKE CHILD TO MEDICAL FACILITY!**

Parent/Guardian Signature_____ Date_____

Doctor's Signature_____ Date_____
(Required)