

SEIZURE ACTION PLAN

Effective Date: _____

THIS STUDENT IS BEING TREATED FOR A SEIZURE DISORDER. THE INFORMATION BELOW SHOULD ASSIST YOU IF A SEIZURE OCCURS DURING SCHOOL HOURS.

Student: _____ Birthdate: _____

Mother/Guardian: _____ Home Telephone: _____

Cell: _____ Work Telephone: _____

Father/Guardian: _____ Home Telephone: _____

Cell: _____ Work Telephone: _____

Treating Physician: _____ Telephone: _____

Significant Medical History: _____

Current Medications: _____

Allergies: _____

SEIZURE EMERGENCY PROTOCOL

Diazepam rectal gel _____ mg rectally PRN for seizure > _____ minutes OR for _____ or more seizures in _____ hours. Following administration of Diazepam rectal gel, call parents to pick up child.

Use Vagal Nerve Stimulator (VNS) magnet.

Other: _____

Call 911 if:

Seizure does not stop by itself or with VNS within _____ minutes.

Seizures does not stop within _____ minutes of giving Diazepam rectal gel.

Child does not start waking up within _____ minutes after seizure is over (NO Diazepam rectal gel given).

Child does not start waking up within _____ minutes after seizure is over (AFTER Diazepam rectal gel given).

Following a seizure:

Child should rest in clinic.

Child may return to class (specify time frame: _____)

Notify parents immediately.

Send a copy of the seizure record home with child for parents.

Notify physician.

Other: _____

Seizure Information: Student may experience some or all of the listed symptoms during a specific seizure.

<i>Seizure Type(s)</i>	<i>Description</i>
<input type="checkbox"/> Absence	<ul style="list-style-type: none"> • Staring • Eye blinking • Loss of awareness • Other
<input type="checkbox"/> Simple partial	<ul style="list-style-type: none"> • Remains conscious • Distorted sense of smell, hearing, sight • Involuntary rhythmic jerking/twitching on one side • Other _____
<input type="checkbox"/> Complex partial	<ul style="list-style-type: none"> • Confusion • Not fully responsive/unresponsive • May appear fearful • Purposeless, repetitive movements • Other _____
<input type="checkbox"/> Generalized tonic-clonic	<ul style="list-style-type: none"> • Convulsions • Stiffening • Breathing may be shallow • Lips or skin may have blush color • Unconsciousness • Confusion, weariness, or belligerence when seizure ends • Other _____

Seizure usually lasts _____ minutes and returns to baseline in _____ minutes.

Triggers or warning signs: _____

Call parents under the following circumstances:

1. _____

2. _____

Basic Seizure First Aid
<ul style="list-style-type: none"> • Stay calm and track time • Keep child safe • Do not restrain • Do not put anything in mouth • Stay with child until fully conscious • Record seizure in log
For tonic-clonic (grand mal) seizure:
<ul style="list-style-type: none"> • Protect head • Keep airway open/watch breathing • Turn child on side

A seizure is generally considered an EMERGENCY when:
<ul style="list-style-type: none"> • A convulsive (tonic-clonic) seizure lasts longer than 5 minutes • Student has repeated seizures without regaining consciousness • Student has a first-time seizure • Student is injured or has diabetes • Student has breathing difficulties • Student has a seizure in water

Special Considerations and Safety Precautions (regarding school activities, sports, trips, etc.):

Parent/Guardian Signature

Date

Physician Signature

Date