

FENTANYL

THE INVISIBLE KILLER



Drugs may contain deadly level of fentanyl. You cannot see, taste, or smell it. It is almost impossible to tell if a drug is laced with fentanyl unless tested with a fentanyl testing strip.

Fentanyl is a synthetic opioid that is **50-100 times more potent than morphine, often lethal with as little as 2 milligrams.**

one
pill kills

don't make a deadly decision

This small amount is a
LETHAL
dose of fentanyl



one pill kills | don't make a deadly decision

99% of fentanyl overdoses are **accidents.**

The DEA lab testing reveals that

4 out of every **10** pills with fentanyl, contain a potentially

LETHAL DOSE






The number of fentanyl-related deaths in Texas has **almost doubled in just 1 year.**



= 10 fentanyl-related deaths

Fentanyl poisoning is the leading cause of death for people ages 18 to 45 in the U.S.*



Be

SMART!



About Medicine & Pills

ONLY take medicine from your parents, guardians, a doctor, or your school nurse.



Medicine or pills from friends or strangers can make you **SICK**.

Some pills may look like candy but they can **HURT** you.

**THIS IS
NOT
CANDY!**



WHAT TO DO IF YOU THINK SOMEONE IS OVERDOSING:

It may be hard to tell whether a person is high or experiencing an overdose. If you aren't sure, it's best to treat the situation like an overdose—**you could save a life.**

- 1 Call 911 immediately.
- 2 Administer Naloxone or Narcan, if available.
- 3 Try to keep the person awake and breathing.
- 4 Lay the person on their side to prevent choking.
- 5 Stay with them until emergency workers arrive.

Elementary Questions

- What pills are safe to take when you are sick?
- Why is it dangerous to take pills that are not yours?
- What would you say to a peer who offers you a pill?