

**PESTICIDE STANDARD WRITTEN NOTIFICATION
FOR SCHOOLS, DAY CARE PROGRAMS, AND SCHOOL-AGE CHILDCARE PROGRAMS**

- The school, day care center, and/or school-age childcare program is responsible for sending this standard written notification form to employees, pupils, parents etc. to insure that they receive this information at least 2 working days prior to any pesticide use.
- It is recommended that the Pest Management Professional use this ready-to-copy standard written notification form for the purpose of providing pesticide use information to the school, day care center, and/or school-age childcare program. The Pest Management Professional should save this form for copying.

School: R.E. Shaw Elementary School 58 Elmwood School Millbury MA

Name of School , Day care center, and/or School age childcare program

Pest Management Company: Ford's Hometown Services 549 Grove St Worcester MA
(Please Print) Name Address

Pest Management Professional: Matt 49339, Shane 44806, Doug 48588,
(Please Print) License number

A. List the Approximate Dates on which the pesticide use shall commence and conclude

Beginning Date 05/03/19 **Ending Date** 05/10/19

B. Record the specific location of the anticipated pesticide use

see graph for treatment area

C. Pesticide Information (Pest Management Professional should be specific as is possible when listing product(s) to be used)


Pesticide Product Name	Pesticide Type	EPA Registration #	Description/Purpose of treatment and/or application
1. Maxxthor SC	Bifenthrin	81824-5	border area for ticks
2.			
3.			
4.			
5.			

This standard written notification must be accompanied by the following 2 documents. These materials are available from the MDAR web page www.mass.gov/agr. Follow the links to the Children's Protection page.

- Chemical Specific Fact Sheet(s)
- Consumer Information Bulletin for school, day care center, and/or school-age childcare program.

Shaw Elementary School - 58 Elmwood St - Milbury

11/16/15





04/13

MAXXTHOR SC

For use by individuals/firms licensed or registered by the state to apply termiticide products when used as a termiticide. States may have more restrictive requirements regarding qualifications of persons using this product. Consult the structural pest control regulatory agency of your state prior to use of this product.

To control structural pests indoors and outdoors in and around residential, commercial, industrial, institutional and public structures and buildings.

To control pests of lawns and ornamental plants in lawns, parks, grounds, landscapes, recreational areas and athletic fields.

Active Ingredient:	By Wt.
Bifenthrin*	7.9%
Other Ingredients:	92.1%
TOTAL:	100.0%

*Cis isomers 97% minimum, trans isomers 3% maximum.

EPA Reg. No. 81824-5 EPA Est. No. 81824-NC-001

Maxxthor SC contains 2/3 pounds active ingredient per gallon.

(PRECAUCION AL USUARIO: Si usted no puede leer o entender ingles, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.)

(TO THE USER: If you cannot read and understand English, do not use this product until the label has been fully explained to you.)

For product use information call 1-866-FOR-THOR (367-8467) or visit www.maxxthor.com.

KEEP OUT OF REACH OF CHILDREN

CAUTION

ENSYSTEX II, Inc.

2175 Village Drive
Fayetteville, NC 28304

Net Contents: As marked on container

FIRST AID

If swallowed	<ul style="list-style-type: none">• Call poison control center or doctor immediately for treatment advice.• Have person sip a glass of water if able to swallow.• Do not induce vomiting unless told to do so by the poison control center or doctor.• Do not give anything to an unconscious person.
If inhaled	<ul style="list-style-type: none">• Move person to fresh air.• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible.• Call a poison control center or doctor for further treatment advice.
If on skin or clothing	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15 to 20 minutes.• Call a poison control center or doctor for treatment advice.
If in eyes	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice.

HOTLINE NUMBER

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-(888)-398-3772 for emergency medical treatment information.

NOTE TO PHYSICIAN

This product is a pyrethroid. If large amounts have been ingested, the stomach and intestine should be evacuated. Treatment is symptomatic and supportive. Digestible fats, oils, or alcohol may increase absorption and so should be avoided.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION

Harmful if swallowed, inhaled or absorbed through skin. Avoid contact with skin, eyes or clothing. Avoid breathing spray mist. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove contaminated clothing and wash before reuse.

Personal Protective Equipment: All pesticide handlers (mixers, loaders and applicators) must wear long-sleeved shirt and long pants, socks, shoes and chemical-resistant gloves. All pesticide handlers must wear a respiratory protection device when working in a non-ventilated space (such as a NIOSH approved respirator with any R, P or HE filter or a NIOSH approved respirator with an organic vapor (OV) cartridge or canister with any R, P or HE pre-filter). All pesticide handlers must wear protective eyewear, such as goggles, faceshield or safety glasses, when working in a non-ventilated space or when applying as a termiticide by rodding or sub-slab injection. After the product is diluted in accordance with label directions for use (or if an in-line injector system is used) shirt, pants, socks, shoes and waterproof gloves are sufficient.

Environmental Hazards

This pesticide is extremely toxic to fish and aquatic invertebrates. Drift and run-off from treated areas may be hazardous to aquatic organisms in neighboring areas. Care should be used when spraying to avoid fish and reptile pets in/around ornamental ponds. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies and drainage systems. This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

Physical and Chemical Hazards

Do not apply this product or solutions of this product around electrical equipment, such as electrical conduits, motor housings, junction boxes, switch boxes, etc. due to the possibility of shock hazard.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do not apply as a broadcast application to interior wall and floor surfaces of homes except as otherwise permitted.

Do not apply by aircraft or through an irrigation or chemigation system.

Do not allow spray to drift onto ponds, streams or lakes.

Do not apply in greenhouses or nurseries.

Not for use on sod farm turf, golf course turf or grass grown for seed.

Do not water the treated area to the point of run-off.

Do not make applications during rain.

Application is prohibited directly into sewers or drains, or to any area like a gutter where drainage to sewers, storm drains, water bodies, or aquatic habitat can occur. Do not allow the product to enter any drain during application.

Termite Control Treatment: When treating adjacent to an existing structure, the applicator must check the area to be treated and immediately adjacent areas of the structure for visible and accessible cracks and holes to prevent any leaks or significant exposures to persons occupying the structure. People present or residing in the structure during application must be advised to remove their pets and themselves from the structure if they see any signs of leakage. After application, the applicator is required to check for leaks. All leaks resulting in the deposition of termiticide in locations other than those prescribed on this label must be cleaned up prior to leaving the application site. Do not allow people or pets to contact contaminated areas or to reoccupy contaminated areas of the structure until the clean-up is completed.

Additional Application Restrictions for Pre-Construction Termiticide Applications:

The treatment site must be covered prior to a rain event in order to prevent run-off of the pesticide into non-target areas.

The applicator must either cover the soil him/herself or provide written notification of the above requirement to the contractor on site and to the person commissioning the application (if different from the contractor). If notice is provided to the contractor or the person commissioning the application, then they are responsible under FIFRA to ensure that 1) if the concrete slab cannot be poured over the treated soil within 24 hours of application the treated soil is covered with a waterproof covering (such as polyethylene sheeting), and 2) the treated soil is covered if precipitation is predicted to occur before the concrete slab is scheduled to be poured.

Do not treat soil that is water-saturated or frozen.

Do not treat when raining.

Do not allow treatment to run off from the target area.

Do not apply within 10 feet of storm drains. Do not apply within 25 feet of aquatic habitats (such as, but not limited to, lakes; reservoirs; rivers; permanent streams; marshes or ponds; estuaries; and commercial fish farm ponds).

Do not make on-grade applications when sustained wind sweeps are above 10 mph (at application site) at nozzle end height.

Additional Application Restrictions for Residential Outdoor Surface and Space Sprays:

All outdoor applications must be limited to spot or crack-and-crevice treatments only, except for the following permitted uses:

(1) Treatment to soil or vegetation around structures;

(2) Applications to lawns, turf, and other vegetation;

(3) Applications to building foundations, up to a maximum height of 3 feet.

Other than applications to building foundations, all outdoor applications to impervious surfaces such as sidewalks, driveways, patios, porches and structural surfaces (such as windows, doors and eaves) are limited to spot and crack-and-crevice applications only.

Application is prohibited directly into sewers or drains, or to any area like a gutter where drainage to sewers, storm drains, water bodies, or aquatic habitat can occur. Do not allow the product to enter any drain during or after application.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: Keep out of reach of children and animals. Store in original containers only. Store in a cool, dry place and avoid excess heat. Handle and open container in a manner so as to prevent spillage. Do not put concentrate or dilute material into food or drink containers.

Pesticide Disposal: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label directions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

Container Disposal: Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying.

Nonrefillable containers less than or equal to 5 gallons: Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Nonrefillable containers greater than 5 gallons: Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip the container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over on to its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

In Case of Spill: Confine it, avoid contact, isolate area and keep animals and unprotected persons away. Form dike around spill area and/or absorb spill with absorbent materials, such as sand, cat litter or clay. Place damaged package in a holding container and identify contents. Contact Ensysyst II at 1-888-398-3772 for any assistance.

Tip and Measure Container Dispensing Directions

1. Remove the cap from the measuring chamber and remove seal over container opening.
2. Replace the cap securely.
3. Tip the container until the measuring chamber contains the desired amount of product.
4. Return container to its level position.
5. Remove the cap from the measuring chamber and pour product into the application equipment.
6. Replace cap securely.

APPLICATION FOR CONTROL OF SUBTERRANEAN TERMITES

General

Maxxthor SC, in the form of a dilute insecticidal solution, prevents and controls subterranean termite infestations in and around structures and other items by creating a continuous chemically treated zone (horizontal and/or vertical as needed) between the wood and other cellulose material in the structure and termite colonies in the soil. In order to establish a barrier between the wood in the structure and the termites in the soil, adequately disperse the solution of this product in the soil. Maxxthor SC can also be used to directly treat termite infested wood, however such treatments should be considered to be supplementary to, and not a replacement for, soil based applications of this product.

To effectively control termites with this product, the service technician should be familiar with current termite control practices including trenching, rodding, sub-slab and void injection, soil surface fan spraying, excavated soil treatment and brush, spray and injection applications to wood. Correct use of these techniques is necessary to effectively control infestations by subterranean termites such as *Coptotermes*, *Heterotermes*, *Reticulitermes* and *Zootermopsis*. The service technician should consider the biology and behavior of the termite specie(s) to be controlled to determine which control practices to use.

Treatment standards and procedures for subterranean termite control may vary due to regulations, water table level, structure design, soil types, construction practices and other factors. For advice concerning current control practices with respect to specific local conditions, consult resources in structural pest control and state cooperative extension and regulatory agencies. Follow all federal, state and local regulations and treatment standards for protection of a structure from subterranean termites.

Effective termite control may also include mechanical alteration of the structure. Elimination of leaks or points of moisture accumulation within or on the exterior of the structure that result in an increase in the moisture content of wooden structural components is advised. Removal of non-essential cellulose containing materials that are in contact with the ground under or around the structure can reduce termite foraging in the area. When untreated wooden parts of the structure touch the ground and such contact cannot be broken, creating a barrier between the soil and such components using a solution of the product may protect the components and the structure against termite attack.

Maxxthor SC is labeled for use against subterranean termites as a 0.06% - 0.12% solution in water, however the 0.06% finished solution should be used for typical control situations. When difficult or problem soils or construction types are encountered, it may be necessary to use 0.12% Maxxthor SC mixed in reduced volumes of water. All treatment directions contained in this label may not be necessary to provide adequate protection against termites.

Avoid contamination of water supplies due to backflow under reduced water system pressure by using anti-backflow equipment or procedures to prevent siphoning of any solution back into a water supply. Do not contaminate cisterns or wells. Do not treat soil that is water saturated or frozen. Do not treat while precipitation is occurring. Do not apply solution to an area or site if the soil at the area or site is in such a state or condition that runoff or movement of the solution from the treated area or site is likely to occur. Consult state and local specifications for recommended distances of wells from treated areas, or if such regulations do not exist, refer to Federal Housing Administration Specifications (H.U.D.) for guidance.

For the purposes of this label and its directions, crawl spaces are to be considered to be inside of the structure.

Mixing Directions For Maxxthor SC For Use As A Termiticide

Mix Maxxthor SC for use as a termiticide in the following manner:

1. Fill tank 1/4 to 1/3 full.
2. Start pump to begin by-pass agitation and place end of treating tool in tank to allow circulation through hose.
3. Add appropriate amount of Maxxthor SC.
4. Add remaining amount of water.
5. Let pump run and allow recirculation through the hose for 2 to 3 minutes.

Maxxthor SC may also be mixed into full tanks of water, but substantial agitation is required to ensure uniformity of the solution.

Dilution of Maxxthor SC for Use as a Termiticide

For termite control, use rates for Maxxthor SC are expressed and the solution is mixed according to the percentage (%) concentration it forms when mixed in water. Use the mixing table or alternately the formulas below to determine the amount of Maxxthor SC to add to any quantity of water.

To prepare a 0.06% water solution, ready to use, dilute 3 quarts of Maxxthor SC with 99.25 gallons of water. To prepare a 0.12% water solution, ready to use, dilute 6 quarts of Maxxthor SC with 98.5 gallons of water.

Mixing Table for Maxxthor SC for Use as a Termiticide			
Solution Percentage Concentration Desired	Gallons of Finished Solution Desired	Amount of Maxxthor SC to add	Water to mix with Maxxthor SC
0.06%	1	1 oz	127 oz.
	5	5 oz	4.9 gal.
	25	25 oz.	24.8 gal.
	50	1 qt.+16 oz.	49.6 gal.
	100	3 qt	99.25 gal.
0.12%*	1	2 oz	126 oz.
	5	10 oz	4.9 gal.
	25	1 qt.+18 oz.	24.6 gal.
	50	3 qt.	49.2 gal.
	100	6 qt	98.5 gal.

*Only use the 0.12% rate in accordance with the *Adjustments to Application Volume* section. May also be used in accordance with the *FOAM APPLICATION and APPLICATIONS TO PROTECT UNDERGROUND ITEMS FROM TERMITE ATTACK* sections.

Calculating an Amount of Maxxthor SC to Mix

To mix any amount of Isothor SC for termite control, determine:

A = Units of water into which the Maxxthor SC will be mixed. Express any partial units as decimal fractions (1/2 = .5). Any unit of measure, such as gallons or quarts, can be used for A. Answers to equations below are in same units as A.

Maxxthor SC to add to A for 0.06% = A / 132.3

Maxxthor SC to add to A for 0.12% = A / 65.7

To convert gallons to fluid ounces, multiply number of gallons X 128

128 fluid ounces = 16 cups = 8 pints = 4 quarts = 1 gallon

32 fluid ounces = 1 quart

Application Volume

To provide maximum control and protection against termite infestation, apply the specified volume of the finished water solution containing the specified amount of Maxxthor SC as set out below or as otherwise directed in this label.

Prescribed Horizontal Barrier Rate: Unless otherwise directed, horizontal barriers are created by applying a 0.06% solution at a rate of one gallon of solution per 10 square feet. (One gallon of 0.06% solution contains 1.0 fluid ounce of Maxxthor SC.)

Prescribed Vertical Barrier Rate: Unless otherwise directed, vertical barriers are created by applying a 0.06% solution at a rate of four gallons of solution per 10 linear feet per foot of depth. (Four gallons of 0.06% solution contains 4.0 fluid ounces of Maxxthor SC.)

Adjustments to Application Volume

If soil will not accept the labeled application volumes, the volume may be reduced provided there is a corresponding increase in concentration so that the amount of active ingredient applied to the soil remains the same.

Note: Large reductions of application volume reduce the likelihood of obtaining a continuous barrier. Variance is allowed when volume and concentration are consistent with label directed rates and a continuous barrier can still be achieved. When volume is reduced, the spacing of holes created for sub slab injection and soil rodding may need to be reduced to account for decreased dispersion of the solution in the soil.

For example, adjust the amount of solution applied to deliver a horizontal barrier of 10 square feet from 1 gallon to as low as 0.5 gallons and as high as 2 gallons while maintaining the amount of Maxxthor SC (1 fluid ounce) applied per 10 square feet.

For example, adjust the amount of solution applied to deliver a vertical barrier 10 feet long by one foot deep from 4 gallons to as low as 2 gallons and as high as 8 gallons while maintaining the amount of Maxthor SC (4 fluid ounces) applied per 10 linear feet.

PRE-CONSTRUCTION TREATMENT

All Structures

Pre-construction treatment: Do not apply at a lower dosage and/or concentration than specified on this label for applications prior to the installation of the finished grade.

Prior to each application, applicators must notify the general contractor, construction superintendent, or similar responsible party, of the intended termiticide application and intended sites of application and instruct the responsible person to notify construction workers and other individuals to leave the area to be treated during application and until the termiticide is absorbed into the soil.

Effective control of subterranean termites can be accomplished during construction by using a 0.06% solution of Maxthor SC to establish vertical and/or horizontal barriers between the structure and the soil as directed. To meet current termite proofing requirements, follow the procedures in the latest edition of the Housing and Urban Development Minimum Property Standards.

Horizontal Barriers Under Slabs on Ground Including Basements

Create a horizontal barrier on the entire surface of soil or substrate that will be covered by a slab, including slab floors, garages, carports, basements, porches and entrance platforms by treating the soil or substrate with the solution at the Prescribed Horizontal Barrier Rate.

If the fill under the slab is a coarse material such as washed gravel, make sure that a sufficient enough amount of dilution is applied that the solution reaches the soil beneath the fill.

Apply solution using a coarse spray nozzle. If the slab over the treated area will not be poured on the same day as the application (and there are no foundation walls in place around the treated soil) cover treated soil with a water-proof barrier such as polyethylene sheeting.

Vertical Barriers

Create a vertical barrier along the inside and outside of foundation walls, around piers, plumbing and utility service entrances and other points of possible future termite access and entry by treating the soil at these points at the Prescribed Vertical Barrier Rate. When trenching and rodding into the trench, or trenching alone, it is important that the solution reaches the top of the footing. Rod holes must be spaced so as to achieve a continuous termiticidal barrier, but they should in no case be more than 12 inches apart. Trenches need not be wider than 6 inches. Mix the solution into the soil as it is being replaced in the trench. Care should be taken to avoid washing soil out from around footings thereby undermining the stability of the structure. An inside vertical barrier may not be required for a monolithic slab.

If distance from final grade to top of footing will be less than four feet, it is permissible to wait until final grade is established to apply the vertical barrier. When treating foundations deeper than 4 feet, apply the termiticide as the backfill is being replaced, or if the construction contractor fails to notify the applicator in time to permit this, treat the foundation to a minimum depth of 4 feet after the backfill has been installed. The applicator must trench and rod into the trench or trench along the foundation walls and around pillars and other foundation elements, at the rate prescribed from grade to a minimum depth of 4 feet. When the top of the footing is exposed, the applicator must treat the soil adjacent to the footing to a depth not to exceed the bottom of the footing. However, in no case should a structure be treated below the footing.

Hollow Block Foundations and Voids

Hollow block foundations and voids may be treated at a rate of 2 gallons of solution per 10 linear feet to create a continuous treated zone within the voids at the footing.

POST CONSTRUCTION TREATMENT

All Structures

Do not apply treatment until the identity and location of all wells, radiant heat pipes, water and sewer lines, electrical conduits and sub-slab heating and air conditioning ducts is established. Caution must be taken to avoid puncturing these elements and/or injecting solution into them. All holes in commonly occupied areas into which material has been applied must be plugged. Plugs must be of a non-cellulose material or covered by an impervious, non-cellulose material.

Vertical Barrier Depth: For applications made after the final grade is installed, the applicator must trench and rod into the trench or trench along the foundation walls and around pillars and other foundation elements and treat at the rate prescribed from grade to the top of the footing. When the footing is more than four (4) feet below grade, the applicator must trench and rod into the trench or trench along the foundation walls and treat at the rate prescribed to a minimum depth of four feet. The actual depth of treatment will vary depending on the soil type, degree of compaction, and location of termite activity. When the top of the footing is exposed, the applicator must treat the soil adjacent to the footing to a depth not to exceed the bottom of the footing. However, in no case should a structure be treated below the footing.

Structures Containing Concrete Slabs on Ground (Monolithic/Floating/Supported) including Basements

To make an application beneath existing slabs, it may be necessary to drill holes in the slab or adjacent foundation and to apply solution. Holes should be spaced such that when treatment is applied through them, a continuous treated zone is applied beneath the slab.

Vertical Barriers Along Exterior of Foundation Walls: Trench and rod into the trench or trench along the outside of foundation walls and treat at the Prescribed Vertical Barrier Rate to the depth specified under Vertical Barrier Depth. Where physical obstructions such as concrete walkways adjacent to foundation elements or soil type and/or conditions make trenching prohibitive, treatment may be made by rodding alone.

Vertical Barriers Along Interior of Foundation Walls: Vertical barriers may be established on the interior side of foundation walls by sub-slab injection of the solution at the Prescribed Vertical Barrier Rate. Injection openings can be drilled either vertically through the slab along the interior of the foundation wall or horizontally from the exterior through the foundation wall low enough on the wall to allow for the deposition of the solution beneath the slab along the interior side of the foundation wall. Drill holes should be spaced so as to achieve a continuous chemical barrier but in no case farther apart than 12 inches. Special care must be taken to distribute the solution evenly. Vertical barriers may also be established beneath the slab along both sides of interior footing-supported walls, one side of interior partitions and along all cracks and expansion joints and utility service entrances and bath traps.

Horizontal Barriers Beneath Slabs on Ground: Create a horizontal barrier by treating at the Prescribed Horizontal Barrier Rate beneath slabs by either drilling and long rodding from the exterior or by grid pattern drilling and injection vertically through the slab. Long rodding should be used only when grid pattern drilling and injection and horizontal short rodding and injection cannot be used to deliver the sub slab treatment.

Bath Traps: Exposed soil beneath and around areas where plumbing and utility services penetrate the slab should be treated at the rate of 1 gallon of solution per square foot of soil.

Structures Containing Accessible Crawl Spaces

For crawl spaces, apply vertical termiticide barriers at the rate of 4 gallons of solution per 10 linear feet per foot of depth from grade to the top of the footing, or if the footing is more than 4 feet below grade, to a minimum depth of 4 feet. Apply by trenching and rodding into the trench, or trenching. Treat both sides of foundation and around all piers and pipes. Where physical obstructions such as concrete walkways adjacent to foundation elements prevent trenching, treatment may be made by rodding alone. When soil type and/or conditions make trenching prohibitive, rodding may be used. When the top of the footing is exposed, the applicator must treat the soil adjacent to the footing to a depth not to exceed the bottom of

the footing. Read and follow the mixing and use direction section of the label if situations are encountered where the soil will not accept the full application volume.

1. Rod holes and trenches must not extend below the bottom of the footing.

2. Rod holes must be spaced so as to achieve a continuous termiticide barrier but in no case more than 12 inches apart.

3. Trenches must be a minimum of 6 inches deep or to the bottom of the footing, whichever is less, and need not be wider than 6 inches. When trenching in sloping (tiered) soil, the trench must be stepped to ensure adequate distribution and to prevent termiticide from running off. The solution must be mixed with the soil as it is replaced in the trench.

4. When treating plenums or crawl spaces, turn off the air circulation system of the structure until application has been completed and all solution has been absorbed by the soil.

Structures Containing Inaccessible Crawl Spaces

For inaccessible interior areas, such as areas where there is insufficient clearance between floor joists and ground surfaces to allow operator access, excavate, if possible, and treat according to the instructions for accessible crawl spaces. Otherwise, apply one, or a combination of the following two methods.

1. To establish a horizontal barrier, apply to the soil surface, 1 gallon of solution per 10 square feet overall using a nozzle pressure of less than 25 p.s.i. and a coarse application nozzle (e.g., Delavan Type RD Raindrop RD-7 or larger, or Spraying Systems Co. 8010LP TeeJet or comparable nozzle). For an area that cannot be reached with the application wand, use one or more extension rods to make the application to the soil. Do not broadcast or power spray with higher pressures.

2. To establish a horizontal barrier, drill through the foundation wall or through the floor above and treat the soil perimeter at a rate of 1 gallon of solution per 10 square feet. Drill spacing must be at intervals not to exceed 16 inches. Many states have smaller intervals, so check state regulations which may apply.

When treating plenums or crawl spaces, turn off the air circulation system of the structure until application has been completed and all termiticide has been absorbed by the soil.

Masonry Voids

Drill and treat voids in multiple masonry elements of the structure extending from the structure to the soil in order to create a continuous treatment barrier in the area to be treated. Apply at the rate of 2 gallons of solution per 10 linear feet of footing using a nozzle pressure of less than 25 p.s.i. When using this treatment access holes must be drilled below the sill plate and should be as close as possible to the footing as is practical. Treatment of voids in block or rubble foundation walls must be closely monitored: Applicators must inspect areas of possible runoff as a precaution against application leakage in the treated areas. Some areas may not be treatable or may require mechanical alteration prior to treatment.

All leaks resulting in the deposition of termiticide in locations other than those prescribed on this label must be cleaned up prior to leaving the application site. Do not allow people or pets to contact contaminated areas or to reoccupy the contaminated areas of the structure until the clean-up is completed.

Note: When drilling veneer walls, care should be taken to not drill beyond the depth of the void behind the veneer into another construction layer behind the veneer. It is however permissible to drill through the veneer and into concrete blocks behind the veneer and to treat the veneer and the concrete blocks at the same time.

Note: Not for use in voids insulated with rigid foam.

TREATMENT OF STRUCTURES WITH WELLS AND CISTERNS

Do not contaminate wells or cisterns.

Structures with Wells/Cisterns Inside Foundations

Structures that contain wells or cisterns within the foundation of a structure can only be treated using the following techniques:

1. Do not treat soil while it is beneath or within the foundation or along the exterior perimeter of a structure that contains a well or cistern. The treated backfill method must be used if soil is removed and treated outside/away from the foundation. The treated backfill technique is described as follows:

a. Trench and remove soil to be treated onto heavy plastic sheeting or similar material or into a wheelbarrow.

b. Treat the soil at the rate of 4 gallons of dilute solution per 10 linear feet per foot of depth of the trench, or 1 gallon per 1.0 cubic feet of soil. See *Mixing Directions for Maxthor SC for Use as a Termiticide* section of the label. Mix thoroughly into the soil taking care to contain the liquid and prevent runoff or spillage.

c. After the treated soil has absorbed the solution, replace the soil into the trench.

2. Treat infested and/or damaged wood in place using an injection technique such as described in the APPLICATION TO WOOD INDORS to PROTECT AGAINST WOOD DESTROYING INSECTS or APPLICATION TO WOOD OUTDOORS to PROTECT AGAINST WOOD DESTROYING INSECTS sections of this label.

Structures with Adjacent Wells/Cisterns and/or Other Water Bodies

Applicators must inspect all structures with nearby water sources including wells, cisterns, surface ponds, streams, and other bodies of water and evaluate, at a minimum, the treatment listed below prior to making an application.

1. Prior to treatment, if feasible, expose the water pipe(s) coming from a well to the structure, if the pipe(s) enter the structure within 3 feet of grade.

2. Prior to treatment, applicators are advised to take precautions to limit the risk of applying the termiticide into subsurface drains that could empty into any bodies of water. These precautions include evaluating whether application of the termiticide to the top of the footer may result in contamination of the subsurface drain. Factors such as depth to the drain system and soil type and degree of compaction should be taken into account in determining the depth of treatment.

3. When appropriate (for example, on the water side of the structure), the treated backfill technique (described above) can also be used to minimize offsite movement of termiticide.

FOAM APPLICATION

Maxthor SC, in the form of a foam, can be used to deliver Maxthor SC as a termiticide any time it appears likely this form of delivery will improve the dispersal of Maxthor SC into and within the intended target area. Foam can be particularly useful to deliver Maxthor SC where it either cannot be depended upon to be delivered as just a solution or due to a need to reduce the amount of water used in order to avoid water damage to the target or adjacent areas. In some situations, for example under some slabs, a solution cannot be depended upon to disperse as completely as a foam because of deflection of the liquid stream or some other structural obstacle or defect.

Depending on the circumstances, foam applications of Maxthor SC may be used alone or in combination with liquid solution applications, provided that the cumulative amount of active ingredient per unit of area applied is equivalent to that which would be contained in a 0.06% solution-only application applied to the same area.

Using foam generating equipment, a solution of Maxthor SC, ranging in concentration from 0.06% to 0.12%, may be converted into a foam according to the foaming agent and foaming equipment manufacturer's recommendations.

First, form a solution of Maxthor SC of the appropriate percentage concentration and volume. Then add the recommended volume of a foaming agent. Verify that the foaming agent is compatible with Maxthor SC.

Applications may be made behind veneers, piers, chimney bases, into rubble foundations, into block voids, structural voids or other similar voids, under slabs, stoops, porches or to the soil in crawlspaces.

RETREATMENT

Retreatments for subterranean termites can only be performed if there is clear evidence of reinfestation or disruption of the barrier due to construction, excavation or landscaping and/or evidence of the breakdown of the termiticide barrier in the soil. These vulnerable or reinfested areas may be retreated in accordance with

application techniques described in this product's labeling. The timing and type of these retreatments will vary, depending on factors such as termite pressure, soil types, soil conditions and other factors which may reduce the effectiveness of the barrier.

Annual retreatment of the structure is prohibited unless there is clear evidence that reinfestation or barrier disruption has occurred.

APPLICATION IN CONJUNCTION WITH TERMITE BAITS

Spot only applications of Maxxthor SC can be used as a supplement to termite baiting systems. For the purposes of this section, spot only applications are defined as the use of Maxxthor SC according to any of the permitted and applicable post-treatment application techniques contained in this label, alone or in combination, to the extent needed or deemed necessary or useful as an adjunct to a termite bait product labeled as a stand alone protection against termite attack. Stand alone termite bait product is defined as a termite bait product that provides sufficient structural protection when applied without other termite control products.

APPLICATION TO PROTECT UNDERGROUND ITEMS FROM SUBTERRANEAN TERMITE ATTACK

To protect components installed underground such as wires, conduits, cables and pipes buried in soil against termite attack, create an envelope of Maxxthor SC treated soil around the components along the entire underground length of the component. First, treat soil through which components will be run with 0.06% to 0.12% solution of Maxxthor SC at a rate of 2 gallons of solution per 10 linear feet. Install components, laying them on the treated soil. Cover components with untreated soil and then treat this covering soil using the same percent solution at 2 gallons of solution per 10 linear feet.

Underground components to be protected may be located within the foundation of a structure or outside of a structure such as within a utility right of way, for example. Do not treat items that are electrically energized at the time of application. If the soil will not absorb the indicated amount of solution, as little as 1 gallon of 0.12% solution per 10 linear feet can be used. Treat points where services emerge from the ground at a rate of 1 to 2 gallons of solution at the point of emergence.

APPLICATIONS TO PROTECT POLES, POSTS AND OTHER WOODEN ITEMS FROM SUBTERRANEAN TERMITE ATTACK

Maxxthor SC can be used to protect the below ground portions of wooden structural components from termites. Form a treated zone around components below ground by vertically rodding the soil around their perimeter to a depth of six inches below their maximum depth of placement in the soil and applying a 0.06% solution of Maxxthor SC at a rate of 0.4 gallons of solution per linear foot of perimeter around the component per foot of treated depth. Measure the perimeter of the component six inches from the outside of the component.

APPLICATIONS TO CONTROL WOOD INFESTING INSECTS

Mix and dilute Maxxthor SC according to the instructions contained in the APPLICATION FOR CONTROL OF SUBTERRANEAN TERMITES section.

APPLICATIONS TO CONTROL EXPOSED WOOD INFESTING INSECTS LOCATED ABOVE GROUND

Maxxthor SC can be applied above ground as a non-soil treatment to control and kill exposed worker and winged reproductive (swarmer) termites and carpenter ants. This type of application is only supplementary to and should not be considered as a replacement for soil treatment in the case of termite infestation.

Apply a 0.06% solution as a general fan spray within attics, crawl spaces, unfinished basements and other void areas where termites have been found. Apply treatment directly to swarming termites and areas where they congregate.

APPLICATIONS TO TERMITE CARTON NESTS LOCATED IN ABOVE GROUND WALL VOIDS

Apply a 0.06% solution of Maxxthor SC directly into above ground termite carton nests including nests located in wall voids using a directional injector. Apply as a solution or foam under pressure to distribute solution thoroughly throughout the nest. It may be necessary to inject solution at one or more points and at varying depths within the nest to adequately distribute solution within the interior of the nest.

APPLICATION TO WOOD INDOORS TO PROTECT AGAINST WOOD INFESTING INSECTS

A 0.06% solution of Maxxthor SC applied as a liquid or foam to wooden structural components within structures can be used to protect them against attack from wood infesting insects such as termites, carpenter ants and wood boring beetles or borers. This type of application is only supplementary to and should not be considered a replacement for soil treatment in the case of termite infestation.

Apply solution as a general fan spray onto the surface of the wood or inject solution under pressure into the wood as a liquid or foam. Inject by either injecting solution through a directional injector directly into existing voids and galleries or by drilling wood to form treatment channels through which the solution can be injected into the insect galleries. Multiple treatment channels of varying depth may need to be drilled to adequately distribute the solution within the wood interior. Application can also be made with a paintbrush.

Before application, locate heat pipes, ducts, water and sewer lines and electrical conduits. Take precautions to avoid puncturing and/or injecting solution into these items. Do not apply solution inside of electrified enclosures, switches or sockets.

Plastic sheeting must be placed below any indoor overhead surfaces being treated that are located anywhere except within a soil-based crawl space. Wear protective clothing, unvented goggles, gloves and respirator when making an overhead application or when applying in poorly ventilated indoor areas. Do not touch surfaces until spray has dried.

When treating in home food preparation and storage areas, cover all food preparation surfaces and utensils prior to beginning treatment. Surfaces or items that cannot be covered or removed should be thoroughly washed after treatment and before use. Food that cannot be covered should be removed. Before application, remove pets, birds and cover and disconnect aquariums. Do not allow humans and pets to touch treated surfaces until they have dried.

APPLICATION TO WOOD OUTDOORS TO PROTECT AGAINST WOOD DESTROYING INSECTS

On and around structures

A 0.06% solution of Maxxthor SC applied as a liquid or foam to exterior structural components can be used to protect them from the attack of wood infesting insects such as termites, carpenter ants and wood boring beetles or borers. Apply solution by general fan spray to the point of runoff onto the surface of the wood or inject solution under pressure into the wood as a liquid or foam. Inject by either injecting solution directly into existing voids and galleries or drilling wood to form treatment channels through which the solution can be injected into the insect galleries. Multiple treatment channels of varying depth may need to be drilled to adequately distribute the solution within the wood interior. Application can also be made with a paintbrush.

To control carpenter ants, also apply solution around doors and windows and other areas carpenter ants have been observed or can be expected to forage.

Within wooden components and trees

Locate the cavity that the insects (such as carpenter ants) are inhabiting. Inject a 0.06% solution of Maxxthor SC into the cavity as a solution or foam. Apply a sufficient volume of solution to completely flood or fill the cavity. Drill injection channels if necessary.

APPLICATION FOR LAWN PESTS

Maxxthor SC controls the listed turfgrass insect and mite pests. Maxxthor SC is not for use on turfgrass grown for sale, commercial seed production, for other commercial use or for research. Not for use on golf course turf.

Application Sites

For use on any type of landscape or recreational turfgrass in any type of setting or at any type of site except as otherwise prohibited. Permitted sites include lawns, grounds, recreational areas, parks, landscapes and athletic fields. New York State only: Do not apply Maxxthor SC containing solutions to grass or turf within 100 feet of a body of water (lake, pond, river, stream, wetland or drainage ditch).

Application Precautions and Preparations

Keep children and pets off treated areas until spray has dried.

If necessary, test the effects of applications of different rates and volumes of mixed solution on a small patch of a type of grass (with observations over one week to detect the occurrence of negative effects) before application of solutions to large areas of that type of grass.

Maxxthor SC can be mixed with other pesticides, including insect growth regulators. Follow the label directions of all the products mixed, making sure not to exceed the labeled application rate of any individual product in the mixture. Any tank mixture that has not been tested before should be tested before full scale use by first mixing a small quantity of the mixture to ensure there is no physical or chemical incompatibility.

Application Methods

Apply solution as a broadcast treatment in a quantity sufficient to wet all foliage. If necessary, consult resources in horticulture in your area (such as your Cooperative Extension Service) to determine appropriate application timing and cultural practices to control different types of pests.

Reapplication

Reapplications may be necessary particularly in the event of high pest pressure. Reapply as needed based on pest reinfestation. Reapplications must not occur more often than every 7 days. New York State only: Do make a single reapplication of Maxxthor SC if there is renewed insect activity, but not sooner than two weeks after first application.

Lawn Application Use Rates and Volumes

Use rates for Maxxthor SC for lawn pests are stated in fluid ounces of Maxxthor SC per 1000 square feet.

Apply Maxxthor SC at 0.18 to 1.0 fluid ounces per 1000 square feet depending on the target pest. Rates for specific pests within this range are given below. However, applications of up to 1.0 fluid ounce per 1000 square feet are permitted at the discretion of the applicator to control any pest.

Depending on the length and/or density of grass being treated, application volumes of up to 10 gallons per 1000 square feet can be made in order to obtain uniform coverage. If a minimal volume application is made (2 gallons per 1000 square feet) and the target pest, such as mole crickets or chinch bugs for example, are located in the thatch or below the surface, irrigate the treated area with at least 0.25 inches of water immediately after the application.

The calculated amount of Maxxthor SC can be applied in any volume of water as long as the maximum label rate of 1.0 fluid ounce per 1000 feet is not exceeded. Do not exceed the maximum label rate by applying solution to an area smaller than intended when it was mixed and diluted unless such under dosing will not result in an application rate per 1000 square feet in excess of the maximum label rate.

Lawn Pest Application Use Rates

The application rates listed below provide control of the listed pests under normal conditions. At the discretion of the applicator, up to 1.0 fluid ounce of Maxxthor SC per 1000 square feet can be used to control any of the listed pests. Use the higher application rates when maximum residual control is needed.

Use Rate Table for Maxxthor SC for Lawn Applications	
Use Rate	Fluid ounces Maxxthor SC per 1,000 square feet (Range)
A	0.18⇒0.25
B	0.25⇒0.50
C	0.50 ⇒1.00

Lawn Pests Grouped by Use Rates

Use Rate A: Armyworms, Cutworms, Sod Webworms

Use Rate B: Annual Bluegrass or Hyperodes Weevil (Adult), (Banks Grass Mite, Billbugs (Adult), Black Turfgrass Ataenius (Adult), Centipede, Chinch Bug, Cricket, Earwigs, Fleas (Adult), Grasshoppers, Leafhoppers, Mealybugs, Millipedes, Mites, Pillbugs, Sowbugs

Use Rate C: Ants, Fleas (Larvae), Imported Fire Ant, Japanese Beetle (Adult), Mole Cricket (Nymph and Adult) Ticks

Mixing Table for Maxxthor SC for Lawn Applications				
Application Volume: Gallons/ 1000 sq. ft.	Use Rate: Fluid oz. / 1000 sq. ft.	Use these amounts of Maxxthor SC diluted to these volumes of finished spray		
		5 gal.	10 gal.	25 gal.
2	.18	.45	.90	2.25
	.25	.63	1.25	3.13
	.50	1.25	2.50	6.25
	1.00	2.50	5.00	12.50
5	.18	.18	.36	.90
	.25	.25	.50	1.25
	.50	.50	1.00	2.50
	1.00	1.00	2.00	5.00
10	.18	--	.18	.45
	.25	--	.25	.63
	.50	--	.50	1.25
	1.00	--	1.00	2.50

Calculating Amounts of Maxxthor SC to Mix for Lawn Pests

To mix and apply any amount of Maxxthor SC for lawn pests, determine:
A = Square feet of area to be treated / 1000 (5,500 sq. ft./1000 = 5.5)
B = Use Rate per 1000 square feet for the target pest(s) in fluid ounces Maxxthor SC taken from the Use Rate Table. If treating for more than one type of pest, select the highest rate.
Calculate the amount of Maxxthor SC to mix for lawn pests as follows:
Fluid Ounces Maxxthor SC to use = A X B
Mix this amount of Maxxthor SC in the amount of water needed to make the application. After mixing, the percent of Maxxthor SC active ingredient contained in the mixture can be determined using the following formula.
% a. i. = (0.0617 X fluid oz. of Maxxthor SC added) / gallons of water in container

Application Against Specific Lawn Pests

Annual Bluegrass or Hyperodes Weevil (Adult): Consult your Cooperative Extension Service for advice on application timing in your area.

Armyworms, Cutworms, Sod Webworms: Do not water or mow grass within 24 hours of application for optimum control of these surface feeding insects.

Billbug Adults: Make application when adults first appear in the spring or when chewed or brown grass indicates damage. Consult your Cooperative Extension Service for advice on more exact application timing in your area.

Chinch Bugs: Water grass immediately after application to help move active ingredient deeper into thatch where these insects live. The highest application rate may be necessary to achieve control during the summer.

Imported Fire Ant: Combine broadcast treatments to control newly invading ants and mound treatments to control existing ant colonies. Treat mounds according to the Structural Pests Outside and Around Structures section of this label. For best results, make application during cool weather (65 to 80 degrees F) or in the early morning or late evening.

Mole Crickets: Make application late in the day and water grass immediately after application to move treatment down into the grass where these insects live. If soil is not moist, water before treatment also to bring crickets closer to surface before application. Treating at the time of peak egg hatch increases control of hatching nymphs. Frequent applications may be necessary to control larger nymphs later in the year.

Ticks: (including ticks that may transmit Lyme Disease and Rocky Mountain Spotted Fever) - Treat the entire area where ticks could be present. Begin treatments in the spring. Ticks may be re-introduced by host animals in the surrounding area. Retreats may be necessary to achieve and maintain control during periods of high pest pressure.

APPLICATION FOR ORNAMENTAL PESTS

Maxxthor SC controls listed insects and mites on trees, scrubs, foliage plants, non-bearing fruit and nut trees and flowers. Non-bearing trees are perennial plants that will not produce a harvestable agricultural commodity within the next 12 months. Maxxthor SC is not for use on plants being grown for sale, for commercial seed production or for research purposes.

Application Sites

For use on ornamental plants including trees, shrubs, ground covers, bedding plants and foliage plants being used for decorative or climate modification purposes. Plants on which use is permitted include those being grown in any type of setting or at any type of site not otherwise prohibited. Permitted sites include ornamental gardens, parks, landscapes, lawns, grounds and interior plantscapes.

Application Preparation

If necessary, test the effects of applications of different rates and volumes of mixed solution on a small number of a type of plant (with observations over one week to detect the occurrence of negative effects) before application of solution to large numbers of that type of plant.

Maxxthor SC can be mixed with other pesticides, including insect growth regulators. Follow the label directions of all the products mixed, making sure not to exceed the label rate of any individual product in the mixture. Any tank mixture that has not been tested before should be tested before full scale use by first mixing a small quantity of the mixture to ensure there is no physical or chemical incompatibility.

Application Methods

Apply solution in a quantity sufficient to wet all foliage. It may also be necessary to treat non-foliage parts of plants such as trunks and bark to control some pests based on where the pest may be located on the plant at a particular life stage.

If necessary, consult resources in horticulture in your area (such as your Cooperative Extension Service) to determine appropriate application timing, cultural practices and placement of treatment on different parts of plants to control different pests.

Reapplication

Reapplications may be necessary as plant growth occurs and new foliage appears or in the event of high pest pressure. Reapply as needed based on pest re-infestation. Reapplications must not occur more often than every 7 days.

Ornamental Application Use Rates and Volumes

Use rates for Maxxthor SC for ornamental pests are stated in fluid ounces of Maxxthor SC per 1000 square feet.

Ornamental application use rates based on the target pest, range between 0.125 to 1.0 fluid ounce of Maxxthor SC per 1,000 square feet. Recommended rates for specific pests within this range are given below. However, applications of up to 1.0 fluid ounce per 1000 square feet are permitted at the discretion of the applicator to control any pest.

The calculated amount of Maxxthor SC can be applied in any volume of water as long as the maximum label rate of 1.0 fluid ounce per 1000 feet is not exceeded. Do not exceed the maximum label rate by applying solution to an area smaller than intended when it was mixed and diluted unless such underdosing will not result in an application rate per 1000 square feet in excess of the maximum label rate.

Ornamental Pest Application Rates

The application rates listed below provide control of the listed pests under normal conditions. At the discretion of the applicator, up to 1 fluid ounce of Maxxthor SC per 1000 square feet can be used to control any of the listed pests. Use the higher application rate when maximum residual control is needed.

Use Rate Table for Maxxthor SC for Ornamental Applications	
Use Rate	Fluid ounces Maxxthor SC per 1,000 square feet (Range)
A	0.125⇒0.250
B	0.250⇒0.500
C	0.500⇒1.000

Ornamental Pests Grouped by Use Rates

Use Rate A: Bagworms, Cutworms, Elm Leaf Beetle, Fall Webworm, Gypsy Moth Caterpillar, Lace Bug, Leaf Feeding Caterpillar, Tent Caterpillar
Use Rate B: Ants, Aphids, Bees, Beet Armyworm, Black Vine Weevil (Adult), Brown Soft Scale, Broad Mite, Budworms, California Red Scale (Crawler), Centipedes, Citrus Thrip, Clover Mite, Crickets, Diaprepes (Adult), Earwig, European Red Mite, Flea Beetles, Fungus Gnat (Adult), Grasshoppers, Leafhoppers, Leafrollers, Mealybugs, Millipedes, Mites, Mosquitoes, Orchid Weevil, Pillbugs, Pine Needle Scale (Crawler), Plant Bugs (including Lygus spp.), San Jose Scale (Crawlers), Scorpions, Sowbugs, Spiders, Spittlebugs, Thrips, Tip Moth Twig Borers, Wasps, Weevils, Whiteflies
Use Rate B except not for use in California - Adelgids, Beetles, Cicadas, Japanese Beetle (adult), Psyllids, Spittlebugs, Treehoppers
Use Rate C: Imported Fire Ant, Leafminers, Pecan Leaf Scorch Mite, Pine Shoot Beetle (Adults), Spider Mites

Mixing Table for Maxxthor SC for Ornamental Applications				
Application Volume: Gallons/1000 sq. ft.	Use Rate: Fluid oz. / 1000 sq. ft.	Use these amounts of Maxxthor SC diluted to these volumes of finished spray		
		5 gal.	10 gal.	25 gal.
2	.125	.31	.63	1.56
	.250	.63	1.25	3.13
	.500	1.25	2.50	6.25
	1.000	2.50	5.00	12.50
5	.125	.13	.25	.63
	.250	.25	.50	1.25
	.500	.50	1.00	2.50
	1.000	1.00	2.00	5.00
10	.125	--	.13	.31
	.250	--	.25	.63
	.500	--	.50	1.25
	1.000	--	1.00	2.50

Calculating Amounts of Maxxthor SC to Mix for Ornamental Pests

To mix and apply any amount of Maxxthor SC for ornamental pests, determine:
A = Square feet of area to be treated /1000 (5,500 sq. ft./1000 = 5.5)
B = Use Rate per 1000 square feet for the target pest(s) in fluid ounces Maxxthor SC taken from the Use Rate Table. If treating for more than one pest, select the highest rate.
Calculate the amount of Maxxthor to mix for ornamental pests as follows:
Fluid Ounces Maxxthor SC to use = A X B
Mix this amount of Maxxthor SC in the amount of water needed to make the application. After mixing, the percent of Maxxthor SC active ingredient contained in the mixture can be determined using the following formula.
% a. i. = (0.0617 X fluid oz. of Maxxthor SC added) / gallons of water in container

Application Against Specific Ornamental Pests

Bagworms: Spray when bagworms first begin to hatch. Apply directly to the larvae. Treatment is most effective against young larvae.

STRUCTURAL PESTS (OTHER THAN WOOD INFESTING INSECTS)

Maxxthor SC controls a wide range of structural pests including nuisance pests inside and outside of structures Maxxthor SC can be applied in and around any type of residential or commercial structure, building or mode of transport including food/feed handling establishments unless otherwise prohibited. Permitted sites include but are not limited to the interior and exterior of homes, office buildings, mobile and modular homes, apartments and stores. Do not apply within aircraft cabins.

Follow additional application restrictions for residential outdoor surface and space sprays under DIRECTIONS FOR USE.

Dilution of Maxxthor SC for Structural Pests

For structural pests, use rates for Maxxthor SC are expressed and mixed according to the percentage (%) concentration solution it forms when mixed in water. Each 0.166 fluid ounce (1 teaspoon) of Maxxthor SC that is added to one gallon of water increases the concentration of Maxxthor SC in that one gallon of water by 0.01%. For example, to make a 0.06% solution in one gallon of water, mix 1 fluid ounce (6 teaspoons) of Maxxthor SC in one gallon of water. Use the conversion table and formulas below to determine the amount of Maxxthor SC to add to any quantity of water.

Mixing Table for Maxxthor SC for Structural Pests		
% to mix	Fluid ounces to add per gallon to mix this %	Teaspoons to add per gallon to mix this %
0.01	0.166	1
0.02	0.333	2
0.03	0.500	3
0.04	0.666	4
0.05	0.833	5
0.06	1.000	6

29.57 milliliters = 2 tablespoons = 6 teaspoons=1 Fluid ounce

Calculating Amount of Maxxthor SC to Mix for Structural Pests

Calculate the amount of Maxxthor SC to mix for structural pests as follows:
A = Volume of water, in gallons, into which the Maxxthor SC will be mixed. Express any partial gallons as decimal fractions (1/2 = .5)
B = Fluid ounces (or tablespoons) Maxxthor SC per gallon from Mixing Table. Select the desired % concentration based on the site of application and the pest(s) to be controlled. Read across to find amount of Maxxthor SC to add per gallon in fluid ounces or teaspoons.
Fluid ounces (or teaspoons) Maxxthor SC to mix = A X B
Mix this amount of Maxxthor SC in the predetermined amount of water (A). Food utensils such as teaspoons and measuring cups should not be used for food purposes after use with pesticides.

STRUCTURAL PESTS OUTSIDE AND AROUND STRUCTURES

Pests Controlled

Ants (including Fire Ant), Armyworms, Bees, Biting Flies, Boxelder Bug, Centipedes, Chiggers, Chinch Bug, Clover Mites, Crickets, Cutworms, Dichondra Flea Beetle, Earwigs, Elm Leaf Beetle, Firebrats, Fleas, Flies, Gnats, Grasshoppers, Hornets, Midges, Millipedes, Mosquitoes, Moths, Pillbugs, Roaches (including Cockroaches), Scorpions, Silverfish, Sod Webworms, Sowbugs, Spider Mites, Spiders (including Black Widow Spider), Springtails, Ticks (including Brown Dog Tick) and Wasps.

Pest controlled (but not for use against in California)

Beetles including Japanese Beetle

Application Methods

Apply Maxthor SC as a solution in the form of a general surface, spot, crack and crevice, pinstream or coarse spray. Do not apply as a space spray. May also be applied with a paintbrush.

Application Use Rates and Volumes

Use a 0.02% to 0.06% solution of Maxthor SC. Use a spray volume of up to 10 gallons of solution per 1,000 square feet. Higher application volumes may be used if necessary to sufficiently wet vegetation and landscaping with the spray solution.

Re-Application

Treatments must not be repeated more often than once every 7 days. The best efficacy and longest residual control is achieved when the highest concentration is used.

Application Locations

The following procedure must be followed to help achieve maximum control of the pest.

Apply spray to the exterior surfaces of structures and to grounds, lawns, landscaping, plants and hard surface areas adjacent to structures. Can also be applied to any areas where pests congregate or have been seen.

Treat non-porous surfaces only in areas protected from rainfall and spray from sprinklers with low volume applications at the rate of one gallon product diluted in water (see above dilution rates) per 1000 square feet.

Applications to vertical exterior surfaces (e.g., foundations) are permitted to a maximum height of 3 feet from ground level. Sections of vertical exterior surfaces that abut non-porous horizontal surfaces can only be treated if either 1) these sections are protection from rainfall and spray from sprinklers or 2) they do not drain into a sewer, storm drain, or curbside gutter (e.g., not to sections that abut driveways or sidewalks that drain into streets).

Perimeter Band Treatment

To help prevent pest infestation of structures, create a treated zone or band on the structure, soil and vegetation around the entire perimeter of a structure. Apply solution to all surfaces within a band beginning 6 to 10 feet from the exterior foundation of the structure that extends back to the structure and then continues 2 to 3 feet up the exterior surface of the structure from the ground. Application volume will depend upon the nature of the surface being treated. Mulch areas, for example, require more volume and hard surface areas requiring less.

*For sections of foundation that abut non-porous horizontal surfaces, the treated areas must be protected from rainfall and spray from sprinklers or they do not drain into a sewer, storm drain or curbside gutter (e.g., not to sections that abut driveways or sidewalks that drain into streets).

Control of Specific Structural Pests Outdoors

Ants (Nuisance ants other than Carpenter Ants) Outdoors: To achieve the highest level of control, locate and directly treat ant nests. Apply solution to ant trails, around doors and windows and at points where ants can be expected to forage or congregate.

For Ant (including Fire Ant) Mounds: Treat mounds and area within a 2 foot radius of the center of the mound with 1-2 gallons of a 0.06% solution. When mounds exceed 12 inches in size use the highest volume of solution. Applications in cool weather (65 to 80 degrees F) or the cooler parts of the day are most effective. Do not apply treatment during the heat of the day.

Bees, Wasps, Hornets, and Yellow Jackets: Always use the highest rate. Make application in late evening when insects are at rest at the nest for best results and to avoid stings. Spray nest openings in ground, in bushes and wherever insects may be nesting. Spray to the point of saturation. Spray as many insects as possible. Remove and destroy treated nests to prevent emergence of newly hatched insects.

Boxelder Bug: Apply directly to where insects have congregated at points of entry. It may be necessary to also treat trees.

Centipedes, Earwigs, Beetles, Millipedes, Pillbugs, Sowbugs: Thoroughly treat mulch areas. The reduction of moisture in mulch areas may also aid in control of these pests.

Mosquitoes: Apply solution to lawns and landscaping, under decks and to building foundations. Refer to *APPLICATIONS FOR ORNAMENTAL PESTS* section for mixing and applying large amounts of solution against mosquitoes.

Fleas and Ticks: Treat the entire area where insects could be present. Begin treatments in the spring. Insects may be re-introduced by host animals in the surrounding area. Retreatment may be necessary to achieve and maintain control during periods of high pest pressure.

STRUCTURAL PESTS INSIDE STRUCTURES (OTHER THAN FOOD/FEED HANDLING ESTABLISHMENTS) INCLUDING MODES OF TRANSPORT

Pests Controlled

Ants, Bedbug, Bees, Beetles, Boxelder Bug, Carpet Beetles, Centipedes, Clothes Moth, Cockroaches, Crickets, Earwigs, Firebrats, Flies, Gnats, Midges, Millipedes, Pillbugs, Scorpions, Sowbugs, Silverfish, Spiders, Ticks and Wasps

Application Preparation

When treating in home food preparation and storage areas, cover all food preparation surfaces and utensils prior to beginning treatment. Surfaces or items that cannot be covered or removed should be thoroughly washed after treatment and before use. Food that cannot be covered should be removed. Before application, remove pets, birds and cover and disconnect aquariums. Do not allow humans and pets to touch treated surfaces until they have dried. During any application to overhead areas of structure, cover surfaces below with plastic sheeting or similar material, except for soil surfaces in crawlspaces. Wear protective clothing, unvented goggles, gloves and respirator when making an overhead application or when applying in poorly ventilated indoor areas.

Application Use Rates

Use a 0.02% to 0.06% solution of Maxthor SC.

Application Methods

Apply Maxthor SC as a solution in the form of a general surface, spot, crack and crevice, pinstream or coarse spray. Maxthor SC may also be applied with a paintbrush. Do not apply as a space spray.

Maxthor SC may be converted to foam and used to treat structural voids. First form a solution of Maxthor SC of the appropriate percentage concentration and volume. Then add recommended volume of a compatible foaming agent. Verify that the foaming agent is compatible with Maxthor SC.

Application Locations

Apply to and around any areas pests or their evidence is seen or found, could hide or rest or could enter the premises including cracks and crevices, behind and under cabinets and appliances, around doors and windows, in attics and storage areas. Spot treatments to floor or rugs beneath furniture are permitted but do not apply to entire floor area.

Control of Specific Structural Pests Indoors

Cockroaches, Crickets, Firebrats, Scorpions, Silverfish, Spiders, and Ticks: Apply wherever these pests may hide, enter the structure or congregate such as cracks and crevices, baseboards, water pipe openings, around doors and windows, behind and under cabinets and appliances and within storage and attic areas.

Bedbug: To aid in control, apply to cracks and crevices wherever evidence of bedbugs has been found including within empty dressers and closets, on bed frames and box springs and behind high and low wall moldings and wallpaper edges. Do not apply to bed linens, blankets, pillows, mattresses or clothes. Remove all clothes and other articles from dressers or closets before application within them. Not recommended for use as a sole control agent against bedbugs. If evidence of bedbugs is found on or in mattresses, use products approved for application to this item.

Boxelder Bug, Centipedes, Earwigs, Beetles, Millipedes, Pillbugs, Sowbugs: Apply to points of pest entry into structure such as around windows and doors. Apply anywhere else pests may congregate or be located.

Bees, Wasps, Hornets, Yellow-Jackets: Always use the highest rate. Apply solution to hiding and breeding places, contacting as many insects as possible. Best results are achieved when application is made in the evening when insects are at rest and to avoid stings. Spray to the point of saturation. Spray as many insects as possible. Remove and destroy treated nests to prevent emergence of newly hatched insects.

Ants (Nuisance ants other than Carpenter Ants) Indoors: To achieve the highest level of control, locate and directly treat ant nests and ant trails. Apply solution in areas infested by or expected to be infested by ants

CONTROL OF STRUCTURAL PESTS WITHIN FOOD/FEEDAREAS OF FOOD/FEED HANDLING ESTABLISHMENTS

Food/feed handling establishments are defined as places other than private residences in which exposed food/feed is held, processed, prepared or served. This includes areas for receiving, storing, packing (canning, bottling, wrapping, boxing) food. Also included are areas of edible waste storage and enclosed processing systems (mills, dairies, edible oils, syrups) of food. Serving areas where food is exposed and the facility is in operation are also considered food areas.

Application Location

Applications may be made to both food/feed and nonfood areas of food/feed handling establishments.

Application Methods

Maxthor SC can be applied within food/feed handling establishments in the form of a general surface, spot or crack and crevice spray.

General Surface Application

Do not use this application method when facility is in use or when foods are exposed. Do not apply solution directly to food. Cover or remove all food processing and/or handling equipment before application. After application in areas where food is commercially prepared or processed, wash all equipment, benches, shelving and other surfaces which food will contact. Rinse thoroughly with fresh, clean water. Clean food handling or processing equipment and thoroughly rinse with clean, fresh water.

Spot and Crack and Crevice Application

Spot or crack and crevice applications may be made while the facility is in operation. Food should be covered or removed from area being treated. Do not apply directly to food or food handling surfaces.

ATTENTION

Remove or cover and disconnect aquariums during application.

Do not apply a broadcast application to interior surfaces of living areas.

Do not apply to pets, crops, sources of electricity or firewood.

During any application to overhead areas of structure, cover surfaces below with plastic sheeting or similar material, except for soil surfaces in crawlspaces.

Do not allow dripping or runoff to occur during indoor applications.

Do not allow spray to contact food, foodstuffs, food contacting surfaces, food utensils or water supplies.

Thoroughly wash dishes and food handling utensils with soap and water if they become contaminated by application of this product.

Do not allow people or pets to touch or walk on treated surfaces until spray has dried.

Do not apply this product in nursing home or patient rooms or in any rooms while occupied by the elderly or infirmed.

Do not apply in classrooms when they are in use.

Do not apply when occupants are present in the immediate area in institutions such as health care facilities, libraries, schools, offices, etc.

Do not apply in livestock buildings such as barns.

Maxthor SC will not stain or damage any surface that water alone will not stain or damage.

IMPORTANT READ BEFORE USE

NOTICE: Read the entire Directions for Use, Conditions of Sale, Disclaimer of Warranties and Limitations of Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

CONDITIONS OF SALE: The Directions for Use of this product are believed to be adequate and must be followed carefully. However, because of manner of use and other factors beyond the control of Ensysyex II, Inc., it is impossible for Ensysyex II to eliminate all risks associated with the use of this product such as ineffectiveness or unintended consequences. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold Ensysyex II harmless for any claims relating to such factors.

DISCLAIMER OF WARRANTIES: Seller warrants that this product conforms to the chemical description in the label and is reasonably fit for the purposes stated on the Directions for Use when used in accordance with the Directions for Use under normal conditions of use. ENSYSTEX II MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE, NOR ANY OTHER EXPRESS OR IMPLIED WARRANTIES WITH RESPECT TO THE SELECTION, PURCHASE, OR USE OF THIS PRODUCT THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. Any warranties, express or implied, having been made are inapplicable if this product has been used contrary to label instructions, under abnormal conditions or under conditions not reasonably foreseeable by (or beyond the control of) seller or Ensysyex II, Inc., and buyer assumes the risk of any such use.

LIMITATIONS OF LIABILITY: To the extent consistent with applicable law, Ensysyex II shall not be liable for any incidental, consequential or special damages resulting from the use or handling of this product. THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF ENSYSTEX II AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF ENSYSTEX II, THE REPLACEMENT OF THE PRODUCT.

This Conditions of Sale and Limitation of Warranty and Liability may not be amended by any oral or written agreement.

Maxthor is a registered trademark of Ensysyex II, Inc.

04/13

Safety Data Sheet

Maxxthor SC

Emergency Phone 1-800-424-9300 (Chemtrec)

1. PRODUCT AND COMPANY IDENTIFICATION

Product Name: Maxxthor SC
Chemical Name: Bifenthrin
Recommended use of the chemical and restrictions on use: Insecticide

Company: Ensystex II, Inc.
Address: 2175 Village Dr., Fayetteville, NC 28304
Daytime Phone: 1-800-398-3772

2. HAZARDS IDENTIFICATION

Hazard classification

This material is hazardous under the criteria of the Federal OSHA Hazard Communication Standard 29CFR 1910.1200
Acute toxicity - Category 4 - Oral
Acute toxicity - Category 4 - Inhalation (Dusts/Mists)
Carcinogenicity - Category 2
Specific target organ - single exposure - Category 1
Specific target organ toxicity - repeated exposure - Category 1

Label Elements

Hazard pictograms



Signal Word: DANGER!

Hazards

H302 - Harmful if swallowed
H332 - Harmful if inhaled
H351 - Suspected of causing cancer
H370 - Causes damage to organs
H372 - Causes damage to organs through prolonged or repeated exposure

Precautionary Statements

Prevention

P202 - Do not handle until all safety precautions have been read and understood
P281 - Use personal protective equipment as required
P264 - Wash face, hands and any exposed skin thoroughly after handling
P270 - Do not eat, drink or smoke when using this product
P260 - Do not breathe dust/fume/gas/mist/vapors/spray

Response

P321 - Specific treatment (see supplemental first aid instructions on the label)
P308 + P311 - If exposed or concerned: Call a POISON CONTROL CENTER or doctor
P304 + P340 - If INHALED: Remove person to fresh air and keep comfortable for breathing
P312 - Call a POISON CONTROL CENTER or doctor if you feel unwell
P301 + P312 - IF SWALLOWED: Call a POISON CONTROL CENTER or doctor if you feel unwell
P330 - Rinse mouth

Storage

P405 - Store locked up.

Disposal

P501 - Dispose of contents / container to an approved waste disposal facility.

Other hazards

No data available

Other Information

Very toxic to aquatic life with long lasting effects

3. COMPOSITION / INFORMATION ON INGREDIENTS

Bifenthrin CAS # 82657-04-3 Weight % - 7.9
Propylene Glycol CAS # 57-55-6

4. FIRST-AID

Description of first-aid measures

General advice: First-Aid responders should pay attention to self-protection and use the recommended protective clothing (chemical resistant gloves, splash protection). If potential for exposure exists, refer to Section 8 for specific personal protective equipment.

Inhalation: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. If the person is not breathing and has no pulse, consider cardiopulmonary resuscitation (CPR); use pocket resuscitation mask, bag valve mask etc., to avoid risk of poisoning rescuer. Consult a physician in all cases.

Eye Contact: Hold eye open and rinse slowly and gently with water for at least 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a Poison Control Center or doctor immediately for treatment advice.

Skin Contact: Take off Contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a Poison Control Center or doctor immediately for treatment advice.

Ingestion: Call a Poison Control Center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the Poison Control Center. Never give anything by mouth to an unconscious person.

Most important symptoms and effects, both acute and delayed: Central nervous system effects.

Indication of any immediate medical attention and special treatment needed

Treat symptomatically. This product is a pyrethroid. If large amounts have been ingested, the stomach and intestines should be evacuated. Treatment is symptomatic and supportive. Digestible fats, oils, or alcohol may increase absorption and so should be avoided.

5. FIRE-FIGHTING MEASURES

Extinguishing Media: Foam. Carbon dioxide (CO₂). Dry chemical. Soft stream or water fog only if necessary.

Advice for firefighters

Explosion Data

Sensitivity to Mechanical Impact Not sensitive.
Sensitivity to Static Discharge Not sensitive

Special protective equipment for firefighters: Wear positive-pressure self-contained breathing apparatus (SCBA) and protective fire fighting clothing (includes fire fighting helmet, coat, trousers, boots and gloves). If protective equipment is not available or not used, fight fire from a protected location or safe distance.

6. ACCIDENTAL RELEASE MEASURES

Personal Precautions: Isolate and post spill area. Wear appropriate safety clothing, respiratory protection devices and eye/face protection (see Section 8). Evacuate unprotected personnel that are nearby.

Environmental precautions: Keep people and animals away from and upwind of spill or leak. Prevent from entering into soil, ditches, sewers, waterways and /or groundwater. See Section 12, Ecological Information.

Methods and materials for containment and cleanup:

Small spills: knock down and dilute vapors with water fog or spray. Apply vapor suppression foams until spill can be cleaned up. Use non-sparking tools in cleanup operations. Large spills: Contact Ensysystex II, Inc. for cleanup assistance. See Section 13, Disposal considerations, for additional information.

7. HANDLING AND STORAGE

Handling: Use good personal hygiene. Do not contaminate other pesticides, fertilizers, water, food or feed by storage or disposal.

Storage: Keep out of reach of children. Product should be stored in compliance with local regulations. Store in a well ventilated, cool, dry area. Keep away from heat sources. Store in original container.

Incompatible products: None known

8. EXPOSURE CONTROLS/PERSONAL PROTECTION

Applicators should refer to the product label for personal protection equipment requirements during application.

Component	Regulation	Type of listing	Value/Notation
Propylene glycol 57-55-6	Safe Work Australia Standards for Airborne Contaminants	8 hr TWA vapor and particulates	474 mg/m ³ (150 ppm)
	Safe Work Australia Standards for Airborne Contaminants	8 hr TWA particulates only	10 mg/m ³

These recommendations are for Manufacturing. Applicators should see the product label for proper personal protective equipment.

Engineering measures: Apply technical measures to comply with the occupational exposure limits.

Exposure controls

Provide general and/or local exhaust ventilation to control airborne levels below the exposure limits.

Respiratory Protection: Atmospheric levels should be maintained below the exposure guidelines. When respiratory protection is required or during emergency conditions, use a NIOSH/MSHA approved respiratory protection.

Hand/Skin Protection: Wear long-sleeved shirt, long pants, socks, protective gloves and shoes

Eye/Face Protection: Chemical proof goggles / face shield

9. PHYSICAL AND CHEMICAL PROPERTIES

Appearance/Odor: Beige liquid, Mild odor
pH: 6.7
Flash Point/Range: > 65.5°C / > 149.9°F Tag Closed Cup
Water solubility: Dispersible in water
Vapor pressure: 15.2 atmospheres at @ 68 °F (20 °C)
Specific Gravity: 1.024 @ 20°C
Bulk Density: 8.53 lb/gal
Flammability (solid, gas): No information available
Flammability limit in Air
 Upper flammability limit: No information available
 Lower flammability limit: No information available
Vapor pressure: No information available
Vapor density: No information available
Density: No information available
Solubility in other solvents: No information available
Partition coefficient: No information available
Autoignition temperature: No information available
Decomposition temperature: No information available

10. STABILITY AND REACTIVITY

Reactivity: None under normal use conditions
Chemical Stability: Stable under normal storage conditions.
Conditions to Avoid: Heat, flames and sparks.
Materials to Avoid: Strong bases.
Hazardous Decomposition Products: Carbon oxides (CO_x), hydrogen chloride, hydrogen fluoride, chlorine, fluorine
Additional Information: Hazardous polymerization will not occur.

11. TOXICOLOGICAL INFORMATION

Acute Toxicity: Inhalation LC50/Rat/ 2.895 mg/L 4 hr (rat) Estimated; Oral LD50/Rat/632mg/kg; Dermal LD50/Rabbit/>2000mg/kg
Irritation: Reacts with mucous membranes
Chronic Toxicity: Inhalation, after repeated exposure, various species,

Skin corrosion/irritation

Non-irritating to skin

Serious eye damage/eye irritation

Non-irritating to eye

Sensitization

Non-sensitizing

Information on toxicological effects

Symptoms: large doses of bifenthrin ingested by laboratory animals produced signs of toxicity including convulsions, tremors and bloody nasal discharge.

Delayed and immediate effects as well as chronic effects from short and long-term exposure

Chronic toxicity: Bifenthrin: Long-term exposure caused neurotoxicity (tremors and impaired gait) in the early exposure in animal studies, but tremors disappeared with continued exposure

Carcinogenicity: Bifenthrin: weak treatment-related response for liver adenocarcinomas and benign bladder tumors (lesions) in male mice.

Neurological effects: Bifenthrin causes clinical signs of neurotoxicity (tremors, impaired gait, excessive salivation) following acute or subchronic exposure. Tremors disappeared with continued exposure.

No teratogenic effect

No carcinogenic effects

No reproductive toxicity effects

No mutagenic effects

Target organ effects: Bifenthrin: Central Nervous System

Neurological effects: Bifenthrin causes clinical signs of neurotoxicity (tremors, impaired gait, excessive salivation) following acute or subchronic exposure. Tremors disappeared with continued exposure.

Aspiration hazard: No information available

12. ECOTOXICOLOGICAL INFORMATION

Toxicity

Acute toxicity to fish

LC₅₀, 96 Hour, 0.1 µg/l

Acute toxicity to aquatic crustacea

EC₅₀, 48 Hour, 0.11 µg/l

Acute toxicity to algae/aquatic plants

EC₅₀ 72 Hour 0.822 mg/L

No Observed Effect Level Aquatic (NOEC)

Fish 21d 0.012 µg/l

Crustacea 21d 0.0013 µg/l

Persistence and degradability

Bifenthrin: Moderately persistent. Does not readily hydrolyze. Not readily biodegradable.

Bioaccumulative potential

Bifenthrin: Has a potential for bioaccumulation

Mobility in soil

Bifenthrin: Immobile. Not expected to reach groundwater.

13. DISPOSAL CONSIDERATIONS

Waste disposal methods: Improper disposal of excess pesticide, spray mixture, or rinsate is prohibited. If these wastes cannot be disposed of by use according to the label instructions, contact appropriate authorities for guidance.

Contaminated Packaging: Containers must be disposed of in accordance with local, state and federal regulations. Refer to product label for container disposal instructions.

14. TRANSPORT INFORMATION

DOT This material is not a hazardous material as defined by U.S. Department of Transportation 49 CFR Parts 100 - 185.

TDG: Classification below is only applicable when shipped by vessel and is not applicable when shipped by road or rail only.

UN #: UN2191

Proper Shipping Name: Environmentally hazardous substance, liquid, n.o.s.

Hazard Class: 9

Packing Group: III

Marine Pollutant: Bifenthrin

Description: UN3802, Environmentally hazardous substance, liquid, n.o.s. (Bifenthrin), 9 PG

Classification for SEA transport (IMO-IMDG):

Proper Shipping Name: Environmentally hazardous substance, liquid, n.o.s.

UN #: UN3082

Hazard Class: 9

Packing group: III

EmS No.: F-A, S-F

Marine pollutant: Bifenthrin

Description: UN3082, Environmentally hazardous substance, liquid, n.o.s. (Bifenthrin), 9 PG III

Classification for AIR transport (IATA/ICAO):

UN#: UN3082

Proper Shipping Name: Environmentally hazardous substance, liquid, n.o.s.

Hazard Class: 9

Packing group: III

Limited quantity: 30 kg G

15. REGULATORY INFORMATION

The information herein is given in good faith, but no warranty, expressed or implied, is made. Consult Ensystex II for further information.

US Federal Regulations

SARA 313

Section 313 of Title III of the Superfund Amendments and Reauthorization Act of 1986 (SARA). This product contains a chemical or chemicals which are subject to the reporting requirements of the Act and Title 40 of the Code of Federal Regulations, Part 372:

Chemical Name	CAS#	Weight %	SARA 313 Threshold Values %
Bifenthrin	82657-04-3	7.9	1.0

Superfund Amendments and Reauthorization Act of 1986 Title III (Emergency Planning and Community Right-to-Know Act of 1986) Sections 311 and 312

Acute health hazard Yes

Chronic health hazard Yes

Fire hazard No

Sudden release of pressure hazard No

Reactive hazard No

Clean Water Act

This product does not contain any substances regulated as pollutants pursuant to the Clean Water Act (40 CFR 122.21 and 40 CFR 122.42)

CERCLA

This material, as supplied, does not contain any substances regulated as hazardous substances under the Comprehensive Environmental Response Compensation and Liability Act (CERCLA) (40 CFR 302) or the Superfund Amendments and Reauthorization Act (SARA) (40 CFR 355). There may be specific reporting requirements of the local, regional, or state level pertaining to releases of this material.

FIFRA Information

This chemical is a pesticide product registered by the Environmental Protection Agency and is subject to certain labeling requirements under federal pesticide law. These requirements differ from the classification criteria and hazard information required for safety data sheets, and for workplace labels of non-pesticide chemicals. Following is the hazard information as required on the pesticide label:

Caution

Signal Word:

16. Other Information

Hazard rating System

NFPA

Health	Fire	Reactivity	Specific Hazard
1	0	0	None

HMIS

Health	Fire	Reactivity	
2	0	0	Protective Equipment: B

Disclaimer: The information of this SDS is based on the present state of our knowledge. The user's working conditions however, are beyond our knowledge and control. The product is not to be used for purposes other than those specified under section 1 without a written permission. It remains the responsibility of the user to ensure that the necessary steps are taken to meet the laws and regulations. Handling of the product may only be done by people above 18 years of age, who are satisfactorily informed of how to do the work, the hazardous properties and necessary safety precautions. The information given in this SDS is to describe the product only in terms of health and safety requirements and should not, therefore, be construed as guaranteeing specific properties.

Revised 07/21/2015