

□ **Objective Set 1: Classifying Numbers as Real, Rational, Irrational, Integer, Whole or Natural**

[Khan Academy Number Sets](#)

Circle each set of numbers that each number belongs to. Circle ALL that apply.

a.) $\sqrt{25}$

Real

Rational

Irrational

Integer

Whole

Natural

b.) -3.14

Real

Rational

Irrational

Integer

Whole

Natural

□ **Objective Set 2: Computing basic number operations (add, subtract, multiply, divide – with FRACTIONS!)**

[Khan Academy Order of Operations](#)

Simplify each of the expressions below.

a.) $\frac{3}{4} - \frac{7}{9} + 2$

$LCD = 36$

$$\frac{27}{36} - \frac{28}{36} + \frac{72}{36} = \boxed{\frac{71}{36}}$$

b.) $\frac{3}{7} \div \frac{4}{14}$

$$\frac{3}{7} \cdot \frac{14}{4} = \boxed{\frac{3}{2}}$$

c.) $-92 + (-19) + 82$

$$\boxed{-29}$$

d.) $\left(\frac{-1}{5}\right)^2 = \boxed{\frac{1}{25}}$

e.) $\frac{3-9}{3+7}$

$$\frac{-6}{10} = \boxed{-\frac{3}{5}}$$

f.) $7(2) - (3+5)^2 \div 4 - 2$

$$14 - 8^2 \div 4 - 2$$

$$14 - 64 \div 4 - 2$$

$$14 - 16 - 2$$

$$-2 - 2$$

$$\boxed{-4}$$

□ **Objective Set 3: Simplifying expressions.**

[Khan Academy Distributive Property](#)

[Khan Academy Multiplying Polynomials](#)

[Khan Academy Exponent Products](#)

[Khan Academy Exponent Quotients](#)

[Khan Academy Exponent Properties Part 3](#)

Simplify each of the expressions below.

a.) $3(x - 4x^2) + 3x - 4(2x + 1)$

$$3x - 12x^2 + 3x - 8x - 4$$

$$\boxed{-12x^2 - 2x - 4}$$

b.) $7 - 2(4ab - 3a) + 7(2b - 4ab)$

$$7 - 8ab + 6a + 14b - 28ab$$

$$\boxed{-36ab + 6a + 14b + 7}$$

c.) $(3m - 4b)^2$

$$(3m - 4b)(3m - 4b)$$

$$\boxed{9m^2 - 24mb + 16b^2}$$

d.) $-2(3x + 4y - 8xy)$

$$\boxed{-6x - 8y + 16xy}$$

e.) $(2x - 5)(3x^2 + 4x - 7)$

$$6x^3 + 8x^2 - 14x - 15x^2 - 20x + 35$$

$$\boxed{6x^3 - 7x^2 - 34x + 35}$$

f.) $\frac{3xy^2}{21x^4yz^{-4}}$

$$\boxed{\frac{1yz^4}{7x^3}}$$

g.) $\left(\frac{-14a^2bc^3}{2a^4b^2c}\right)\left(\frac{3a}{7c^3}\right)$

$$\boxed{\frac{-3}{abc}}$$

h.) $(3mx^2)^{-1}$

$$\boxed{\frac{1}{3mx^2}}$$

i.) $\left(\frac{-2x^2y^{-3}}{4x^3y^2}\right)^2$

$$\left(\frac{-1}{2x^5y^5}\right)^2$$

$$\boxed{\frac{1}{4x^{10}y^{10}}}$$

□ **Objective Set 4: Evaluate expressions.**

[Khan Academy Evaluate Expressions](#)

Evaluate each expression below for the given values.

a.) $(2x-3)^2 - 4x + 7$ for $x = -2$

$$\begin{array}{r} 4x^2 - 12x + 9 \\ -4x + 7 \\ \hline 4x^2 - 16x + 16 \end{array}$$
$$4(-2)^2 - 16(-2) + 16$$
$$16 + 32 + 16 = \boxed{64}$$

c.) $|2-x| + 8$ for $x = 5$

$$|2-5| + 8$$
$$|-3| + 8$$
$$3 + 8 = \boxed{11}$$

b.) $\left(\frac{3x^2}{-y}\right) + z$ for $x = -1, y = 2, z = 5/3$

$$\left(\frac{3(-1)^2}{-2}\right) + \frac{5}{3}$$
$$\left(\frac{3 \cdot 1}{-2}\right) + \frac{5}{3}$$
$$\left(\frac{-3}{2}\right) + \frac{5}{3} = \boxed{\frac{1}{6}}$$

□ **Objective Set 5: Solving equations and inequalities – then graphing the solution on a number line.**

[Khan Academy Solving Linear Equations](#)

[Khan Academy Solving Linear Inequalities](#)

Solve the following. Graph when indicated.

a.) $2x - 5 = 7$


$$\begin{array}{r} 2x - 5 = 7 \\ +5 +5 \\ \hline 2x = 12 \\ \boxed{x = 6} \end{array}$$

b.) $\frac{9}{5} = \frac{a-1}{2}$

$$18 = 5(a-1)$$
$$18 = 5a - 5$$
$$23 = 5a$$
$$\boxed{a = \frac{23}{5}}$$

c.) $2(b-4) + 8 = 3b + 7$

$$2b - 8 + 8 = 3b + 7$$
$$2b = 3b + 7$$
$$-b = 7$$
$$\boxed{b = -7}$$




d.) $3|x| - 4 = 11$

$3|x| = 15$

$|x| = 5$


$x = 5, x = -5$



e.) $3x - 4 < 7$

$3x < 11$

$x < \frac{11}{3}$



f.) $2(x - 4) + 3(4 - 5x) = 25 - 7(3 + 2x) + x$

$2x - 8 + 12 - 15x = 25 - 21 - 14x + x$

$-13x + 4 = -13x + 4$

all real numbers $\mathbb{R}$


g.) $\left(\frac{1}{2}(4x - 5) + 3\right)^2 \geq (-6)^2$

$4x - 5 + 6 \geq -12$

$4x + 1 \geq -12$

$4x \geq -13$

$x \geq -\frac{13}{4}$




□ **Objective Set 6: Graphing equations of lines in standard form, slope-intercept form, point-slope form and horizontal/vertical form.**

[Khan Academy Graphing a Line in Slope Intercept Form](#)

[Khan Academy Converting to Slope Intercept Form](#)

Graph each line on the grid provided.


a.) $x = -2$



b.) $y = \frac{3}{2}x - 4$

$m = \frac{3}{2}$


$b = (0, -4)$



c.) $y + 7 = -2(x - 3)$


$(3, -7)$

$m = -2$




d.) $3x - 4y = 12$

$(4, 0)$
 $(0, -3)$




e.) $3(x - 4) = 3y + 6$

$3x - 12 = 3y + 6$
 $3x - 3y = 18$
 $(6, 0) \quad (0, -6)$



f.) parallel to the x-axis
through $(2, -6)$



□ **Objective Set 7: Writing equations of lines in slope-intercept and point-slope form.**

[Khan Academy Writing Equations of Lines](#)

Write an equation of the line for the given information.

a.) goes through $(2, 3)$ and $(-4, 6)$

$m = \frac{6-3}{-4-2}$
 $m = \frac{3}{-6}$
 $m = -\frac{1}{2}$

$y - 3 = -\frac{1}{2}(x - 2)$
 $y - 3 = -\frac{1}{2}x + 1$
 $+3 \quad +3$
 $y = -\frac{1}{2}x + 4$

b.) has a slope of $-\frac{3}{4}$ and y-intercept of 5

$y = -\frac{3}{4}x + 5$


c.) that models this relationship

Time(secs)	0	1	2	3
Distance(feet)	15	25	35	45

$y = 10x + 15$

$x = \text{time}$
 $y = \text{distance}$

d.)



$m = -\frac{1}{3}$

$b = (0, 0)$

$y = -\frac{1}{3}x$

e.) Is perpendicular to $y = \frac{3}{4}x + 5$ and goes through $(-9, 6)$

$$m = \frac{4}{3}$$

$$\begin{aligned} y - 6 &= \frac{4}{3}(x + 9) \\ y - 6 &= \frac{4}{3}x + 12 \\ y &= \frac{4}{3}x + 18 \end{aligned}$$

$$y = \frac{4}{3}x + 18$$

□ **Objective Sets 8 & 9: Recognizing whether a relation is a function and indicating its domain and range.**

[Khan Academy Recognizing a Function and its Domain and Range](#)

Indicate whether or not the relation is a function. If the relation is a function, identify the Domain and Range.

a.)

X	-2	-2	-2	-2
F(x)	3	5	7	9

no, -2 has more than 1 output


b.) $(-2, 3) (3, 4) (4, 3) (5, 6)$

yes, every input has 1 output

$$D: \{-2, 3, 4, 5\}$$

$$R: \{3, 4, 6\}$$

c.)



No - fails the vertical line test

d.) $h(x) = 2|x|$

yes. Every x has one y value

$$D: \mathbb{R} \text{ or } \mathbb{R}^1$$

$$R: y \geq 0$$

□ **Objective Set 10: Factoring Polynomials using the Greatest Common Factor, Difference of Two Squares, Lead Coefficient of One OR a combination of two of these ☺**

[Khan Academy GCF Factoring](#)

[Khan Academy Factoring Strategies](#)

Factor each of the following completely.

a.) $3xy - 15x^2y^3$

$$3xy(1 - 5xy^2)$$

b.) $49x^2 - 25y^2$

$$(7x - 5y)(7x + 5y)$$

c.) $x^2 - 6x - 55$

$$(x - 11)(x + 5)$$

d.) $a^4 + a^2 - 6$

$$(a^2 + 3)(a^2 - 2)$$

e.) $2y^2 + 30y + 112$

$$2(y^2 + 15y + 56)$$

$$2(y + 7)(y + 8)$$

f.) $a^4 - 81$

$$(a^2 - 9)(a^2 + 9)$$

$$(a - 3)(a + 3)(a^2 + 9)$$

g.) $3x^2 + 75$

$$3(x^2 + 25)$$

h.) $4a^4 - 16a^3 + 18a^2$

$$4a^2(a^2 - 4a + 4.5)$$

□ **Objective Set 11: Solving Right Triangles using right triangle trigonometry.**


[Khan Academy 45-45-90 Triangle Ratios](#)

[Khan Academy 30-60-90 Triangle Ratios](#)

[Khan Academy Triangle Trigonometry](#)

Find the missing angle or side length using properties of right triangles.

a.)




$$\sin 45 = \frac{x}{20}$$

$$\frac{\sqrt{2}}{2} = \frac{x}{20}$$

$$\boxed{x = 10\sqrt{2}}$$

b.)




$$\cos 30^\circ = \frac{x}{74}$$

$$74 \cos 30^\circ = x$$

$$74 \cdot \frac{\sqrt{3}}{2} = x$$

$$\boxed{37\sqrt{3} = x}$$

c.)



$$\tan \theta = \frac{8}{3}$$

$$\theta = \tan^{-1}\left(\frac{8}{3}\right)$$

This answer requires
a calculator!

$$\boxed{\theta = 69.44^\circ}$$