

Kingsway Regional School District
 Anaphylaxis/Severe Allergy Action Plan and Doctor's Order Sheet
 Valid for 1 School Year: _____

Name: _____ Grade: _____ Date of Birth: _____

Previous episode of Anaphylaxis Yes No Asthmatic Yes* No *Higher risk for severe reaction

Student has been trained and is capable of self-administration of the medication(s) named below.

Student IS NOT trained to self-administer the medication named below.








I give approval for my child to self-administer this medication at school, and I understand that under NJ state law, a trained delegate will be assigned to administer epinephrine to my child in the absence of a school nurse, or if my child is unable to self-administer the medication. I also understand that a delegate may not give antihistamines. In the absence of a school nurse, any antihistamine order will be disregarded and a trained delegate will administer epinephrine.

Extremely reactive to the following allergens: _____

THEREFORE:

- If checked, give epinephrine immediately if the allergen was **LIKELY** eaten, for **ANY** symptoms.
- If checked, give epinephrine immediately if the allergen was **DEFINITELY** eaten, even if no symptoms are apparent.





**FOR ANY OF THE FOLLOWING:
SEVERE SYMPTOMS**

 LUNG Shortness of breath, wheezing, repetitive cough	 HEART Pale or bluish skin, faintness, weak pulse, dizziness	 THROAT Tight or hoarse throat, trouble breathing or swallowing	 MOUTH Significant swelling of the tongue or lips
 SKIN Many hives over body, widespread redness	 GUT Repetitive vomiting, severe diarrhea	 OTHER Feeling something bad is about to happen, anxiety, confusion	OR A COMBINATION of symptoms from different body areas.

↓ ↓ ↓

1. **INJECT EPINEPHRINE IMMEDIATELY.**
2. **Call 911.** Tell emergency dispatcher the person is having anaphylaxis and may need epinephrine when emergency responders arrive.
 - Consider giving additional medications following epinephrine:
 - » Antihistamine
 - » Inhaler (bronchodilator) if wheezing
 - Lay the person flat, raise legs and keep warm. If breathing is difficult or they are vomiting, let them sit up or lie on their side.
 - If symptoms do not improve, or symptoms return, more doses of epinephrine can be given about 5 minutes or more after the last dose.
 - Alert emergency contacts.
 - Transport patient to ER, even if symptoms resolve. Patient should remain in ER for at least 4 hours because symptoms may return.

MILD SYMPTOMS

 NOSE Itchy or runny nose, sneezing	 MOUTH Itchy mouth	 SKIN A few hives, mild itch	 GUT Mild nausea or discomfort
--	--	--	--

FOR MILD SYMPTOMS FROM MORE THAN ONE SYSTEM AREA, GIVE EPINEPHRINE.

FOR MILD SYMPTOMS FROM A SINGLE SYSTEM AREA, FOLLOW THE DIRECTIONS BELOW:

1. Antihistamines may be given, if ordered by a healthcare provider.
2. Stay with the person; alert emergency contacts.
3. Watch closely for changes. If symptoms worsen, give epinephrine.

MEDICATIONS/DOSES

Epinephrine Brand or Generic: _____

Epinephrine Dose: 0.1 mg IM 0.15 mg IM 0.3 mg IM

Antihistamine Brand or Generic: _____

Antihistamine Dose: _____


Other (e.g., inhaler-bronchodilator if wheezing): _____

Parent/Guardian's Signature: _____ Date: _____

Physician Stamp and Signature: _____ Date: _____


HOW TO USE AUVI-Q® (EPINEPHRINE INJECTION, USP), KALEO

1. Remove Auvi-Q from the outer case.
2. Pull off red safety guard.
3. Place black end of Auvi-Q against the middle of the outer thigh.
4. Press firmly until you hear a click and hiss sound, and hold in place for 2 seconds.
5. Call 911 and get emergency medical help right away.




HOW TO USE EPIPEN® AND EPIPEN JR® (EPINEPHRINE) AUTO-INJECTOR AND EPINEPHRINE INJECTION (AUTHORIZED GENERIC OF EPIPEN®), USP AUTO-INJECTOR, MYLAN AUTO-INJECTOR, MYLAN

1. Remove the EpiPen® or EpiPen Jr® Auto-Injector from the clear carrier tube.
2. Grasp the auto-injector in your fist with the orange tip (needle end) pointing downward.
3. With your other hand, remove the blue safety release by pulling straight up.
4. Swing and push the auto-injector firmly into the middle of the outer thigh until it 'clicks'.
5. Hold firmly in place for 3 seconds (count slowly 1, 2, 3).
6. Remove and massage the injection area for 10 seconds.
7. Call 911 and get emergency medical help right away.




HOW TO USE IMPAX EPINEPHRINE INJECTION (AUTHORIZED GENERIC OF ADRENACLICK®), USP AUTO-INJECTOR, IMPAX LABORATORIES

1. Remove epinephrine auto-injector from its protective carrying case.
2. Pull off both blue end caps: you will now see a red tip.
3. Grasp the auto-injector in your fist with the red tip pointing downward.
4. Put the red tip against the middle of the outer thigh at a 90-degree angle, perpendicular to the thigh.
5. Press down hard and hold firmly against the thigh for approximately 10 seconds.
6. Remove and massage the area for 10 seconds.
7. Call 911 and get emergency medical help right away.



HOW TO USE TEVA'S GENERIC EPIPEN® (EPINEPHRINE INJECTION, USP) AUTO-INJECTOR, TEVA PHARMACEUTICAL INDUSTRIES

1. Quickly twist the yellow or green cap off of the auto-injector in the direction of the "twist arrow" to remove it.
2. Grasp the auto-injector in your fist with the orange tip (needle end) pointing downward.
3. With your other hand, pull off the blue safety release.
4. Place the orange tip against the middle of the outer thigh (upper leg) at a right angle (perpendicular) to the thigh.
5. Swing and push the auto-injector firmly into the middle of the outer thigh until it 'clicks'.
6. Hold firmly in place for 3 seconds (count slowly 1, 2, 3).
7. Remove and massage the injection area for 10 seconds.
8. Call 911 and get emergency medical help right away.



ADMINISTRATION AND SAFETY INFORMATION FOR ALL AUTO-INJECTORS:

1. Do not put your thumb, fingers or hand over the tip of the auto-injector or inject into any body part other than mid-outer thigh. In case of accidental injection, go immediately to the nearest emergency room.
2. If administering to a young child, hold their leg firmly in place before and during injection to prevent injuries.
3. Epinephrine can be injected through clothing if needed.
4. Call 911 immediately after injection.

OTHER DIRECTIONS/INFORMATION (may self-carry epinephrine, may self-administer epinephrine, etc.):

Treat the person before calling emergency contacts. The first signs of a reaction can be mild, but symptoms can worsen quickly.

EMERGENCY CONTACTS — CALL 911

RESCUE SQUAD: _____

DOCTOR: _____ PHONE: _____

PARENT/GUARDIAN: _____ PHONE: _____

OTHER EMERGENCY CONTACTS

NAME/RELATIONSHIP: _____ PHONE: _____

NAME/RELATIONSHIP: _____ PHONE: _____

NAME/RELATIONSHIP: _____ PHONE: _____



KINGSWAY REGIONAL HIGH SCHOOL

Committed to Excellence

*Mrs. Christina Santiago, RN, ATC
School Nurse, 9th & 10th, ext. 4287*

*Mrs. Kate Richards, RN, BSN, CPN
School Nurse 7th, ext. 3022*

*Mrs. Barbara Neal, RN, BSN
School Nurse, 11th & 12th, ext. 4287
High School fax (856) 467-4136*

*Mrs. Megan Anastasia, RN, BSN
School Nurse, 8th, ext. 3022
Middle School fax (856) 467-5968*

Indemnification/Hold Harmless Agreement for Pupil Self-Administration of Medication

We, the parents/guardians of _____
(Pupil Name)

hereby acknowledge and agree that the Kingsway Regional High School District shall incur no liability as a result of any injury arising from the self-administration of medication by

(Pupil Name)

We further agree that, pursuant of N.J.S.A. 18A:40-12.3(d), we shall indemnify, hold harmless and defend the Kingsway Regional High School District, its employees and agents, from and against any and all costs, expenses (including reasonable counsel fees), liabilities, judgements, losses, damages, suits, actions, fines, penalties, claims or demands of any kind and asserted by or on behalf of any person or entity arising out of or in any way connected with the self-administration of medication by _____
(Pupil Name)

Parent/Guardian Signature

Date

Home of the Dragons