



CAMP HESS KRAMER GINDLING HILLTOP CAMP WILSHIRE BOULEVARD TEMPLE

Before campers arrive on site, you must update their immunizations and health history record in Campminder. This needs to be updated every year.

COVID-19: While vaccinations are strongly encouraged, Wilshire Boulevard Temple Camps will not require proof of COVID-19 vaccinations to attend camp.

Immunization Schedule

For a complete overview and further details, please visit the

[Center for Disease Control and Prevention's website:](https://www.cdc.gov/vaccines/schedules/hcp/imz/child-adolescent.html)

<https://www.cdc.gov/vaccines/schedules/hcp/imz/child-adolescent.html>

Vaccine	Prior to Age 4	Ages 4-6	Ages 7-12	Ages 13-17
Diphtheria, Tetanus, and Pertussis (DTaP)	Dose #1 at 2 months of age Dose #2 at 4 months of age Dose #3 at 6 months of age Dose #4 at 15 to 18 months of age	Dose #5 at 4-6 years of age	Age 11 or 12 is recommended age for single dose	Catch-up vaccination if not vaccinated at age 11-12
Measles, Mumps, Rubella	Dose #1 at age 12 months of age	Dose #2 at 4-6 years of age	None unless they have not had two prior doses	None unless they have not had two prior doses
Varicella (Chicken Pox)	Dose #1 at age 12 months of age	Dose #2 at 4-6 years of age	None unless they have	None unless they have not

			not had two	had two prior
MMRV (alternative to separate MMR/Varicel la dosing)	Dose #1 at age 12-15 months of age	Dose #2 at 4-6 years of age	None unless they have not had two prior doses	None unless they have not had two prior doses
Hepatitis B	Dose #1 at birth Dose #2 1-2 months Dose #3 6-18 months (Alternative schedules exist)	Additional doses not needed if immunize d with 3-4 doses appropriat ely spaced as an infant	Additional doses not needed if immunized with 3-4 doses as an infant. If not immunized as an infant, alternative dosing schedule age 11-15 two doses Recombivax 4-6 months apart	Additional doses not needed if immunized with 3-4 doses as an infant or had two dose adolescent series at ages 11-15.
Polio	Dose #1 2 months Dose #2 4 months Dose #3 6-18 months	Dose #4 at age 4 Note: If dose #3 given after 4th birthday and at least 6 months from dose #2 may omit dose#4	Needed only if not immunized previously or foreign travel to designated destinations	Needed only if not immunized previously or foreign travel to designated destinations

Meningitis Vaccine age 11-12 (MenACWY) formerly MCV4

Booster MenACWY at age 16 -18