Diabetes Medical Management Plan (DMMP)

This plan should be completed by the student's personal diabetes health care team, including the parents/guardians. It should be reviewed with relevant school staff and copies should be kept in a place that can be accessed easily by the school nurse, trained diabetes personnel, and other authorized personnel.

Date of plan:	This plan is valid for the current school year:			
Student information				
Student's name:		Date of birth:		
Date of diabetes diagnosis:	□ Type 1	☐ Type 2 ☐ Other:		
School:		School phone number:		
Grade:	Homeroom teacher:			
School nurse:		Phone:		
Contact information				
Parent/guardian 1:				
Address:				
Telephone: Home:				
Email address:				
Address:				
Telephone: Home:				
Email address:				
Address:				
Telephone:		cy number:		
Email address:				
Other emergency contacts:				
Name:	Relat	ionship:		
Telenhone: Home:		Cell:		



Checking blood	glucose					
Brand/model of blood	glucose meter:					
Target range of blood	glucose:					
Before meals: □ 90-	-130 mg/dL □ Othei	r:				
Check blood glucose le	evel:					
☐ Before breakfast	☐ After breakfast		Hours after breakfast	□ 2 h	ours after a cori	rection dose
☐ Before lunch	☐ After lunch		Hours after lunch	☐ Bet	fore dismissal	
☐ Mid-morning						
☐ As needed for signs/symptoms of low or high blood glucose ☐ As needed for signs/symptoms of illnes					s/symptoms of illness	
Preferred site of testing	ng: ☐ Side of fingerti	p 🗆 Oth	ner:	_		
Note: The side of the fi	ingertip should always	be used t	o check blood glucose lev	el if hy	poglycemia is su	ispected.
Student's self-care blo	ood glucose checking s	kills:				
☐ Independently chec	ks own blood glucose					
☐ May check blood gl	ucose with supervisior	1				
☐ Requires school nur	se or trained diabetes	personne	el to check blood glucose			
☐ Uses a smartphone	or other monitoring to	echnology	to track blood glucose va	alues		
Continuous glucose m	onitor (CGM): 🗆 Yes	s □ No	Brand/model:			
Alarms set for: Sev	ere Low:	Lov	v: Hi	igh:		
Predictive alarm: Low	v: Hi	gh:	Rate of change	: Low:		High:
Threshold suspend set	ting:					
CGM may be used for	insulin calculation if gl	ucose is b	etween mg/dL _	Yes	No	
CGM may be used for	hypoglycemia manage	ment	Yes No			
CGM may be used for	hyperglycemia manag	ement	_ Yes No			
 Do not disconnect If the adhesive is p If the CGM becom Refer to the manual 	should be given at leas t from the CGM for spo peeling, reinforce it wi nes dislodged, return e ufacturer's instructions	t three incorts activit th approviverything on how t	ches away from the CGM ties.	s. Do no	ot throw any pai	·
The student troubles	hoots alarms and malf	unctions.			☐ Yes	□No
	hat to do and is able t		h a HIGH alarm.		☐ Yes	□No
	hat to do and is able t	o deal wit	h a LOW alarm.		☐ Yes	□ No
The student knows w		M indica	tes a ranid trending rise o	or	☐ Yes	□ No
	The student knows what to do when the CGM indicates a rapid trending rise or fall in the blood glucose level.					
The student should be	escorted to the nurse	if the CGI	M alarm goes off: ☐ Yes	s 🗆 No	0	
Other instructions for						



Hypoglycemia treatment				
Student's usual symptoms of hypoglycemia (list below):				
If exhibiting symptoms of hypoglyc product equal to grams of o		vel is less tha	nmg/dL,	give a quick-acting glucose
Recheck blood glucose in 15 minute	es and repeat treatment if b	olood glucose	level is less than	ı mg/dL.
Additional treatment:				
If the student is unable to eat or d movement): • Position the student on his or leading to the student o			is having seizure	e activity or convulsions (jerking
Administer glucagon	Name of glucagon use	d:		
Injection:				
□ 1 mg	□½ mg □ Ot	her (dose)		
• Route:	☐ Subcutaneous (SC)			
 Site for glucagon injection: 	☐ Buttocks	☐ Arm	☐ Thigh	☐ Other:
Nasal route:				
☐ 3 mg				
Route:Site:	☐ Intranasal (IN)☐ Nose			
 Call 911 (Emergency Medical S Contact the student's health ca If on insulin pump, stop by place 	are provider.	_		vith EMS to hospital.
Hyperglycemia treatment				
Student's usual symptoms of hype	rgiycemia (iist below):			
Check ☐ Urine ☐ Blood for For blood glucose greater than insulin (see correction dose or Notify parents/guardians if blooms for insulin pump users: see Additional Allow unrestricted access to the	ders). ood glucose is over Iditional Information for St ne bathroom.	st hour mg/dL. udent with In	s since last insul	in dose, give correction dose of
Additional treatment for ketones:				

• Follow physical activity and sports orders. (See Physical Activity and Sports)

If the student has symptoms of a hyperglycemia emergency, call 911 (Emergency Medical Services) and contact the student's parents/guardians and health care provider. Symptoms of a hyperglycemia emergency include: dry mouth, extreme thirst, nausea and vomiting, severe abdominal pain, heavy breathing or shortness of breath, chest pain, increasing sleepiness or lethargy, or depressed level of consciousness.



Insulin therapy			
Insulin delivery device:	☐ Syringe	☐ Insulin pen	☐ Insulin pump
Type of insulin therapy at school:	☐ Adjustable (basal-bolus) ins	sulin	oy 🗆 No insulin
Adjustable (Basal-bolus) Insulin T • Carbohydrate Coverage/Corr	• •	·	
• Carbohydrate Coverage:			
Insulin-to-carbohydrate rati	o:		
Breakfast: 1 unit of insulin p Lunch: 1 unit of insulin per_ Snack: 1 unit of insulin per_		•	
	Carbohydrate Dose Ca	culation Example	
Total Grams of Carb	ohydrate to Be Eaten ÷ Insulin-	to-Carbohydrate Ratio =	Units of Insulin
Correction Dose: Blood glucose co	orrection factor (insulin sensitiv	ity factor) = Target bl	ood glucose =mg/dL
	Correction Dose Calcu	ulation Example	
(Current Blood Gluco	ose — Target Blood Glucose) ÷ C	Correction Factor = Unit	ts of Insulin
Correction dose scale (use instead	d of calculation above to determ	nine insulin correction dose):	
Blood glucose to m	g/dL, give units E	Blood glucose to	_ mg/dL, give units
Blood glucose to m	g/dL, give units E	Blood glucose to	_ mg/dL, give units
See the worksheet examples in Adv instructions on how to compute the	-	=	
When to give insulin:			
Breakfast			
☐ Carbohydrate coverage only			
☐ Carbohydrate coverage plus co last insulin dose.	rrection dose when blood glucc	se is greater than mg/c	IL and hours since
☐ Other:			
Lunch			
☐ Carbohydrate coverage only			
☐ Carbohydrate coverage plus co last insulin dose.	rrection dose when blood gluco	se is greater than m	g/dL and hours since
☐ Other:			
Snack			
☐ No coverage for snack			
☐ Carbohydrate coverage only			
☐ Carbohydrate coverage plus co last insulin dose.	rrection dose when blood gluco	se is greater than m _{	g/dL and hours since
\square Correction dose only: For blood	d glucose greater than	mg/dL AND at leasth	ours since last insulin dose.
☐ Other:			



Insulin therap	y (continued)				
Fixed Insulin Therapy Name of insulin:					
☐ Units of insulin given pre-breakfast daily					
☐ Units of insulin given pre-lunch daily ☐ Units of insulin given pre-snack daily					
					☐ Other:
Basal Insulin Thera	apy Name of insulin:				
To be given during	g school hours: Pre	e-breakfast dose:	units		
	Pre	e-lunch dose:	units		
	Pre	e-dinner dose:	units		
Other diabetes me	edications:				
Name:	Dose:	Route:	Times gi	ven:	
				ven:	
					
Parents/Guardian	s Authorization to Adjus	t Insulin Dose			
☐ Yes ☐ No F	Parents/guardians autho	rization should be o	btained before adminis	tering a correction dose.	
	Parents/guardians are au range: +/ units o		e or decrease correction	dose scale within the following	
	_			carbohydrate ratio within the +/ grams of carbohydrate.	
	Parents/guardians are aut		or decrease fixed insulin	dose within the following range:	
Student's self-care	insulin administration	skills:			
☐ Independently of	calculates and gives own	injections.			
☐ May calculate/g	give own injections with	supervision.			
☐ Requires school supervision.	nurse or trained diabete	es personnel to calc	ulate dose and student	can give own injection with	
☐ Requires school	nurse or trained diabet	es personnel to calc	ulate dose and give the	injection.	
·		•	_		
Additional inf	formation for stud	lent with insul	in pump		
Brand/model of p	ump:		Type of insulin in pump	:	
				Basal rate:	
				Basal rate:	
		Basal rate:			
Other pump instru	uctions:				
osne. pamp matre					
Type of infusion se					
. The or illingion se	~ · · <u></u>				



Additional information for stude	nt with insulin pum	p (continued)		
Appropriate infusion site(s):				
☐ For blood glucose greater than mg failure or infusion site failure. Notify paren	within hours after	hours after correction, consider pump		
☐ For infusion site failure: Insert new infusion	n set and/or replace reserv	oir, or give insulin by syrin	ge or pen.	
☐ For suspected pump failure: Suspend or re	•		-	
Physical Activity				
May disconnect from pump for sports activiti	es: 🗆 Yes, for	hours	□No	
Set a temporary basal rate:		temporary basal for	hours \square No	
Suspend pump use:	☐ Yes, for h		□ No	
Student's Self-care Pump Skills: Check "Y Counts carbohydrates Calculates correct amount of insulin for carb		nt can perform the skill	independently.	
Administers correction bolus	•	☐ Yes	□No	
Calculates and sets basal profiles			□No	
Calculates and sets temporary basal rate			□ No	
Changes batteries		☐ Yes	□ No	
Disconnects pump		□ Yes	□ No	
Reconnects pump to infusion set		□ Yes	□ No	
Prepares reservoir, pod, and/or tubing		□ Yes	□ No	
Inserts infusion set Troubleshoots alarms and malfunctions		□ Yes	□ No □ No	
Troubleshoots alarms and manunctions		☐ Yes	□ NO	
Meal plan				
Meal/Snack	Time	Carbohydra	te Content (grams)	
Breakfast			to	
Mid-morning snack			to	
Lunch		to		
Mid-afternoon snackto			to	
Other times to give snacks and content/amo				
Instructions for when food is provided to the	e class (e.g., as part of a cla	ss party or food sampling	event):	
Parent/guardian substitution of food for mea	ls, snacks and special even	ts/parties permitted.		
Special event/party food permitted: \qed Par	ents'/Guardians' discretion	n ☐ Student discretion		
Student's self-care nutrition skills: ☐ Independently counts carbohydrates				
☐ May count carbohydrates with supervision	1			
☐ Requires school nurse/trained diabetes pe		rates		



Physical activity and sports	
A quick-acting source of glucose such as $\ \square$ glucose tabs and/or $\ \square$ sugar-containing physical education	juice must be available at the site of on activities and sports.
Student should eat ☐ 15 grams ☐ 30 grams of carbohydrate ☐ other:	
□ before □ every 30 minutes during □ every 60 minutes during □ after vigorou	s physical activity 🛛 other:
If most recent blood glucose is less thanmg/dL, student can participate in physicorrected and abovemg/dL.	cal activity when blood glucose is
Avoid physical activity when blood glucose is greater thanmg/dL or if urine/b	blood ketones are moderate to large.
(See Administer Insulin for additional information for students on insulin pumps.)	
Disaster/emergency and drill plan	
To prepare for an unplanned disaster, emergency (72 hours) or drill, obtain emergency School nurse or other designated personnel should take student's diabetes supplies an destination to make available to student for the duration of the unplanned disaster, en	d medications to student's
☐ Continue to follow orders contained in this DMMP.	
☐ Additional insulin orders as follows (e.g., dinner and nighttime):	
□ Other:	
Signatures	
This Diabetes Medical Management Plan has been approved by:	
This Diabetes Medical Management Flan has been approved by.	
Student's Physician/Health Care Provider	Date
I, (parent/guardian) give permiss	ion to the school nurse or another
qualified health care professional or trained diabetes personnel of (school)	
and carry out the diabetes care tasks as outlined in (student) Management Plan. I also consent to the release of the information contained in this Diabetes.	Diabetes Medical
school staff members and other adults who have responsibility for my child and who may	=
maintain my child's health and safety. I also give permission to the school nurse or another	
contact my child's physician/health care provider.	
Acknowledged and received by:	
Student's Parent/Guardian	Date
Student's Parent/Guardian	
	Date
	Date

