

Student's Name:

Canon McMillan School District Parental Release of Allergy Information

Your child has been identified with an allergy that could have serious implications to his/her health. For your child's safety, the allergy information contained in his/her Individualized Medical Plan (IMP) and/or their Allergy Action Plan will be shared with all employees having contact with him/her. This will include classroom teachers, related arts teachers, nurses, paraeducators, custodians, bus drivers, substitute employees and any other Canon-McMillan School District employee that will have contact with your child. This information will be shared on a "need to know" basis and will be kept confidential by these employees.

Our goal is to provide a safe and healthy environment for every child. Sharing this information with appropriate personnel will provide important communication between home and school to help achieve this goal.

Please complete this form and return it to your child's school nurse or principal.

Ctadent's Name.	
Allergies:	
Yes, I give permission for my child's allerg school personnel as outlined above.	gy information to be shared with the appropriate
Parent's Signature:	Date:

Grade.

	Canon-McMinan School District - F	Hiergy Av	MORE FAI	
	D.O.B.: Te	acher:	· ·	Place Child's Picture
ALLERGY	TO:		1610-1610-1610-1610-161 0	Here
Asthmatic	Yes* ☐ No ☐ *Higher risk for severe reaction			
	♦ STEP 1: TREATME	NT •		
Symptoms		;	Give Checked M **(To be determine authorizing treatment	d by physician
If a food	allergen has been ingested, but no symptoms:		□ Epinephrine	□ Antihistamine
Mouth	Itching, tingling, or swelling of lips, tongue, mouth		□ Epinephrine	□ Antihistamine
Skin	Hives, itchy rash, swelling of the face or extremities		I Epinephrine	□ Antihistamine
Gut	Nausea, abdominal cramps, vomiting, diarrhea		□ Epinephrine	□ Antihistamine
Throat †	Tightening of throat, hoarseness, hacking cough	Marking and printers and printers are a second	□ Epinephrine	디 Antihistamine
Lung†	Shortness of breath, repetitive coughing, wheezing		□ Epinephrine	□ Antihistamine
Heart †	Weak or thready pulse, low blood pressure, fainting, pale, blu-	eness	□ Epinephrine	□ Antihistamine
Other †			□ Epinephrine	[.] □ Antihistamine
■ Ifreaction	on is progressing (several of the above areas affected), give:		□ Epinephrine	□ Antihistamine
Antihistamir	ne: give	mantant .	n	add Pillian in the and a state of the state
Other: give _	medication/dose/route	yar a a a a a a a a a a a a a a a a a a		**************************************
IMPORTAN	VT: Asthma inhalers and/or antihistamines cannot be depend			in anaphylaxis.
1. Call 911 (c	or Rescue Squad:). State that an allergic reaction has b	een treated,	and additional epi	nephrine may be needed.
2. Dr	Phone Number	**	managaman salaman managamban ya masha Nobi Mashili Marita (1808)	III/ II A TO THE TOTAL THE TOTAL TO THE TOTAL THE TOTAL TO THE TOTAL THE TOTAL TO T
3. Parent	Phone Number	r(s):		***************************************
4. Emergency Nan	contacts: ne/Relationship Phone Number(s)			
a	1)	······································	2)	di MANATON NA BARILLIA, ara en al districti de se di Menatonia ancienta de se anticolo de la compansión de se a
	1)			
***************************************	ENT/GUARDIAN CANNOT BE REACHED, DO NOT HESITATE T			
Parent/Guard	ian's Signature	-	Date	And the state of t
Doctor's Sign	nature		Date	

(Required)

TRAINED STAFF MEMBERS				
	Room			
2	Room			
3,	Room			

EpiPen® and EpiPen® Jr. Directions

Pull off gray activation cap.



 Hold black tip near outer thigh (always apply to thigh).



Swing and jab firmly into outer thigh until Auto-Injector mechanism functions. Hold in place and count to 10. Remove the EpiPen® unit and massage the injection area for 10 seconds. Twinject® 0.3 mg and Twinject® 0.15 mg Directions



- Remove caps labeled "1" and "2."
- Place rounded tip against outer thigh, press down hard until needle penetrates. Hold for 10 seconds, then remove.



SECOND DOSE ADMINISTRATION:
If symptoms don't improve after
10 minutes, administer second dose:

 Unscrew rounded tip. Pull syringe from barrel by holding blue collar at needle base.



- Slide yellow collar off plunger.
- Put needle into thigh through skin, push plunger down all the way, and remove.



Once EpiPen® or Twinject® is used, call the Rescue Squad. Take the used unit with you to the Emergency Room. Plan to stay for observation at the Emergency Room for at least 4 hours.

For children with multiple food allergies, consider providing separate Action Plans for different foods.

**Medication checklist adapted from the Authorization of Emergency Treatment form developed by the Mount Sinai School of Medicine, Used with permission.





Authorization for Medication During School Hours

(prescription and over-the-counter)

Student's Name:	
School:	Grade:
The student named above must receive the fol- maintain sufficient health to participate in the s manufacturer's container or the pharmacy labe	, —
Name of Medication:	
Prescribed Dosage:	
Time Schedule:	
Length of Time (days/weeks):	
Reason for Administration:	
Possible Side Effects:	
Asthma Inhalers	
My child (check only one) is able to self-a	administer the medication. self-administer the medication.
If the student can self-administer, he/she has p	permission to carry the inhaler.
Epi-Pen	
My child (check only one) is permitted to in its not permitted	carry the epi-pen with them. I to carry the epi-pen with them.
I do hereby release, discharge and hold harmle agents and employees, from any and all liabilit administration of the above medication to my o	y and claims whatsoever arising from the
Physician's Signature:	Date:
Parent/Guardian's Signature:	Date: