

# Summary of Immunization Requirements for School Attendance

(Refer to [Kentucky 902 KAR 2:060](#) for details on exceptions and exemptions)

By Age:	A child is required to have the following immunizations/doses for attendance:	
5 & 6 years	DTaP or DTP	(DT may be used if contradiction to Pertussis) ✓ 5 doses, or ✓ 4 doses if 4th dose given at age 4 or older and at least 6 months after previous dose
	IPV or OPV	✓ 4 doses with 4th dose given at age 4 or after and at least 6 months after previous dose (if 4 or more doses was given before age 4, will still need an additional dose at age 4 or after and at least 6 months from previous dose), or ✓ 3 doses if 3rd dose was given at age 4 or after and at least 6 months from previous dose
	Hepatitis B	✓ 3 doses (last dose must be given after the 6-month birthday)
	Hepatitis A	✓ 2 doses (spaced at least 6 months apart)
	MMR	✓ 2 doses
	Varicella	✓ 2 doses (unless varicella immunity due to history of chicken pox verified by healthcare provider)
11 & 12 years	DTaP or DTP	(DT may be used if contradiction to Pertussis) ✓ 5 doses, or ✓ 4 doses if 4th dose given at age 4 or older and at least 6 months after previous dose, or ✓ 2 doses (any combination of DTaP, DTP, DT or TD) with 1 dose Tdap or Td received after ✓ 2 doses of Td after the dose of Tdap may take the place of DTaP vaccines
	IPV or OPV	✓ 4 doses with 4th dose given at age 4 or after and at least 6 months after previous dose (if 4 or more doses was given before age 4, will still need an additional dose at age 4 or after and at least 6 months from previous dose), or ✓ 3 doses if 3rd dose was given at age 4 or after and at least 6 months from previous dose
	Hepatitis B	✓ 3 doses (last dose must be given after the 6-month birthday), or ✓ 2 doses of Adult Hep B given after age 11
	Hepatitis A	✓ 2 doses (spaced at least 6 months apart)
	MMR	✓ 2 doses
	Varicella	✓ 2 doses (unless varicella immunity due to history of chicken pox verified by healthcare provider)
	Tdap	✓ 1 dose (Td may be used if contraindication to Pertussis vaccine)
MenACWY	✓ 1 dose	
16 years & Older	DTaP or DTP	(DT may be used if contradiction to Pertussis) ✓ 5 doses, or ✓ 4 doses if 4th dose given at age 4 or older and at least 6 months after previous dose, or ✓ 2 doses (any combination of DTaP, DTP, DT or TD) with 1 dose Tdap or Td received after, or ✓ 2 doses of Td after the dose of Tdap may take the place of DTaP vaccines
	IPV or OPV	✓ 4 doses with 4th dose given at age 4 or after and at least 6 months after previous dose (if 4 or more doses was given before age 4, will still need an additional dose at age 4 or after and at least 6 months from previous dose), or ✓ 3 doses if 3rd dose was given at age 4 or after and at least 6 months from previous dose
	Hepatitis B	✓ 3 doses (last dose must be given after the 6-month birthday), or ✓ 2 doses of Adult Hep B given after age 11
	Hepatitis A	✓ 2 doses (spaced at least 6 months apart)
	MMR	✓ 2 doses
	Varicella	✓ 2 doses (unless varicella immunity due to history of chicken pox verified by healthcare provider)
	Tdap	✓ 1 dose (Td may be used if contraindication to Pertussis vaccine)
MenACWY	✓ 2 doses, or ✓ 1 dose if received at age 16 or older	



\*NOTE: Refer to [Kentucky 902 KAR 2:060](#) for ages 7-10 and 13-15 for required regulations.