



Seizure Action Plan

Effective Date _____

This student is being treated for a seizure disorder. The information below should assist you if a seizure occurs during school hours.

Student's Name _____ Date of Birth _____

Parent/Guardian _____ Phone _____ Cell _____

Other Emergency Contact _____ Phone _____ Cell _____

Treating Physician _____ Phone _____

Significant Medical History _____

Seizure Information

Seizure Type	Length	Frequency	Description

Seizure triggers or warning signs: _____

Student's response after a seizure: _____

Basic First Aid: Care & Comfort

Please describe basic first aid procedures: _____

Does student need to leave the classroom after a seizure? Yes No

If YES, describe process for returning student to classroom: _____

Basic Seizure First Aid

- Stay calm & track time
- Keep child safe
- Do not restrain
- Do not put anything in mouth
- Stay with child until fully conscious
- Record seizure in log

For tonic-clonic seizure:

- Protect head
- Keep airway open/watch breathing
- Turn child on side

A seizure is generally considered an emergency when:

- Convulsive (tonic-clonic) seizure lasts longer than 5 minutes
- Student has repeated seizures without regaining consciousness
- Student is injured or has diabetes
- Student has a first-time seizure
- Student has breathing difficulties
- Student has a seizure in water

Emergency Response

A "seizure emergency" for this student is defined as: _____

Seizure Emergency Protocol

(Check all that apply and clarify below)

- Contact school nurse at _____
- Call 911 for transport to _____
- Notify parent or emergency contact
- Administer emergency medications as indicated below
- Notify doctor
- Other _____

Treatment Protocol During School Hours (include daily and emergency medications)

Emerg. Med. ✓	Medication	Dosage & Time of Day Given	Common Side Effects & Special Instructions

Does student have a Vagus Nerve Stimulator? Yes No If YES, describe magnet use: _____

Special Considerations and Precautions (regarding school activities, sports, trips, etc.)

Describe any special considerations or precautions: _____

Physician Signature _____ Date _____

Parent/Guardian Signature _____ Date _____

Food Allergy Action Plan

Name: _____ D.O.B.: / /

Allergy to: _____

Weight: _____ lbs. Asthma: Yes (higher risk for a severe reaction) No

Place
Student's
Picture
Here

Extremely reactive to the following foods: _____

THEREFORE:

- If checked, give epinephrine immediately for ANY symptoms if the allergen was *likely* eaten.
 If checked, give epinephrine immediately if the allergen was *definitely* eaten, even if no symptoms are noted.

Any **SEVERE SYMPTOMS** after suspected or known ingestion:

One or more of the following:

- LUNG: Short of breath, wheeze, repetitive cough
 HEART: Pale, blue, faint, weak pulse, dizzy, confused
 THROAT: Tight, hoarse, trouble breathing/swallowing
 MOUTH: Obstructive swelling (tongue and/or lips)
 SKIN: Many hives over body

Or combination of symptoms from different body areas:

- SKIN: Hives, itchy rashes, swelling (e.g., eyes, lips)
 GUT: Vomiting, crampy pain



1. **INJECT EPINEPHRINE IMMEDIATELY**
2. Call 911
3. Begin monitoring (see box below)
4. Give additional medications:*
 -Antihistamine
 -Inhaler (bronchodilator) if asthma

*Antihistamines & inhalers/bronchodilators are not to be depended upon to treat a severe reaction (anaphylaxis). **USE EPINEPHRINE.**

MILD SYMPTOMS ONLY:

- MOUTH: Itchy mouth
 SKIN: A few hives around mouth/face, mild itch
 GUT: Mild nausea/discomfort



1. **GIVE ANTIHISTAMINE**
2. Stay with student; alert healthcare professionals and parent
3. If symptoms progress (see above), **USE EPINEPHRINE**
4. Begin monitoring (see box below)

Medications/Doses

Epinephrine (brand and dose): _____

Antihistamine (brand and dose): _____

Other (e.g., inhaler-bronchodilator if asthmatic): _____

Monitoring

Stay with student; alert healthcare professionals and parent. Tell rescue squad epinephrine was given; request an ambulance with epinephrine. Note time when epinephrine was administered. A second dose of epinephrine can be given 5 minutes or more after the first if symptoms persist or recur. For a severe reaction, consider keeping student lying on back with legs raised. Treat student even if parents cannot be reached. See back/attached for auto-injection technique.

Parent/Guardian Signature _____

Date _____

Physician/Healthcare Provider Signature _____

Date _____

TURN FORM OVER

Form provided courtesy of FAAN (www.foodallergy.org) 7/2010

Date of Plan: _____

Diabetes Medical Management Plan

Effective Dates: _____

This plan should be completed by the student's personal health care team and parents/guardian. It should be reviewed with relevant school staff and copies should be kept in a place that is easily accessed by the school nurse, trained diabetes personnel, and other authorized personnel.

Student's Name: _____

Date of Birth: _____ Date of Diabetes Diagnosis: _____

Grade: _____ Homeroom Teacher: _____

Physical Condition: Diabetes type 1 Diabetes type 2

Contact Information

Mother/Guardian: _____

Address: _____

Telephone: Home _____ Work _____ Cell _____

Father/Guardian: _____

Address: _____

Telephone: Home _____ Work _____ Cell _____

Student's Doctor/Health Care Provider:

Name: _____

Address: _____

Telephone: _____ Emergency Number: _____

Other Emergency Contacts:

Name: _____

Relationship: _____

Telephone: Home _____ Work _____ Cell _____

Notify parents/guardian or emergency contact in the following situations:

Diabetes Medical Management Plan *Continued*

Blood Glucose Monitoring

Target range for blood glucose is 70-150 70-180 Other _____

Usual times to check blood glucose _____

Times to do extra blood glucose checks (*check all that apply*)

before exercise

after exercise

when student exhibits symptoms of hyperglycemia

when student exhibits symptoms of hypoglycemia

other (explain): _____

Can student perform own blood glucose checks? Yes No

Exceptions: _____

Type of blood glucose meter student uses: _____

Insulin

Usual Lunchtime Dose

Base dose of Humalog/Novolog /Regular insulin at lunch (circle type of rapid-/short-acting insulin used) is _____ units or does flexible dosing using _____ units/ _____ grams carbohydrate.

Use of other insulin at lunch: (circle type of insulin used): intermediate/NPH/lente _____ units or basal/Lantus/Ultralente _____ units.

Insulin Correction Doses

Parental authorization should be obtained before administering a correction dose for high blood glucose levels. Yes No

_____ units if blood glucose is _____ to _____ mg/dl

_____ units if blood glucose is _____ to _____ mg/dl

_____ units if blood glucose is _____ to _____ mg/dl

_____ units if blood glucose is _____ to _____ mg/dl

_____ units if blood glucose is _____ to _____ mg/dl

Can student give own injections? Yes No

Can student determine correct amount of insulin? Yes No

Can student draw correct dose of insulin? Yes No

_____ Parents are authorized to adjust the insulin dosage under the following circumstances: _____

For Students With Insulin Pumps

Type of pump: _____ Basal rates: _____ 12 am to _____

_____ to _____

_____ to _____

Type of insulin in pump: _____

Type of infusion set: _____

Insulin/carbohydrate ratio: _____ Correction factor: _____

Diabetes Medical Management Plan *Continued*

Student Pump Abilities/Skills:

Needs Assistance

- Count carbohydrates Yes No
- Bolus correct amount for carbohydrates consumed Yes No
- Calculate and administer corrective bolus Yes No
- Calculate and set basal profiles Yes No
- Calculate and set temporary basal rate Yes No
- Disconnect pump Yes No
- Reconnect pump at infusion set Yes No
- Prepare reservoir and tubing Yes No
- Insert infusion set Yes No
- Troubleshoot alarms and malfunctions Yes No

For Students Taking Oral Diabetes Medications

Type of medication: _____ Timing: _____

Other medications: _____ Timing: _____

Meals and Snacks Eaten at School

Is student independent in carbohydrate calculations and management? Yes No

<i>Meal/Snack</i>	<i>Time</i>	<i>Food content/amount</i>
Breakfast	_____	_____
Mid-morning snack	_____	_____
Lunch	_____	_____
Mid-afternoon snack	_____	_____
Dinner	_____	_____

Snack before exercise? Yes No

Snack after exercise? Yes No

Other times to give snacks and content/amount: _____

Preferred snack foods: _____

Foods to avoid, if any: _____

Instructions for when food is provided to the class (e.g., as part of a class party or food sampling event):

Exercise and Sports

A fast-acting carbohydrate such as _____ should be available at the site of exercise or sports.

Restrictions on activity, if any: _____

Student should not exercise if blood glucose level is below _____ mg/dl or above _____ mg/dl or if moderate to large urine ketones are present.

TOOLS