

DEPARTMENT OF SCHOOL HEALTH SERVICES

Danville Independent Schools

TO: Parent/guardian of: _____
Date of birth: _____
Enrolled in grade _____
Attends _____ school

You have checked on school records that this student has a **severe or serious allergy**. It is important to have at least annual health information when s/he needs help at school. **Please complete this form and return it to the school nurse tomorrow so a plan to help your child can be shared with identified school personnel.** It is the **responsibility of parents** to provide necessary special food and medicine needed at school. If you have any questions, you may call the nurse at the student's school or the school health office at the number above.

CHECK ANY LIFE THREATENING ALLERGY YOUR STUDENT HAS:

- A- ☐ insect stings (list type) _____
B- ☐ food (list type) _____
C- ☐ animals (list type) _____
D- ☐ other (list) _____

INDICATE THE SIGNS THAT ARE USUALLY PRESENT DURING ALLERGY ATTACK by placing letter(s) of allergies checked above by the symptoms listed below.

_____ difficulty breathing	_____ rash
_____ difficulty swallowing	_____ nausea
_____ loss of consciousness	_____ flushed or unusually pale skin
_____ swelling:	_____ other (list) _____
how much? _____	_____
where? _____	_____

Has emergency medical treatment been needed in the past year for allergies? ☐ No ☐ Yes
(when): _____

Allergies are currently being treated by Dr. : _____

ARE MEDICATIONS NEEDED TO CONTROL THE ALLERGY (IES)? ☐ No ☐ Yes (list below)

Medications	Amount taken	Time of Day

(Turn page over)

Do any of these medications need to be taken at school? ☐ no ☐ yes

Which one(s)? _____ * (see below) _____

PLEASE ADVISE THE SCHOOL NURSE AT ONCE OF ANY CHANGES IN DOSE AND/OR TYPE OF MEDICATION

THE USUAL TREATMENT AT SCHOOL FOR A STUDENT HAVING A SEVERE ALLERGIC REACTION IS TO:

1. Assist student with the prescribed medication.
2. Observe the student for inadequate breathing; signs of shock; unusual swelling. If any of these signs are present, 9-1-1 will be called.
3. Report to parent.

If you want additional help given, or have other concerns, describe here; ** (see below)

X

Parent Signature

Date

Remember to advise the school immediately of changes in phone numbers, address, responsible emergency contact persons, doctor, hospital preferences.

- * Your district has a policy regarding taking medicines at school. Please check with your child's school for direction.
- ** Tests and activity restrictions require written directions from the student's doctor.

Name: _____ D.O.B.: _____

Allergy to: _____

Notes: _____ Ibs, Asthma: ☐ Yes (higher risk for a severe reaction) ☐ No

NOTE: Do not depend on antihistamines or inhalers (bronchodilators) to treat a severe reaction. **USE EPINEPHRINE.**

Extremely reactive to the following allergens: _____

THEREFORE:

- ☐ If checked, give epinephrine immediately if the allergen was **LIKELY** eaten, for **ANY** symptoms.
- ☐ If checked, give epinephrine immediately if the allergen was **DEFINITELY** eaten, even if no symptoms are apparent.

FOR ANY OF THE FOLLOWING:
SEVERE SYMPTOMS



LUNG

Shortness of breath, wheezing, repetitive cough



HEART

Pale or bluish skin, faintness, weak pulse, dizziness



THROAT

Tight or hoarse throat, trouble breathing or swallowing



MOUTH

Significant swelling of the tongue or lips



SKIN

Many hives over body, widespread redness



GUT

Repetitive vomiting, severe diarrhea



OTHER

Feeling something bad is about to happen, anxiety, confusion

OR A
COMBINATION
of symptoms
from different
body areas.

1. **INJECT EPINEPHRINE IMMEDIATELY.**
2. **Call 911.** Tell emergency dispatcher the person is having anaphylaxis and may need epinephrine when emergency responders arrive.
 - Consider giving additional medications following epinephrine:
 - » Antihistamine
 - » Inhaler (bronchodilator) if wheezing
 - Lay the person flat, raise legs and keep warm. If breathing is difficult or they are vomiting, let them sit up or lie on their side.
 - If symptoms do not improve, or symptoms return, more doses of epinephrine can be given about 5 minutes or more after the last dose.
 - Alert emergency contacts.
 - Transport patient to ER, even if symptoms resolve. Patient should remain in ER for at least 4 hours because symptoms may return.

MILD SYMPTOMS



NOSE

Itchy or runny nose, sneezing



MOUTH

Itchy mouth



SKIN

A few hives, mild itch



GUT

Mild nausea or discomfort

FOR MILD SYMPTOMS FROM MORE THAN ONE SYSTEM AREA, GIVE EPINEPHRINE.

FOR MILD SYMPTOMS FROM A SINGLE SYSTEM AREA, FOLLOW THE DIRECTIONS BELOW:

1. Antihistamines may be given, if ordered by a healthcare provider.
2. Stay with the person; alert emergency contacts.
3. Watch closely for changes. If symptoms worsen, give epinephrine.

MEDICATIONS/DOSES

Epinephrine Brand or Generic: _____

Epinephrine Dose: ☐ 0.1 mg IM ☐ 0.15 mg IM ☐ 0.3 mg IM

Antihistamine Brand or Generic: _____

Antihistamine Dose: _____

Other (e.g., inhaler-bronchodilator if wheezing): _____

PATIENT OR PARENT/GUARDIAN AUTHORIZATION SIGNATURE

DATE

PHYSICIAN/HCP AUTHORIZATION SIGNATURE

DATE