

Green Brook Family Medicine

Ronald M Frank, MD FAAFP
Sean M Cook, MD
Clair Carragino, APN-C

328 Greenbrook Road
Green Brook, New Jersey 08812

Phone 732.356.0266
Fax 732.356.5022

December 12, 2018

Naloxone Standing Orders

Intranasal Naloxone may be administered by the school nurse or appropriately trained school employees to any individual on school grounds suspected of experiencing an opioid overdose.

1. For individuals who are suspected to be experiencing an opioid overdose:
 - a. Activate EMS/Call 911
 - b. Provide a 4mg Narcan Nasal spray dose as instructed in the attached QUICK START GUIDE.
 - c. Narcan 4mg intranasal spray may be repeated (alternating nostrils) every 2-3 minutes as needed
2. Things to remember:
 - a. The individual may require CPR
 - b. The individual should be placed on their side after receiving Narcan to prevent aspiration if they vomit.
 - c. When the individual responds, they may become agitated and you need to watch out for your safety.
 - d. They must be transported by EMS to a hospital for further treatment.
 - e. Even after an individual respond, they can relapse and need more NARCAN

NARCAN[®] (naloxone HCl) NASAL SPRAY

QUICK START GUIDE Opioid Overdose Response Instructions

Use NARCAN Nasal Spray (naloxone hydrochloride) for known or suspected opioid overdose in adults and children.

Important: For use in the nose only.

Do not remove or test the NARCAN Nasal Spray until ready to use.

1 Identify Opioid Overdose and Check for Response


Ask person if he or she is okay and shout name.

Shake shoulders and firmly rub the middle of their chest.

Check for signs of opioid overdose:

- Will not wake up or respond to your voice or touch
- Breathing is very slow, irregular, or has stopped
- Center part of their eye is very small, sometimes called "pinpoint pupils"

Lay the person on their back to receive a dose of NARCAN Nasal Spray.




2 Give NARCAN Nasal Spray

Remove NARCAN Nasal Spray from the box.

Peel back the tab with the circle to open the NARCAN Nasal Spray.



Hold the NARCAN nasal spray with your thumb on the bottom of the plunger and your first and middle fingers on either side of the nozzle.




Gently insert the tip of the nozzle into either nostril.

- Tilt the person's head back and provide support under the neck with your hand. Gently insert the tip of the nozzle into **one nostril**, until your fingers on either side of the nozzle are against the bottom of the person's nose.



Press the plunger firmly to give the dose of NARCAN Nasal Spray.

- Remove the NARCAN Nasal Spray from the nostril after giving the dose.



Get emergency medical help right away.

3 Call for emergency medical help, Evaluate, and Support


Move the person on their side (recovery position)

after giving NARCAN Nasal Spray.

Watch the person closely.

If the person does not respond by waking up, to voice or touch, or breathing normally another dose may be given. NARCAN Nasal Spray may be dosed every 2 to 3 minutes, if available.

Repeat Step 2 using a new NARCAN Nasal Spray to give another dose in the other nostril. If additional NARCAN Nasal Sprays are available, repeat step 2 every 2 to 3 minutes until the person responds or emergency medical help is received.



For more information about NARCAN Nasal Spray, go to www.narcannasalspray.com, or call 1-844-4NARCAN (1-844-462-7226).

©2015 ADAPT Pharma, Inc. NARCAN[®] is a registered trademark licensed to ADAPT Pharma Operations Limited. A1009.01