

Syllabus

Course Title: 7th Grade Accelerated Mathematics

Grade: 7

DESCRIPTION OF COURSE: Seventh grade Accelerated Math is a course aligned to the PA Core standards. There are five major units of study including the Number System, Expressions and Equations, Functions, Geometry, and Statistics and Probability.

REQUIRED TOPICS OF STUDY	SUGGESTED INSTRUCTIONAL TIME	STANDARDS/ ELIGIBLE CONTENT
The Number System	45 Days	<p>CC.2.1.8.E.1 Distinguish between rational and irrational numbers using their properties.</p> <p>M08.A-N.1.1.1 Determine whether a number is rational or irrational. For rational numbers, show that the decimal expansion terminates or repeats (limit repeating decimals to thousandths).</p> <p>M08.A-N.1.1.2 Convert a terminating or repeating decimal to a rational number (limit repeating decimals to thousandths).</p> <p>CC.2.1.8.E.4 Demonstrate an understanding of rational and irrational numbers.</p> <p>M08.A-N.1.1.3 Estimate the value of irrational numbers without a calculator (limit whole number radicand to less than 144). Example: $\sqrt{5}$ is between 2 and 3 but closer to 2.</p> <p>M08.A-N.1.1.4 Use rational approximations of irrational numbers to compare and order irrational numbers.</p> <p>M08.A-N.1.1.5 Locate/identify rational and irrational numbers at their approximate locations on a number line.</p>

Expressions and Equations	49 Days	<p>CC.2.2.8.B.1 Apply concepts of integer exponents to generate equivalent expressions.</p> <p>M08.B-E.1.1.1 Apply one or more properties of integer exponents to generate equivalent numerical expressions without a calculator (with final answers expressed in exponential form with positive exponents). Properties will be provided. Example: $3^{12} \times 3^{-15} = 3^{-3} = 1/(3^3)$</p> <p>M08.B-E.1.1.2 Use square root and cube root symbols to represent solutions to equations of the form $x^2 = p$ and $x^3 = p$, where p is a positive rational number. Evaluate square roots of perfect squares (up to and including 12^2) and cube roots of perfect cubes (up to and including 5^3) without a calculator. Example: If $x^2 = 25$ then $x = \pm\sqrt{25}$.</p> <p>M08.B-E.1.1.3 Estimate very large or very small quantities by using numbers expressed in the form of a single digit times an integer power of 10 and express how many times larger or smaller one number is than another. Example: Estimate the population of the United States as 3×10^8 and the population of the world as 7×10^9 and determine that the world population is more than 20 times larger than the United States' population.</p> <p>M08.B-E.1.1.4 Perform operations with numbers expressed in scientific notation, including problems where both decimal and scientific notation are used. Express answers in scientific notation and choose units of appropriate size for measurements of very large or very small quantities (e.g., use millimeters per year for seafloor spreading). Interpret scientific notation that has been generated by technology (e.g., interpret 4.7EE9 displayed on a calculator as 4.7×10^9).</p> <p>CC.2.2.8.B.2 Use and evaluate square roots and cube roots to represent solutions to equations</p>

M08.B-E.2.1.1 Graph proportional relationships, interpreting the unit rate as the slope of the graph. Compare two different proportional relationships represented in different ways. Example: Compare a distance-time graph to a distance-time equation to determine which of two moving objects has greater speed.

M08.B-E.2.1.2 Use similar right triangles to show and explain why the slope m is the same between any two distinct points on a non-vertical line in the coordinate plane.

M08.B-E.2.1.3 Derive the equation $y = mx$ for a line through the origin and the equation $y = mx + b$ for a line intercepting the vertical axis at b .

CC.2.2.8.B.3 Analyze and describe linear relationships between two variables, using slope.

M08.B-E.3.1.1 Write and identify linear equations in one variable with one solution, infinitely many solutions, or no solutions. Show which of these possibilities is the case by successively transforming the given equation into simpler forms until an equivalent equation of the form $x = a$, $a = a$, or $a = b$ results (where a and b are different numbers)

M08.B-E.3.1.2 Solve linear equations that have rational number coefficients, including equations whose solutions require expanding expressions using the distributive property and collecting like terms.

M08.B-E.3.1.3 Interpret solutions to a system of two linear equations in two variables as points of intersection of their graphs because points of intersection satisfy both equations simultaneously.

M08.B-E.3.1.4 Solve systems of two linear equations in two variables algebraically and estimate solutions by

		<p>graphing the equations. Solve simple cases by inspection. Example: $3x + 2y = 5$ and $3x + 2y = 6$ have no solution because $3x + 2y$ cannot simultaneously be 5 and 6.</p> <p>M08.B-E.3.1.5 Solve real-world and mathematical problems leading to two linear equations in two variables. Example: Given coordinates for two pairs of points, determine whether the line through the first pair of points intersects the line through the second pair.</p>
<p>Functions</p>	<p>12 Days</p>	<p>CC.2.2.8.C.1 Define, interpret, and compare functions displayed algebraically, graphically, numerically in tables, or by verbal descriptions.</p> <p>M08.B-F.1.1.1 Determine whether a relation is a function.</p> <p>M08.B-F.1.1.2 Determine whether a relation is a function.</p> <p>M08.B-F.1.1.3 Interpret the equation $y = mx + b$ as defining a linear function whose graph is a straight line; give examples of functions that are not linear.</p> <p>CC.2.2.8.C.2 Use concepts of functions to model relationships between quantities.</p> <p>M08.B-F.2.1.1 Construct a function to model a linear relationship between two quantities. Determine the rate of change and initial value of the function from a description of a relationship or from two (x, y) values, including reading these from a table or from a graph. Interpret the rate of change and initial value of a linear function in terms of the situation it models and in terms of its graph or a table of values.</p> <p>M08.B-F.2.1.2 Describe qualitatively the functional relationship between two quantities by analyzing a graph (e.g., where the function is increasing or</p>

		decreasing, linear or nonlinear). Sketch or determine a graph that exhibits the qualitative features of a function that has been described verbally.
Geometry	31 Days	<p>CC.2.3.8.A.1 Apply concepts of volume of cylinders, cones, and spheres to solve real-world and mathematical problems.</p> <p>M08-C-G.3.1.1 Apply formulas for the volumes of cones, cylinders, and spheres to solve real-world and mathematical problems. Formulas will be provided.</p> <p>CC.2.3.8.A.2 Use transformations to demonstrate congruence and similarity of geometric figures.</p> <p>M08-C-G.1.1.1 Identify and apply properties of rotations, reflections, and translations. Example: Angle measures are preserved in rotations, reflections, and translations.</p> <p>M08-C-G.1.1.2 Given two congruent figures, describe a sequence of transformations that exhibits the congruence between them.</p> <p>M08-C-G.1.1.3 Describe the effect of dilations, translations, rotations, and reflections on two-dimensional figures using coordinates.</p> <p>M08-C-G.1.1.4 Given two similar two-dimensional figures, describe a sequence of transformations that exhibits the similarity between them.</p> <p>CC.2.3.8.A.3 Apply the Pythagorean Theorem and its converse to solve mathematical problems in two and three dimensions.</p> <p>M08-C-G.2.1.1 Apply the converse of the Pythagorean theorem to show a triangle is a right triangle.</p>

		<p>M08-C-G.2.1.2 Apply the Pythagorean theorem to determine unknown side lengths in right triangles in real-world and mathematical problems in two and three dimensions. (Figures provided for problems in three dimensions will be consistent with Eligible Content in grade 8 and below.)</p> <p>M08-C-G.2.1.3 Apply the Pythagorean theorem to find the distance between two points in a coordinate system.</p>
Statistics and Probability	15 Days	<p>CC.2.4.8.B.1 Construct, analyze, and interpret bivariate data displayed in scatter plots.</p> <p>M08-D-S.1.1.1 Construct and interpret scatter plots for bivariate measurement data to investigate patterns of association between two quantities. Describe patterns such as clustering, outliers, positive or negative correlation, linear association, and nonlinear association.</p> <p>M08-D-S.1.1.2 For scatter plots that suggest a linear association, identify a line of best fit by judging the closeness of the data points to the line.</p> <p>M08-D-S.1.1.3 Use the equation of a linear model to solve problems in the context of bivariate measurement data, interpreting the slope and intercept. Example: In a linear model for a biology experiment, interpret a slope of 1.5 cm/hr as meaning that an additional hour of sunlight each day is associated with an additional 1.5 cm in mature plant height.</p> <p>CC.2.4.8.B.2 Understand that patterns of association can be seen in bi-variate data utilizing frequencies.</p> <p>M08-D-S.1.2.1 Construct and interpret a two-way table summarizing data on two categorical variables collected from the same subjects. Use relative frequencies calculated for rows or columns to describe possible associations between the two variables.</p>

		Example: Given data on whether students have a curfew on school nights and whether they have assigned chores at home, is there evidence that those who have a curfew also tend to have chores?
--	--	--

RESOURCES: GLENCOE MATH, Accelerated Math copyright 2015
connectED.mcgraw-hill.com

Rev.: (July 2015)