

Keep Me at Home if



<p>Vomiting: 2 or more Occasions during the last 24 hours.</p>	<ol style="list-style-type: none"> 1. I have a rash, don't feel well and I'm itching. I have a rash and fever. 2. Open Sores should be covered. 3. 24 hours have passed after prescribed treatment begun, if it was prescribed. 4. lice: treatment against lice was done. 	<p>Diarrhea that occurs 2 or more times during the last 24 hours. And any time there is blood seen in the stool. Diarrhea with fever</p>	<p>Any suspected communicable infection such as:</p> <ul style="list-style-type: none"> • Skin infections: Impetigo, MRSA, scabies • Respiratory infections The flu, Whooping Cough <p>Alert: Before child can return there must be at least 24 hours after prescription treatment was started or a note from the doctor.</p>	<p>Sore Throat and</p> <ul style="list-style-type: none"> • Poor appetite • Fever • Irritability 	<p>I don't feel well</p> <p>I am very tired pale Poor appetite Confusion I'm irritable I have a fever</p>	<p>Fever of at least 100 degrees F under the armpit or oral 101 and I have one or more of the following:</p> <ul style="list-style-type: none"> • Headache • Ear pain • Rash • Fatigue • Sore Throat <p>Alert: Fever without signs or symptoms in infants under 2 months need medical attention immediately.</p>
---	---	---	---	--	--	---

Keep Me at Home if



- When your child is ill:**
1. Have plans for childcare for your child.
 2. Notify your child's teacher, even if he/she stays at home.