



Cave Creek Unified School District #93

School Health Office: BMES DSES DWES HTES
Phone: 480-575-2102 480-575-2902 480-575-2802 480-272-8502
Fax: 480-488-6708 480-502-2364 480-419-7265 480-907-6643

 LMES STMS CSHS
Phone: 480-437-3002 480-272-8602 480-575-2402
Fax: 480-595-1312 480-272-8699 480-488-6701

GLUCAGON EMERGENCY INJECTION/Nasal Spray

Date of Plan: _____ **Effective Dates:** _____

Student's Name: _____

Date of Birth: _____ **Grade:** _____ **Teacher:** _____

Parent signature/name/phone number: _____

Severe Low Blood Sugar:

Child will be unresponsive, maybe convulsing and unconscious. **This is a life-threatening situation.**

If you find a diabetic in this condition:

• Assign someone to call 911 and Health Office extension: _____

• Put child on his/her side, protect the student's head, and **do not** put anything in his /her mouth.

> **Instructions for injectable glucagon**

- Give Glucagon injection: Mix entire contents of accompanying syringe full of diluting solution into the vial containing glucagon. Mix well and gently withdraw contents into the syringe. With needle pointing up discard all air and any excess solution, over prescribed amount, in syringe. Inject all of the syringe contents into the child's thigh, pause for count of five before removing syringe. (Note syringe is perpendicular to thigh)
- Call parents
- Keep warm, on side; do not give anything by mouth. Vomiting may occur.

> **Instructions for nasal glucagon (BAQSIMI)**

- Remove the Shrink Wrap by pulling on the red stripe.
- Open the Lid and remove the Device from the Tube.
Caution: Do not press the Plunger until ready to give the dose.
- **Hold Device** between fingers and thumb. **Do not** push Plunger yet.
- **Insert Tip** gently in one nostril until finger(s) touch the outside of the nose.
- **Push Plunger** firmly all the way in. **Dose is complete when the Green Line disappears.**

After giving BAQSIMI (nasal spray)

- **If you have not already, call for emergency medical help right away**
- If the person is unconscious, turn the person on their side
- **Throw away the used Device and Tube**
- Encourage the person to eat as soon as possible. When they are able to safely swallow, give the person a fast-acting source of sugar, such as juice. Then encourage the person to eat a snack, such as crackers with cheese or peanut butter
- If the person does not respond after 15 minutes, another dose may be given, if available

➤ **Auto injection - Gvoke (glucagon injection) instructions**



1

Pull red

cap off



2

Push yellow

end down on skin and hold 5 seconds. Window will turn red.

Administer into upper arm, stomach, or thigh.

Call 911, roll patient on their side, keep warm.