



Reasons to Keep Your Child Home from School

You should keep your child at home if he/she might spread an illness to other children.
Children who are sick have a difficult time participating in class and learning.

<p><u>Fever</u> Children with a temperature of 100.4°F (38°C) or higher should stay at home until there is no fever for 24 hours without the use of fever-reducing medications. If your child returns to school with a fever, he/she could still be contagious.</p>	<p><u>COVID-19</u> Notify the school immediately if your child has tested positive or has been in contact with someone who has COVID-19. Follow the school's return guidance for COVID-19.</p>	<p><u>Vomiting</u> Any child who has vomited twice in 24 hours should stay home. Any child who has vomited as a result of a possible infectious cause should stay home. Your child may return to school after 24 hours of NO vomiting or fever.</p>	<p><u>Diarrhea</u> Any child who has watery diarrhea more than once or has diarrhea as a result of a possible infectious cause should stay home. Your child may return after 24 hours of NO diarrhea. Call your child's doctor if fever, rash, dehydration, or weakness are present.</p>
<p><u>Persistent Cough</u> If your child's cough is worse than you would expect with a cold, keep your child home from school. If he/she suddenly develops a cough, has a hard time breathing, or has a fever, call your child's doctor.</p>	<p><u>Sore Throat</u> If your child has a severe sore throat or a sore throat with a fever, headache, or stomachache, keep your child home and call your child's doctor.</p>	<p><u>Pain</u> (ear, head, stomach) If pain disrupts your child's participation in regular activities or is accompanied by fever, nausea/vomiting, or sore throat, keep your child home and call your child's doctor.</p>	<p><u>Chickenpox</u> Your child should stay at home until all bumps are scabbed and no new bumps have shown up for two days. **Most children have received the vaccine for chickenpox (Varicella vaccine)</p>
<p><u>Pink eye/Conjunctivitis</u> If your child has a red eye(s) with drainage that is cloudy or yellow, crusting, and/or pain, call your child's doctor. If your child has bacterial conjunctivitis, he/she may return after beginning treatment.</p>	<p><u>Scabies</u> Your child should stay home for 24 hours after treatment. Scabies is a contagious disease caused by a mite. Itching and a rash will be noted.</p>	<p><u>Head Lice</u> Keep your child home and treat, making sure all live lice are eliminated. Nit removal will need to be done daily. Contact your school for more information. **Child cannot return to school with live lice.</p>	<p><u>Impetigo</u> If your child has impetigo (red, oozing blister areas with yellow-gold scabs on the body or face), he/she should stay home until a doctor says it's OK for the child to return.</p>

When there is doubt in your mind about sending your child to school, consult your child's doctor.
Make sure that your child's school knows how to reach you during the day.
Remember to update your phone contact numbers throughout the school year.