



# Lead Poisoning Prevention

---

## Where is Lead Found?

- **Paint:** Lead can be found in paint that was made before 1978. This paint can be on any painted surface in your home, like doors, windows, and porches.
  - **Dust:** Lead dust in the home is comes from lead painted surfaces that are chipping and peeling. Sanding and scraping old paint when repainting or remodeling can also cause a lead dust problem.
  - **Soil:** Old paint that has fallen off the outside of your house onto the ground may have left lead in the soil.
  - Lead can also be found in ceramic dishes, crystal, food cans from outside the U.S., water pipes, solder and fittings, and some ethnic cosmetics and home remedies.
  - Some jobs and hobbies can expose children and adults to lead. Some examples are painters, house remodelers, plumbers, mechanics, bridge workers, making jewelry, ceramic/pottery or stained glass, and going to indoor firing ranges.
- 

## Is Your Child At Risk For Lead Poisoning?

If you answer yes to any of these questions you may want to have your child tested, even if your child is older.

- Does your child live in or often visit a building built before 1960?
  - Does your child live in or often visit a building built before 1978 that is being or was just repaired or remodeled?
  - Does your child live in or often visit a building that has peeling or chipping paint?
  - Does your child live with an adult or often visit an adult whose job or hobby exposes them to lead?
  - Does your family eat or drink from dishes made outside the U.S.?
  - Does your family use home remedies?
- 

## How does a child get lead poisoned?

- Lead poisoning usually happens when children ingest (eat) dust that has lead in it. Children may also eat chips of lead paint or soil that has lead in it.
- 

## What Does Lead Do to the Body?

- No amount of lead in the body is safe. The damage lead can cause is forever! Lead can damage the brain. It can cause growth problems, hearing loss, and learning problems.
  - Many children do not show signs of lead poisoning. Some signs of high levels of lead poisoning are the same as other childhood illnesses, like the common cold or teething.
-

## What Does Lead Do to the Body? (continued)

- If a pregnant woman is around lead, she and her unborn child may become lead poisoned. Lead can cause lasting damage to the mother and her baby.

---

## How Can You Reduce The Risk?

Replace, fix or manage all lead hazards in a lead safe way.

### Steps you can take to prevent children from being lead poisoned:

- Keep children and pregnant women away from all lead hazards.
- Clean up lead dust and paint chips by wet wiping window sills and window wells and wet mopping floors. Do NOT dry sweep or vacuum, this will spread the lead dust.
- Block places with peeling or chipping paint. Do not use windows that have chipping paint.
- Move your child's bedroom or play area to a room that has no peeling or chipping paint.
- Place washable mats inside and outside entry doors.
- Have people remove their shoes before coming in the home.
- Do not let your child (or pet) play in dirt.
- Wash and dry your child's hands, toys and pacifiers often. Wash and dry your child's hands before playing, eating, and bedtime.
- Use cold water from the tap for drinking, cooking and making formula. Let water run for 1-2 minutes before using.
- Give your child healthy meals and snacks to eat. An empty stomach takes in lead faster than a full stomach.

### Steps adults can take to help prevent themselves or children from becoming lead poisoned from their job or hobby:

- Don't eat, drink or smoke in your work/hobby area.
- Wash your hands and face before eating, smoking or drinking.
- Wear protective clothing (such as disposable gloves, hat, and shoe covers) when you work with lead. Use a NIOSH-approved respirator.
- Shower, wash your hair, and change into clean clothes and shoes before you leave the work area. Leaving dust on your clothes can contaminate your home and car.
- Put your work clothes and shoes in sealed plastic bags.
- Wash work clothes in a different load than the family's laundry.

---

## Does your child need to be tested for lead poisoning?

- Yes, all children, at about ages one and two, must be tested for lead poisoning...it's the law!
- Blood tests will tell how much lead is in your child's blood at the time of the test. If the level is high, your child will need more testing.
- If your child is at risk at other ages, have your child tested at those times too.

---

## Connecticut Lead and Healthy Homes Program

(860) 509-7299

[www.ct.gov/dph/lead](http://www.ct.gov/dph/lead)

[www.ct.gov/dph/healthyhomes](http://www.ct.gov/dph/healthyhomes)





# Eating Right Helps Fight Lead Poisoning

---

Lead tricks the body into thinking it is iron, calcium or zinc. Eating healthy can help decrease the lead from staying in the body.

Don't let your child go through the day on an empty stomach!

---

## Five Basic Food Groups

- Breads, cereals and grains
- Vegetables
- Fruit
- Milk and milk products
- Meat, chicken, fish, nuts, and beans



## Foods Rich in Calcium

- Milk
- Yogurt
- Cheese (*for snacks, in cooking such as macaroni and cheese, pizza, tortillas, vegetables*)
- Foods made of milk (*pudding, soup, ice cream, custard*)
- Sardines or canned salmon (*with bones*)
- Green vegetables (*kale, collard greens, broccoli*)



## Foods Rich in Zinc

- Chicken or turkey
- Lean meat
- Fish
- Milk and cheese
- Clams, oysters, mussels, crab
- Dried beans and lentils
- Eggs



---

## Foods Rich in Iron

- Lean red meat, chicken, turkey and fish
- Iron-fortified hot and cold cereals
- Clams, oysters or mussels (*use canned to make soup or sauce for pasta*)
- Dark green leafy vegetables
- Dried beans, split peas, and other beans (*pinto, red, navy, kidney, garbanzo*)
- Eggs
- Dried fruit



*The iron in vegetables, grains, beans, nuts and eggs may be made more usable to the body when you eat a food high in Vitamin C at the same meal. Oranges, grapefruit, strawberries, cantaloupe, green peppers, cauliflower, broccoli and potatoes are some foods high in Vitamin C.*

---

### Healthy Tips:

- Don't fry foods. Bake or broil them.
  - Try not to eat high fat foods. When you do eat them, eat small portions.
  - Vitamin C helps your body absorb iron.
  - Children under the age of 2 should have whole milk after they no longer drink formula or breast milk. Most children 2 and older can have lower fat milk. Children with milk allergies can have tofu, leafy green vegetables, sardines, or canned salmon for their calcium needs.
  - Younger children need smaller servings than older children or adults. More active people need larger numbers of servings from each of the 5 food groups.
- 

**Connecticut Lead and Healthy Homes Program**

**(860) 509-7299**

**[www.choosemyplate.gov](http://www.choosemyplate.gov)**

**[www.ct.gov/dph/lead](http://www.ct.gov/dph/lead)**

**[www.ct.gov/dph/healthyhomes](http://www.ct.gov/dph/healthyhomes)**



---

**If you require aid/accommodation to fully and fairly benefit from the contents of this document,  
please contact 860-509-7293**



# Imported Home Remedies & Medicines and Lead Poisoning

---

People use home remedies and medicines to try to help someone feel better. Many of the **IMPORTED** brands have lead in them and can make people very sick and cause lead poisoning.

---

**These IMPORTED home remedies are known to have lead in them.**

**They are very dangerous!**

---

<b>Asafoetida:</b>	Spice is used as a digestive aid, in food as a condiment, and in pickles. It typically works as a flavor enhancer and, used along with turmeric, is a standard component of Indian recipes.
<b>Azarcon:</b>	A bright orange powder. It is also called Rueda, Coral, Maria Luisa, Alarcon, and Liga. It is used for an upset stomach or indigestion (“empacho”). It has almost 100% lead.
<b>Albayalde:</b>	A white powder. It is used for an upset stomach or indigestion (“empacho”). It may have up to 37% lead.
<b>Greta:</b>	A yellow powder. It is used for an upset stomach or indigestion (“empacho”). It has almost 100% lead.
<b>Pay~loo~ah:</b>	A red powder. It is used for rashes or fever.
<b>Ghasard:</b>	A brown powder. It is used to aid digestion.
<b>Bala Goli:</b>	A round, flat, black bean that is dissolved in “gripe water.” It is used to treat stomach aches.
<b>Kandu:</b>	A red powder. It is used to treat stomach aches.
<b>Kohl, Kajal Alkohl, Surma:</b>	A black powder. It is used as an eye make-up. It is also put on skin infections and on the umbilical stump of newborn

---

**Any amount of these IMPORTED home remedies is poisonous to children and adults.**

---

---

## How do I know if the imported home remedy I have has lead in it?

- It is difficult to tell if it has lead in it.
- A product may contain high levels of lead even if lead is not listed as an ingredient on the label.
- The only way to know is to test.
- **If your child is lead poisoned provide the home remedy to the local health department for testing.**

---

## What should I do if I or my child has taken an imported home remedy that might have lead in it?

- Stop using the remedy. Keep it away from children.
- Contact your doctor or clinic immediately to get their advice.
- Think about getting you and your child tested for lead poisoning.  
**Please note:** It's the law that all children be tested for lead at 1 year and at 2 years of age, even if they have not taken one of the home remedies that may contain lead.
- Many children can get free medical care. Contact the HUSKY program at 1-877-CTHUSKY (1-877-284-8759) for more information.

---

## What remedies or medicines should I use instead?

- Use U.S. made remedies and medicines.

---

Connecticut Lead and Healthy Homes Program

(860) 509-7299

[www.ct.gov/dph/lead](http://www.ct.gov/dph/lead)

[www.ct.gov/dph/healthyhomes](http://www.ct.gov/dph/healthyhomes)



---

If you require aid/accommodation to fully and fairly benefit from the contents of this document,  
please contact 860-509-7293