

Number Sense Tricks

Bryant Heath

18 October 2007

Contents

1	Numerical Tricks	6
1.1	Introduction: FOILing When Multiplying	6
1.2	Multiplying: The Basics	8
1.2.1	Multiplying by 11 Trick	8
1.2.2	Multiplying by 101 Trick	10
1.2.3	Multiplying by 25 Trick	11
1.2.4	Multiplying by 75 trick	12
1.2.5	Multiplying by Any Fraction of 100, 1000, etc...	13
1.2.6	Double and Half Trick	16
1.2.7	Multiplying Two Numbers Near 100	17
1.2.8	Squares Ending in 5 Trick	19
1.2.9	Squares from 41-59	20
1.2.10	Multiplying Two Numbers Equidistant from a Third Number	21
1.2.11	Multiplying Reverses	23
1.3	Standard Multiplication Tricks	23
1.3.1	Extending Foiling	23
1.3.2	Factoring of Numerical Problems	25
1.3.3	Sum of Consecutive Squares	28
1.3.4	Sum of Squares: Factoring Method	28
1.3.5	Sum of Squares: Special Case	29
1.3.6	Difference of Squares	30
1.3.7	Multiplying Two Numbers Ending in 5	32
1.3.8	Multiplying Mixed Numbers	32
1.3.9	$a \times \frac{a}{b}$ Trick	34
1.3.10	Combination of Tricks	36
1.4	Dividing Tricks	37
1.4.1	Finding a Remainder when Dividing by 4,8, etc...	37
1.4.2	Finding a Remainder when Dividing by 3,9, etc...	38
1.4.3	Finding a Remainder when Dividing by 11	38
1.4.4	Finding Remainders of Other Integers	39
1.4.5	Remainders of Expressions	40
1.4.6	Dividing by 9 Trick	42
1.4.7	Converting $\frac{a}{40}$ and $\frac{b}{80}$, etc... to Decimals	43
1.5	Adding and Subtracting	44
1.5.1	Subtracting Reverses	44
1.5.2	Switching Numbers and Negating on Subtraction	45
1.5.3	$\frac{a}{b \cdot (b+1)} + \frac{b}{(b+1) \cdot (b+2)} + \dots$	46
1.5.4	$\frac{a}{b} + \frac{b}{a}$ Trick	46
1.5.5	$\frac{a}{b} - \frac{na-1}{nb+1}$	48
2	Memorizations	50
2.1	Important Numbers	50
2.1.1	Squares	50
2.1.2	Cubes	52
2.1.3	Powers of 2, 3, 5	54
2.1.4	Important Fractions	56
2.1.5	Special Integers	60
2.1.6	Roman Numerals	62
2.1.7	Platonic Solids	64
2.1.8	π and e Approximations	64

2.1.9	Distance Conversions	65
2.1.10	Conversion between Distance \rightarrow Area, Volume	66
2.1.11	Fluid and Weight Conversions	67
2.1.12	Celsius to Fahrenheit Conversions	68
2.2	Formulas	69
2.2.1	Sum of Series	69
2.2.2	Fibonacci Numbers	72
2.2.3	Integral Divisors	75
2.2.4	Number of Diagonals of a Polygon	78
2.2.5	Exterior/Interior Angles	78
2.2.6	Triangular, Pentagonal, etc... Numbers	79
2.2.7	Finding Sides of a Triangle	81
2.2.8	Equilateral Triangle Formulas	83
2.2.9	Formulas of Solids	84
2.2.10	Combinations and Permutations	85
2.2.11	Trigonometric Values	86
2.2.12	Trigonometric Formulas	90
2.2.13	Graphs of Sines/Cosines	91
2.2.14	Vertex of a Parabola	92
2.2.15	Discriminant and Roots	93
3	Miscellaneous Topics	94
3.1	Random Assortment of Problems	94
3.1.1	GCD/GCF and LCM	94
3.1.2	Perfect, Abundant, and Deficient Numbers	96
3.1.3	Sum and Product of Coefficients in Binomial Expansion	96
3.1.4	Sum/Product of the Roots	98
3.1.5	Finding Units Digit of x^n	99
3.1.6	Exponent Rules	101
3.1.7	Log Rules	103
3.1.8	Square Root Problems	105
3.1.9	Finding Approximations of Square Roots	106
3.1.10	Complex Numbers	107
3.1.11	Function Inverses	109
3.1.12	Patterns	110
3.1.13	Probability and Odds	111
3.1.14	Sets	113
3.2	Changing Bases	114
3.2.1	Converting Integers	114
3.2.2	Converting Decimals	117
3.2.3	Performing Operations	118
3.2.4	Changing Between Bases: Special Case	120
3.2.5	Changing Bases: Sum of Powers	122
3.2.6	Changing Bases: Miscellaneous Topics	122
3.3	Repeating Decimals	123
3.3.1	In the form: .aaaaa...	123
3.3.2	In the form: .ababa...	123
3.3.3	In the form: .abbbb...	124
3.3.4	In the form: .abcbcbc...	125
3.4	Modular Arithmetic	125
3.5	Fun with Factorials!	126
3.5.1	$1 \cdot 1! + 2 \cdot 2! + \cdots + n \cdot n!$	127
3.5.2	$\frac{a! \pm b!}{c!}$	127
3.5.3	Wilson's Theorem	128

3.6	Basic Calculus	129
3.6.1	Limits	129
3.6.2	Derivatives	130
3.6.3	Integration	131
4	Additional Problems	134
5	Solutions	135

Introduction

As most who are reading this book already know, the UIL Number Sense exam is an intense 10 minute test where 80 challenging mental math problems test a student's knowledge of topics ranging from simple multiplication, geometry, algebraic manipulation, to calculus. Although the exam is grueling (with 7.5 seconds per problem, it is hard to imagine it being easy!), there are various tricks to alleviate some of the heavy computations associated with the test. The purpose of writing this book is to explore a variety of these "shortcuts" as well as their applications in order to better prepare the student for taking the Number Sense test. In addition, this book is a source of practice problems so that better proficiency with many different types of problems can be reached leaving more time for the harder and more unique test questions.

The book will be divided into three sections: numerical tricks, necessary memorizations (ranging from conversions to formulas), and miscellaneous topics. The difficulty of tricks discussed in the text range from some of the most basic (11's trick, Subtracting Reverses, etc...) to the more advanced that are on the most recent exams. Although this book will provide, hopefully, adequate understanding of a wide variety of commonly used tricks, it is **not** a replacement for practicing and discovering methods that you feel the most comfortable with. In order to solidify everything exhibited in this book, regular group and individual practice sessions are recommended as well as participation in multiple competitions.

The best way to approach this book would be to read through all the instructional material first then go back and do the practice problems in each section. The reason why this is needed is because many sections deal with combinations of problems which are discussed later in the book. Also, all problems in **bold** reflect questions taken from the state competition exams. Similarly, to maintain consistent nomenclature, all (*) problems are approximation problems where $\pm 5\%$ accuracy is needed.

It should be noted that the tricks exhibited here could very easily not be the faster method for doing problems. I wrote down tricks and procedures that I follow, and since I am only human, there could very easily be faster more to-the-point tricks that I haven't noticed. In fact, as I've been gleaning past tests to find sample problems I've noticed faster methods on how to do some problems (and I've updated the book accordingly). One of the reasons why Number Sense is so great is that there is usually a variety of methods which can be used! This is apparent mostly in the practice problems. I tried to choose problems which reflects the procedures outlined in each section. Sometimes you can employ different methods and come up with an equally fast (or possibly faster depending on which method you prefer) way of solving the problems, so do whatever way you work the fastest and feel the most comfortable with.

1 Numerical Tricks

1.1 Introduction: FOILing When Multiplying

Multiplication is at the heart of every number sense test. Slow multiplication hampers how far you are able to go on the test as well as decreasing your accuracy. To help beginners learn how to speed up multiplying, the concept of FOILing, learned in beginning algebra classes, is introduced as well as some exercises to help in speeding up multiplication. What is nice about the basic multiplication exercises is that *anyone* can make up problems, so practice is unbounded.

When multiplying two two-digit numbers ab and cd swiftly, a method of FOILing (First-Outer+Inner-Last) is used. To understand this concept better, lets take a look at what we do when we multiply $ab \times cd$:

$$ab = 10a + b \text{ and } cd = 10c + d$$
$$(10a + b) \times (10c + d) = 100(ac) + 10(ad + bc) + bd$$

A couple of things can be seen by this:

1. The one's digit of the answer is simply bd or the *First* digits (by first I mean the least significant digit) of the two numbers multiplied.
2. The ten's digit of the answer is $(ad + bc)$ which is the sum of the *Outer* digits multiplied together plus the *Inner* digits multiplied.
3. The hundred's digit is ac which are the *Last* digits (again, by last I mean the most significant digit) multiplied with each other.
4. If in each step you get more than a single digit, you carry the extra (most significant digit) to the next calculation. For example:

	Units:	$3 \times 4 = 12$
$74 \times 23 =$	Tens:	$3 \times 7 + 2 \times 4 + 1 = 30$
	Hundreds:	$2 \times 7 + 3 = 17$
	Answer:	1702

Where the bold represents the answer and the italicized represents the carry.

Similarly, you can extend this concept of FOILing to multiply any n-digit number by m-digit number in a procedure I call "moving down the line." Let's look at an example of a 3-digit multiplied by a 2-digit:

	Ones:	$3 \times 3 = 9$
	Tens:	$3 \times 9 + 2 \times 3 = 33$
$493 \times 23 =$	Hundreds:	$3 \times 4 + 2 \times 9 + 3 = 33$
	Thousands:	$2 \times 4 + 3 = 11$
	Answers:	11339

As one can see, you just continue multiplying the two-digit number "down the line" of the three-digit number employing the FOILing technique at each step then writing down what you get for each digit then moving on (always remembering to carry when necessary). The following is a set of exercises to familiarize you with this process of multiplication:

Problem Set 1.1:

$95 \times 30 =$

$90 \times 78 =$

$51 \times 11 =$

$83 \times 51 =$

$64 \times 53 =$

$65 \times 81 =$

$92 \times 76 =$

$25 \times 46 =$

$94 \times 92 =$

$27 \times 64 =$

$34 \times 27 =$

$11 \times 77 =$

$44 \times 87 =$

$86 \times 63 =$

$54 \times 92 =$

$83 \times 68 =$

$72 \times 65 =$

$81 \times 96 =$

$57 \times 89 =$

$25 \times 98 =$

$34 \times 32 =$

$88 \times 76 =$

$22 \times 11 =$

$36 \times 69 =$

$35 \times 52 =$

$15 \times 88 =$

$62 \times 48 =$

$56 \times 40 =$

$62 \times 78 =$

$57 \times 67 =$

$28 \times 44 =$

$80 \times 71 =$

$51 \times 61 =$

$81 \times 15 =$

$64 \times 14 =$

$47 \times 37 =$

$79 \times 97 =$

$99 \times 87 =$

$49 \times 54 =$

$29 \times 67 =$

$38 \times 98 =$

$75 \times 47 =$

$77 \times 34 =$

$49 \times 94 =$

$71 \times 29 =$

$85 \times 66 =$

$13 \times 65 =$

$64 \times 11 =$

$62 \times 15 =$

$43 \times 65 =$

$74 \times 72 =$

$49 \times 41 =$

$23 \times 70 =$

$72 \times 75 =$

$53 \times 59 =$

$82 \times 91 =$

$14 \times 17 =$

$67 \times 27 =$

$85 \times 25 =$

$25 \times 99 =$

$137 \times 32 =$

$428 \times 74 =$

$996 \times 47 =$

$654 \times 45 =$

$443 \times 39 =$

$739 \times 50 =$

$247 \times 87 =$

$732 \times 66 =$

$554 \times 77 =$

$324 \times 11 =$

$111 \times 54 =$

$885 \times 78 =$

$34 \times 655 =$

$52 \times 532 =$

$33 \times 334 =$

$45 \times 301 =$

$543 \times 543 =$

$606 \times 212 =$

$657 \times 322 =$

$543 \times 230 =$

$111 \times 121 =$

$422 \times 943 =$

$342 \times 542 =$

$789 \times 359 =$

$239 \times 795 =$

$123 \times 543 =$

$683 \times 429 =$

$222 \times 796 =$

1.2 Multiplying: The Basics

1.2.1 Multiplying by 11 Trick

The simplest multiplication trick is the 11's trick. It is a mundane version of "moving down the line," where you add consecutive digits and record the answer. Here is an example:

$$\begin{array}{rcl}
 & \text{Ones:} & 1 \times 3 = \mathbf{3} \\
 & \text{Tens:} & 1 \times 2 + 1 \times 3 = \mathbf{5} \\
 523 \times 11 = & \text{Hundreds:} & 1 \times 5 + 1 \times 2 = \mathbf{7} \\
 & \text{Thousands:} & 1 \times 5 = \mathbf{5} \\
 & \text{Answer:} & \mathbf{5753}
 \end{array}$$

As one can see, the result can be obtained by subsequently adding the digits along the number you're multiplying. Be sure to keep track of the carries as well:

$$\begin{array}{rcl}
 & \text{Ones:} & \mathbf{8} \\
 & \text{Tens:} & 9 + 8 = \mathbf{17} \\
 6798 \times 11 = & \text{Hundreds:} & 7 + 9 + 1 = \mathbf{17} \\
 & \text{Thousands:} & 6 + 7 + 1 = \mathbf{14} \\
 & \text{Ten Thousands:} & 6 + 1 = \mathbf{7} \\
 & \text{Answer:} & \mathbf{74778}
 \end{array}$$

The trick can also be extended to 111 or 1111 (and so on). Where as in the 11's trick you are adding pairs of digits "down the line," for 111 you will be adding triples:

$$\begin{array}{rcl}
 & \text{Ones:} & \mathbf{3} \\
 & \text{Tens:} & 4 + 3 = \mathbf{7} \\
 & \text{Hundreds:} & 5 + 4 + 3 = \mathbf{12} \\
 6543 \times 111 = & \text{Thousands:} & 6 + 5 + 4 + 1 = \mathbf{16} \\
 & \text{Ten Thousands:} & 6 + 5 + 1 = \mathbf{12} \\
 & \text{Hun. Thousands:} & 6 + 1 = \mathbf{7} \\
 & \text{Answer:} & \mathbf{726273}
 \end{array}$$

Another common form of the 11's trick is used in reverse. For example:

$$\begin{array}{rcl}
 & 1353 \div 11 = & \\
 & \text{or} & \\
 & 11 \times x = 1353 & \\
 \text{Ones Digit of } x \text{ is equal to the Ones Digit of } 1353: & & \mathbf{3} \\
 \text{Tens Digit of } x \text{ is equal to:} & 5 = 3 + x_{\text{tens}} & \mathbf{2} \\
 \text{Hundreds Digit of } x \text{ is equal to:} & 3 = 2 + x_{\text{hund}} & \mathbf{1} \\
 \text{Answer:} & & \mathbf{123}
 \end{array}$$

Similarly you can perform the same procedure with 111, and so on. Let's look at an example:

$$\begin{array}{rcl}
 & 46731 \div 111 = & \\
 & \text{or} & \\
 & 111 \times x = 46731 & \\
 \text{Ones Digit of } x \text{ is equal to the Ones Digit of } 46731: & & \mathbf{1} \\
 \text{Tens Digit of } x \text{ is equal to:} & 3 = 1 + x_{\text{tens}} & \mathbf{2} \\
 \text{Hundreds Digit of } x \text{ is equal to:} & 7 = 2 + 1 + x_{\text{hund}} & \mathbf{4} \\
 \text{Answer:} & & \mathbf{421}
 \end{array}$$

The hardest part of the procedure is knowing when to stop. The easiest way I've found is to think about how many digits the answer *should* have. For example, with the above expression, we are dividing a 5-digit number by a roughly 100, leaving an answer which should be 3-digits, so after the third-digit you know you

are done.

The following are some more practice problems to familiarize you with the process:

Problem Set 1.2.1.:

1. $11 \times 54 =$
2. $11 \times 72 =$
3. $11 \times 38 =$
4. $462 \times 11 =$
5. $11 \times 74 =$
6. $66 \times 11 =$
7. $1.1 \times 2.3 =$
8. $52 \times 11 =$
9. $246 \times 11 =$
10. $111 \times 456 =$
11. $198 \div 11 =$
12. $357 \times 11 =$
13. $275 \div 11 =$
14. $321 \times 111 =$
15. $1.1 \times .25 =$
16. $111 \times 44 =$
17. $374 \div 11 =$
18. $87 \times 111 =$
19. $286 \div 11 =$
20. $111 \times 53 =$
21. $297 \div 11 =$
22. $2233 \div 11 =$
23. $198 \times 11 =$
24. $297 \div 11 =$
25. $111 \times 41 =$
26. $111 \times 35 =$
27. $111 \times 345 =$
28. $2003 \times 111 =$
29. $3 \times 5 \times 7 \times 11 =$
30. $121 \times 121 =$
31. $33 \times 1111 =$
32. $22 \times 32 =$
33. $36963 \div 111 =$
34. $20.07 \times 1.1 =$
35. 11% of 22% is: _____% (dec.)

$36. 13 \times 121 =$

$48. 55 \times 33 =$

$37. 27972 \div 111 =$

$49. (*) 32 \times 64 \times 16 \div 48 =$

$38. 2006 \times 11 =$

$50. 2002 \div 11 =$

$39. 11^4 =$

$51. 77 \times 88 =$

$40. 33 \times 44 =$

$52. (*) 44.4 \times 33.3 \times 22.2 =$

$41. 2 \times 3 \times 11 \times 13 =$

$53. 11 \times 11 \times 11 \times 11 =$

$42. 121 \times 22 =$

$54. \mathbf{25553} \div \mathbf{1111} =$

$43. 44 \times 55 =$

$55. 11 \times 13 \times 42 =$

$44. 2 \times 3 \times 5 \times 7 \times 11 =$

$56. 1111 \times 123 =$

$45. 2553 \div 111 =$

$57. 2 \times 5 \times 3 \times 7 \times 11 =$

$46. \mathbf{114} \times \mathbf{121} =$

$58. 121 \times 124 =$

$47. 44 \times 25 \times 11 =$

$59. (*) 33 \times 44 \times 55 =$

1.2.2 Multiplying by 101 Trick

In the same spirit as the multiplying by 11's trick, multiplying by 101 involves adding gap connected digits. Let's look at an example:

$438 \times 101 =$	Ones:	1×8	8
	Tens:	1×3	3
	Hundreds:	$1 \times 4 + 1 \times 8$	12
	Thousands:	$1 \times 3 + 1$	4
	Tens Thousands:	1×4	4
	Answer:		44238

So you see, immediately you can write down the ones/tens digits (they are the same as what you are multiplying 101 with). Then you sum gap digits and move down the line. Let's look at another example:

$8234 \times 101 =$	Ones/Tens:	34	34
	Hundreds:	$2 + 4$	6
	Thousands:	$8 + 3$	11
	Tens Thousands:	$2 + 1$	3
	Hundred Thousands:	8	8
	Answer:		831634

Problem Set 1.2.2

1. $1234 \times 101 =$

6. $202 \times 123 =$

2. $10.1 \times 234 =$

7. If 6 balls cost \$6.06, then 15 balls cost: \$_____

3. $369 \times 101 =$

8. $404 \times 1111 =$

4. $34845 \div 101 =$

9. (*) $(48 + 53) \times 151 =$

5. $22422 \div 101 =$

10. (*) $8888 \times 62.5\% \times \frac{5}{11} =$

1.2.3 Multiplying by 25 Trick

The trick to multiplying by 25 is to think of it as $\frac{100}{4}$. So the strategy is to take what ever you are multiplying with, divide it by 4 then move the decimal over to the right two places. Here are a couple of examples:

$$84 \times 25 = \frac{84}{4} \times 100 = 21 \times 100 = \mathbf{2100}$$

$$166 \times 25 = \frac{166}{4} \times 100 = 41.5 \times 100 = \mathbf{4150}$$

In a similar manner, you can use the same principle to divide numbers by 25 easily. The difference is you multiply by 4 and then move the decimal over to the left two places

$$\frac{415}{25} = \frac{415}{\frac{100}{4}} = \frac{415 \times 4}{100} = \frac{1660}{100} = 16.6$$

Problem Set 1.2.3

1. $240 \times 25 =$

5. $25 \times 33 =$

2. $25 \times 432 =$

6. $64 \div 25 =$

3. $2.6 \times 2.5 =$

7. $25 \times 147 =$

4. $148 \times 25 =$

8. $418 \times 25 =$

9. $616 \div 25 =$
10. $2.5 \times 40.4 =$
11. $1.1 \div 2.5 =$
12. $3232 \times 25 =$
13. (*) $97531 \div 246 =$
14. Which is smaller: $\frac{6}{25}$ or $.25$:
15. $209 \times 25 =$
16. $(18 + 16)(9 + 16) =$
17. (*) $334455 \div 251 =$
18. 21.4 is _____% of 25.
19. $404 \div 25 =$
20. $303 \times 25 =$
21. (*) $97531 \div 246 =$
22. Which is larger: $\frac{7}{25}$ or $.25$:
23. $2006 \div 25 =$
24. $25 \times 307 =$
25. 32 is $2\frac{1}{2}\%$ of: _____
26. (*) $47985 \div 246 =$
27. **$25 \times 2003 =$**
28. $15 \times 25 \times 11 =$
29. $11 \times 24 \times 25 =$
30. $11 \times 18 \times 25 =$
31. (*) $248 \times 250 \times 252 =$

1.2.4 Multiplying by 75 trick

In a similar fashion, you can multiply by 75 by treating it as $\frac{3}{4} \cdot 100$. So when you multiply by 75, first divide the number you're multiplying by 4 then multiply by 3 then move the decimal over two places to the right.

$$76 \times 75 = \frac{76 \cdot 3}{4} \cdot 100 = 19 \times 3 \times 100 = 5700$$

$$42 \times 75 = \frac{42 \cdot 3}{4} \cdot 100 = 10.5 \times 3 \times 100 = 3150$$

Again, you can use the same principle to divide by 75 as well, only you multiply by $\frac{4}{3}$ then divide by 100 (or move the decimal place over two digits to the left).

$$\frac{81}{75} = \frac{81}{\frac{3 \cdot 100}{4}} = \frac{81 \cdot 4}{3 \cdot 100} = \frac{27 \cdot 4}{100} = 1.08$$

Problem Set 1.2.4

1. $48 \times 75 =$

9. $48 \div 75 =$

2. $64 \times 75 =$

10. (*) $566472 \div 748 =$

3. $66 \div 75 =$

11. **$96 \div 75 =$**

4. $84 \times 75 =$

12. $75 \times 11 \times 24 =$

5. (*) $443322 \div 751 =$

13. $4800 \div 75 =$

6. $28 \times 75 =$

14. $75 \times 48 \times 15 =$

7. $75 \times 24 =$

8. (*) 7532×1468

15. $8.8 \times 7.5 \times 1.1 =$

1.2.5 Multiplying by Any Fraction of 100, 1000, etc...

You can take what we learned from the 25's and 75's trick (converting them to fractions of 100) with a variety of potential fractions. $\frac{1}{8}$'s are chosen often because:

$$125 = \frac{1}{8} \cdot 1000$$

$$37.5 = \frac{3}{8} \cdot 100$$

$$6.25 = \frac{5}{8} \cdot 10$$

In addition, you see $\frac{1}{6}$'s, $\frac{1}{3}$'s, $\frac{1}{9}$'s, and sometimes even $\frac{1}{12}$'s for approximation problems (because they do not go evenly into 100, 1000, etc..., they have to be approximated usually).

$$223 \approx \frac{2}{9} \cdot 1000$$

$$8333.3 \approx \frac{5}{6} \cdot 10000 \approx \frac{1}{12} \cdot 100000$$

$$327 \approx \frac{1}{3} \cdot 1000$$

For approximations you will rarely ever see them almost exact to the correct fraction. For example you might use $\frac{2}{3} \cdot 1000$ for any value from 654 – 678. Usually you can tell for the approximation problems what fraction the test writer is really going for. Before doing the problem set, it is recommended to at least familiarize yourself with the fractions in Section 2.1.4.

Problem Set 1.2.5

1. $125 \times 320 =$
2. (*) $8333 \times 24 =$
3. $138 \div 125 =$
4. (*) $57381 \div 128 =$
5. (*) $245632 \div 111 =$
6. (*) **$16667 \div 8333 \times 555 =$**
7. $625 \times 320 =$
8. (*) $774447 \div 111 =$
9. (*) $62.5 \times 3248 =$
10. $12.5 \times 480 =$
11. (*) $17304 \div 118 =$
12. (*) $87\% \text{ of } 5590 =$
13. (*) $457689 \div 111 =$
14. (*) $625 \times 648 =$
15. $375 \times 408 =$
16. (*) $359954 \div 1111 =$
17. $88 \times 12.5 \times .11 =$
18. (*) $719 \times 875 =$
19. (*) $428571 \times 22 =$
20. (*) **$85714.2 \div 714.285 =$**
21. $488 \times 375 =$
22. (*) $6311 \times 1241 =$
23. (*) $884422 \div 666 =$
24. (*) $106.25\% \text{ of } 640 =$
25. (*) **$6388 \times 3.75 =$**
26. $240 \times 875 =$
27. (*) $12.75 \times 28300 \div 102 =$
28. $375 \times 24.8 =$
29. (*) $857142 \times 427 =$
30. $.0625 \times .32 =$
31. (*) $16667 \times 369 =$
32. (*) $918576 \div 432 =$
33. (*) $456789 \div 123 =$
34. (*) $106\% \text{ of } 319 =$
35. (*) $571428 \times .875 =$
36. (*) $123\% \text{ of } 882 =$
37. (*) $95634 \div 278 =$
38. (*) $273849 \div 165 =$
39. (*) **$5714.28 \times 85 =$**
40. (*) $9.08\% \text{ of } 443322 =$
41. (*) $8333 \times 23 =$
42. $.125 \times 482 =$

43. (*) $714285 \times .875 =$
44. (*) 87% of $789 =$
45. (*) $16667 \times 49 =$
46. (*) $123456 \div 111 =$
47. (*) $875421 \div 369 =$
48. (*) $71984 \times 1.371 =$
49. (*) 63% of $7191 =$
50. (*) $5714.28 \times 83 =$
51. (*) $1428.57 \times 62 =$
52. (*) $80520 \div 131 =$
53. (*) $142.857 \times 428.571 =$
54. (*) $12509 \times 635 =$
55. (*) $1234 \times 567 =$
56. (*) $789123 \div 456 =$
57. $625 \times 65 =$
58. (*) $1428.57 \times 73 =$
59. (*) $7142.85 \times 34.2 =$
60. (*) $333 \times 808 \times 444 =$
61. (*) $571428 \times 34 =$
62. (*) $833 \times 612 =$
63. (*) $8333 \times (481 + 358) =$
64. (*) $234678 \div 9111 =$
65. (*) $428.571 \times 87.5 =$
66. (*) $375.1 \times 83.33 \times 1.595 =$
67. (*) $8333 \div 6666 \times 4444 =$
68. (*) $8333 \times 12\frac{1}{2}\% \times .12 =$
69. (*) $639 \times 375 \div 28 =$
70. (*) $6250 \div 8333 \times 8888 =$
71. (*) $416666 \div 555 \times 76 =$
72. (*) $375 \div 833 \times 555 =$
73. (*) $438 \div 9\frac{1}{11}\% \times 11.1 =$
74. (*) $857142 \div 428571 \times 7777 =$
75. (*) $546 \div 45\frac{5}{11}\% \times 10.8 =$
76. (*) $54.5454 \times 66.6 \times 58 =$
77. (*) $456 \div 18.75\% \times \frac{1}{4} =$
78. (*) $818 \div 44\frac{4}{9}\% \times 12.5 =$
79. (*) $62.5 \div 83.3 \times 888 =$
80. (*) $797 \div 87.5\% \times \frac{7}{10} =$
81. (*) $888 \times 87.5\% \div \frac{7}{11} =$
82. (*) $1250 \div 1666 \times 4444 =$
83. (*) $85858 \div 585 =$
84. (*) $(51597 \div 147)^2 =$

1.2.6 Double and Half Trick

This trick involves multiplying by a clever version of 1. Let's look at an example to show the technique:

$$\begin{aligned}15 \times 78 &= \frac{2}{2} \times 15 \times 78 \\ &= 15 \cdot 2 \times \frac{78}{2} \\ &= 30 \times 39 = \mathbf{1170}\end{aligned}$$

So the procedure is you double one of the numbers and half the other one, then multiply. This trick is exceptionally helpful when multiplying by 15 or any two-digit number ending in 5. Another example is:

$$35 \times 42 = 70 \times 21 = \mathbf{1470}$$

It is also good whenever you are multiplying an even number in the teens by another number:

$$\begin{aligned}18 \times 52 &= 9 \times 104 = \mathbf{936} \\ &\text{or} \\ 14 \times 37 &= 7 \times 74 = \mathbf{518}\end{aligned}$$

The purpose of this trick is to save time on calculations. It is a lot easier to multiply a single-digit number than a two-digit number.

Problem Set 1.2.6

1. $1.5 \times 5.2 =$

10. $18 \times 112 =$

2. $4.8 \times 15 =$

11. $27 \times 14 =$

3. $64 \times 1.5 =$

12. $21 \times 15 \times 14 =$

4. $15 \times 48 =$

13. $33.75 = 1.5 \times \text{-----}$

5. $14 \times 203 =$

14. $345 \times 12 =$

6. $14 \times 312 =$

7. $24 \times 35 =$

15. $1.2 \times 1.25 =$

8. $312 \times 14 =$

16. 24% of 44 is =

9. A rectangle has a length of 2.4 and a width of 1.5. Its area is:

17. $14 \times 25 + 12.5 \times 28 =$

1.2.7 Multiplying Two Numbers Near 100

Let's look at two numbers over 100 first:

Express $n_1 = (100 + a)$ and $n_2 = (100 + b)$ where a and b are how much the number's are above 100, then:

$$\begin{aligned} n_1 \cdot n_2 &= (100 + a) \cdot (100 + b) \\ &= 10000 + 100(a + b) + ab \\ &= 100(100 + a + b) + ab \\ &= 100(n_1 + b) + ab = 100(n_2 + a) + ab \end{aligned}$$

1. The Tens/Ones digits are just the difference the two numbers are above 100 multiplied together (ab)
2. The remainder of the answer is just n_1 *plus* the amount n_2 is above 100, or n_2 *plus* the amount n_1 is above 100.

$103 \times 108 =$	Tens/Units:	8×3	24
	Rest of Answer:	$103 + 8$ or $108 + 3$	111
	Answer:		11124

Now let's look at two numbers below 100:

Similarly, $n_1 = (100 - a)$ and $n_2 = (100 - b)$ so:

$$\begin{aligned} n_1 \cdot n_2 &= (100 - a) \cdot (100 - b) \\ &= 10000 - 100(a + b) + ab \\ &= 100(100 - a - b) + ab \\ &= 100(n_1 - b) + ab = 100(n_2 - a) + ab \end{aligned}$$

1. Again, Tens/Ones digits are just the difference the two numbers are above 100 multiplied together (ab)
2. The remainder of the answer is just n_1 *minus* the difference n_2 is from 100, or n_2 *minus* the difference n_1 is from 100.

$97 \times 94 =$	Tens/Ones:	$(100 - 97) \times (100 - 94) = 3 \times 6$	18
	Rest of Answer:	$97 - 6$ or $94 - 3$	91
	Answer:		9118

Now to multiply two numbers, one above and one below is a little bit more tricky:

Let $n_1 = (100 + a)$ which is the number above 100 and $n_2 = (100 - b)$ which is the number below 100. Then:

$$\begin{aligned} n_1 \cdot n_2 &= (100 + a) \cdot (100 - b) \\ &= 10000 + 100(a - b) + ab \\ &= 100(100 + a - b) - ab \\ &= 100(100 + a - b - 1) + (100 - ab) \\ &= 100(n_1 - b - 1) + (100 - ab) \end{aligned}$$

To see what this means, it is best to use an example:

$103 \times 94 =$	Tens/Ones:	$100 - 3 \times 6$	82
	Rest of Answer:	$103 - 6 - 1$	96
	Answer:		9682

So the trick is:

1. The Tens/Ones is just the difference the two numbers are from 100 multiplied together then subtracted from 100.
2. The rest of the answer is just the number that is larger than 100 *minus* the difference the smaller number is from 100 *minus* an additional 1

Let's look at another example to solidify this:

$$\begin{array}{rcll}
 & \text{Tens/Ones:} & 100 - 8 \times 7 & \mathbf{44} \\
 108 \times 93 = & \text{Rest of Answer:} & 108 - 7 - 1 & \mathbf{100} \\
 & \text{Answer:} & & \mathbf{10044}
 \end{array}$$

It should be noted that you can extend this trick to not just integers around 100 but 1000, 10000, and so forth. For the extension, you just need to keep track how many digits each part is. For example, when we are multiplying two numbers over 100 (say 104×103) the first two digits would be $4 \times 3 = 12$, however if we were doing two numbers over 1000 (like 1002×1007) the first *three* digits would be $2 \times 7 = \mathbf{014}$ not 14 like what you would be used to putting. Let's look at the example presented above and the procedure:

$$\begin{array}{rcll}
 & \text{Hundreds/Tens/Ones:} & 2 \times 7 & \mathbf{014} \\
 1002 \times 1007 = & \text{Rest of Answer:} & 1002 + 7 = 1007 + 2 & \mathbf{1009} \\
 & \text{Answer:} & & \mathbf{1009014}
 \end{array}$$

The best way to remember to include the "extra" digit is to think that when you multiply 1002×1007 you are going to *expect* a seven digit number. Now adding $1002 + 7 = 1009$ gives you four of the digits, so you need the first part to produce three digits for you.

Let's look at an example of two numbers below 1000:

$$\begin{array}{rcll}
 & \text{Hundreds/Tens/Ones:} & 7 \times 6 & \mathbf{042} \\
 993 \times 994 = & \text{Rest of Answer:} & 993 - 6 = 994 - 7 & \mathbf{987} \\
 & \text{Answer:} & & \mathbf{987042}
 \end{array}$$

The following are some practice problems so that you can fully understand this trick:

Problem Set 1.2.7

- | | |
|---|------------------------|
| 1. $89 \times 97 =$ | 8. $109 \times 107 =$ |
| 2. $96 \times 97 =$ | 9. $96 \times 89 =$ |
| 3. $103 \times 109 =$ | 10. $92 \times 97 =$ |
| 4. $93 \times 97 =$ | 11. $103 \times 104 =$ |
| 5. $103 \times 107 =$ | 12. $102 \times 103 =$ |
| 6. $93 \times 89 =$ | 13. $92 \times 93 =$ |
| 7. $\mathbf{102} \times \mathbf{108} =$ | 14. $106 \times 107 =$ |

- | | |
|--------------------------|------------------------------------|
| 15. $97 \times 89 =$ | 24. $97 \times 107 =$ |
| 16. $94 \times 98 =$ | 25. $93 \times 104 =$ |
| 17. $94 \times 91 =$ | 26. $96 \times 103 =$ |
| 18. $91 \times 98 =$ | 27. $991 \times 991 =$ |
| 19. $993 \times 994 =$ | 28. $104 \times 97 =$ |
| 20. $103 \times 96 =$ | 29. $1003 \times 1008 =$ |
| 21. $93 \times 103 =$ | 30. $(*) 98^2 + 97^2 =$ |
| 22. $991 \times 989 =$ | 31. $19^2 \times 3^2 \times 2^2 =$ |
| 23. $1009 \times 1004 =$ | |

1.2.8 Squares Ending in 5 Trick

Here is the derivation for this trick. Let $a5$ represent any number ending in 5 (a could be any positive integer, not just restricted to a one-digit number).

$$\begin{aligned}
 (a5)^2 &= (10a + 5)^2 \\
 &= 100a^2 + 100a + 25 \\
 &= 100a(a + 1) + 25
 \end{aligned}$$

So you can tell from this that any number ending in 5 squared will have its last two digits equal to 25 and the remainder of the digits can be found from taking the leading digit(s) and multiplying it by one greater than itself. Here are a couple of examples:

	Tens/Ones:		25
$85^2 =$	Thousand/Hundreds:	$8 \times (8 + 1)$	72
	Answer:		7225

The next example shows how to compute 15^4 by applying the square ending in 5 trick twice, one time to get what 15^2 is then the other to get that result squared.

	Tens/Ones:	25		Tens/Ones:	25
$15^2 =$	Thousands/Hundreds:	$1 \times (1 + 1) = 2$	$225^2 =$	Rest of Answer:	$22 \times (23) = 11 \times 46 = 506$
	Answer:	225		Answer:	50625

In the above trick you *also* use the double/half trick *and* the 11's trick. This just shows that for some problems using multiple tricks might be necessary. Another point to make is that several other tricks use

the squares ending in 5 trick somewhere in the computation (see Section 1.2.10). So although this problem set for this section is rather small, this trick is crucial to applying several other tricks.

Problem Set 1.2.8

- | | |
|--|---|
| 1. 25% of 25 = | 6. 45% of 45 - 45 = |
| 2. .35 × 3.5 = | 7. (*) 12 ⁴ = |
| 3. 12 ² + 2 × 12 × 13 + 13 ² = | 8. 505 × 505 = |
| 4. (115) ² = | 9. A square has an area of 12.25 sq. cm. It's perimeter is: |
| 5. f(x) = 9x ² - 12x + 4, f(19) = | |

1.2.9 Squares from 41-59

There is a quick trick for easy computation for squares from 41 - 59. Let k be a 1-digit positive integer, then any of those squares can be expressed as $(50 \pm k)$:

$$\begin{aligned}(50 \pm k)^2 &= 2500 \pm 100 \cdot k + k^2 \\ &= 100(25 \pm k) + k^2\end{aligned}$$

What this means is that:

1. The tens/ones digits is just the difference the number is from 50 squared (k^2).
2. The remainder of the answer is taken by *adding* (if the number is greater than 50) or *subtracting* (if the number is less than 50) that difference from 25.
3. Note: You could extend this concept to squares outside the range of 41 - 59 as long as you keep up with the carry appropriately.

Let's illustrate with a couple of examples:

46 ² =	Tens/Ones:	(50 - 46) ² = 4 ²	16
	Rest of Answer:	25 - 4	21
	Answer:		2116
57 ² =	Tens/Ones:	(57 - 50) ² = 7 ²	49
	Rest of Answer:	25 + 7	32
	Answer:		3249
61 ² =	Tens/Ones:	(61 - 50) ² = 11 ²	121
	Rest of Answer:	25 + 11 + 1	37
	Answer:		3721

Problem Set 1.2.9

1. $58^2 =$

5. (*) $48 \times 49 \times 50 =$

2. $(510)^2 =$

6. $56^2 =$

3. $47 \times 47 =$

7. $59 \times 59 =$

4. $53^2 =$

8. $41^2 =$

1.2.10 Multiplying Two Numbers Equidistant from a Third Number

To illustrate this concept, let's look at an example of this type of problem: 83×87 . Notice that both 83 and 87 are 2 away from 85. So:

$$83 \times 87 = (85 - 2) \times (85 + 2)$$

Notice this is just the difference of two squares:

$$(85 - 2) \times (85 + 2) = 85^2 - 2^2 = 7225 - 4 = \mathbf{7221}$$

So the procedure is:

1. Find the middle number between the two numbers being multiplied and square it.
2. Subtract from that the difference between the middle number and one of two numbers squared.

For most of these types of problems, the center number will be a multiple of 5, making the computation of its square relatively simple (See: Square's Ending in 5 Trick). The following illustrates another example:

$$61 \times 69 = 65^2 - 4^2 = 4225 - 16 = \mathbf{4209}$$

Problem Set 1.2.10

1. $84 \times 86 =$

6. $88 \times 82 =$

2. $53 \times 57 =$

7. $36 \times 24 =$

3. $48 \times 52 =$

8. **$7.6 \times 8.4 =$**

4. $62 \times 58 =$

9. $5.3 \times 4.7 =$

5. $6.8 \times 7.2 =$

10. $51 \times 59 + 16 =$

$11. 96 \times 104 =$

$12. 81 \times 89 + 16 =$

$13. 34 \times 36 + 1 =$

$14. 73 \times 77 + 4 =$

$15. 62 \times 68 + 9 =$

$16. 32 \times 38 + 9 =$

$17. 18 \times 24 + 9 =$

$18. 61 \times 69 + 16 =$

$19. 43 \times 47 + 4 =$

$20. 88 \times 82 + 9 =$

$21. 57 \times 53 + 4 =$

$22. 38 \times 28 =$

$23. 41 \times 49 - 9 =$

$24. \mathbf{77} \times \mathbf{73} + \mathbf{4} =$

$25. 65 \times 75 - 33 =$

$26. 33 \times 27 + 9 =$

$27. 71 \times 79 + 16 =$

$28. 72 \times 78 + 9 =$

$29. 53 \times 57 + 4 =$

$30. 105 \times 95 =$

$31. 62 \times 68 - 16 =$

$32. 36 \times 26 =$

$33. 83 \times 87 - 21 =$

$34. 23 \times 27 + 4 =$

$35. 29 \times 37 =$

$36. \mathbf{21} - \mathbf{83} \times \mathbf{87} =$

$37. 112 \times 88 =$

$38. (*) 52 \times 48 + 49 \times 51 =$

$39. (*) 4.9^3 \times 3.3^3 =$

$40. (*) 72 \times 68 + 71 \times 69 =$

$41. (*) 42 \times 38 + 41 \times 39 =$

$42. (*) 4.8^3 \times 6.3^3 =$

$43. (*) 4000 + 322 \times 318 =$

$44. \mathbf{118} \times \mathbf{122} + \mathbf{4} =$

$45. (*) 5.1^3 \times 7.9^3 =$

$46. (*) \mathbf{34} \times \mathbf{36} \times \mathbf{34} \times \mathbf{36} =$

1.2.11 Multiplying Reverses

The following trick involves multiplying two two-digit numbers whose digits are reverse of each other.

$$\begin{aligned}ab \times ba &= (10a + b) \times (10b + a) \\ &= 100(a \cdot b) + 10(a^2 + b^2) + a \cdot b\end{aligned}$$

Here is what we know from the above result:

1. The Ones digit of the answer is just the two digits multiplied together.
2. The Tens digit of the answer is the sum of the squares of the digits.
3. The Hundreds digit of the answer is the two digits multiplied together.

Let's look at an example:

	Ones:	3×5	15
$53 \times 35 =$	Tens:	$3^2 + 5^2 + 1$	35
	Hundreds:	$3 \times 5 + 3$	18
	Answer:		1855

Here are some more problems to practice this trick:

Problem Set 1.2.11

- | | |
|---------------------|----------------------|
| 1. $43 \times 34 =$ | 7. $15 \times 51 =$ |
| 2. $23 \times 32 =$ | 8. $14 \times 41 =$ |
| 3. $31 \times 13 =$ | 9. $18 \times 81 =$ |
| 4. $21 \times 12 =$ | 10. $36 \times 63 =$ |
| 5. $27 \times 72 =$ | 11. $42 \times 24 =$ |
| 6. $61 \times 16 =$ | 12. $26 \times 62 =$ |

1.3 Standard Multiplication Tricks

1.3.1 Extending Foiling

You can extend the method of FOILING to quickly multiply two three-digit numbers in the form $cba \times dba$. The general objective is you treat the digits of ba as one number, so after foiling you would get:

	Ones/Tens:	$(ba)^2$
$cba \times dba =$	Hundreds/Thousands:	$(c + d) \times (ba)$
	Rest of Answer:	$c \times d$

Let's look at a problem to practice this extension:

$$\begin{array}{rll}
 412 \times 612 = & \text{Ones/Tens:} & (12)^2 & \mathbf{144} \\
 & \text{Hundreds/Thousands:} & (4 + 6) \times (12) + 1 & \mathbf{121} \\
 & \text{Rest of Answer:} & 4 \times 6 + 1 & \mathbf{25} \\
 & \text{Answer:} & & \mathbf{252144}
 \end{array}$$

By treating the last two digits as a single entity, you reduce the three-digit multiplication to essentially a two-digit multiplication problem. The last two digits need not be the same in the two numbers (usually I do see this as the case though) in order to apply this method, let's look at an example of this:

$$\begin{array}{rll}
 211 \times 808 = & \text{Ones/Tens:} & 08 \times 11 & \mathbf{88} \\
 & \text{Hundreds/Thousands:} & 08 \times 2 + 11 \times 8 & \mathbf{104} \\
 & \text{Rest of Answer:} & 2 \times 8 + 1 & \mathbf{17} \\
 & \text{Answer:} & & \mathbf{170488}
 \end{array}$$

The method works the best when the last two digits don't exceed 20 (after that the multiplication become cumbersome). Another good area where this approach is great for is squaring three-digit numbers:

$$\begin{array}{rll}
 606^2 = 606 \times 606 & \text{Ones/Tens:} & 06 \times 06 & \mathbf{36} \\
 & \text{Hundreds/Thousands:} & 06 \times 6 + 6 \times 06 = 2 \times 6 \times 6 & \mathbf{72} \\
 & \text{Rest of Answer:} & 6 \times 6 & \mathbf{36} \\
 & \text{Answer:} & & \mathbf{367236}
 \end{array}$$

In order to use this procedure for squaring, it would be beneficial to have squares of two-digit numbers memorized. Take for example this problem:

$$\begin{array}{rll}
 431^2 = 431 \times 431 & \text{Ones/Tens:} & 31 \times 31 & \mathbf{961} \\
 & \text{Hundreds/Thousands:} & 31 \times 4 + 4 \times 31 + 9 = 2 \times 4 \times 31 + 9 & \mathbf{257} \\
 & \text{Rest of Answer:} & 4 \times 4 + 2 & \mathbf{18} \\
 & \text{Answer:} & & \mathbf{185761}
 \end{array}$$

If you didn't have 31^2 memorized, you would have to calculate it in order to do the first step in the process (very time consuming). However, if you have it memorized you would not have to do the extra steps, thus saving time.

Here are some practice problems to help with understanding FOILING three-digit numbers.

Problem Set 1.3.1

1. $202^2 =$

2. $406 \times 406 =$

3. $503 \times 503 =$

4. $607^2 =$

5. $208^2 =$

6. $306^2 =$

7. $509 \times 509 =$

8. $804^2 =$

9. $704 \times 704 =$

10. $408^2 =$

27. $203 \times 123 =$

11. $602 \times 602 =$

28. $121 \times 411 =$

12. $303^2 =$

29. $412 \times 112 =$

13. $909^2 =$

30. $505 \times 404 =$

14. $402^2 =$

31. $311 \times 113 =$

15. $707^2 =$

32. $124 \times 121 =$

16. $301 \times 113 =$

33. $918^2 =$

17. $803 \times 803 =$

34. $124 \times 312 =$

18. $404^2 =$

35. $311 \times 122 =$

19. $512^2 =$

36. $524^2 =$

20. $122 \times 311 =$

37. $133 \times 311 =$

21. $612^2 =$

38. $141 \times 141 =$

22. $321 \times 302 =$

39. $511 \times 212 =$

23. $714^2 =$

40. $122 \times 212 =$

24. $234 \times 211 =$

41. $(12012)(12012) =$

25. $112 \times 211 =$

42. $667^2 =$

26. $214 \times 314 =$

1.3.2 Factoring of Numerical Problems

In many of the intermediate problems, there are several questions where factoring can make the problem a lot easier. Outlined in the next couple of tricks are times when factoring would be beneficial towards calculation. We'll start off with some standard problems:

$$\begin{aligned}
21^2 + 63^2 &= 21^2 + (3 \cdot 21)^2 \\
&= 21^2 \cdot (1 + 9) \\
&= \mathbf{4410}
\end{aligned}$$

This is a standard trick of factoring that is common in the middle section of the test. Another factoring procedure is as followed:

$$\begin{aligned}
48 \times 11 + 44 \times 12 &= 11 \cdot (48 + 4 \times 12) \\
&= 11 \cdot (96) \\
&= \mathbf{1056}
\end{aligned}$$

Factoring problems can be easily identified because, at first glance, they look like they require dense computation. For example, the above problem would require two, two-digit multiplication *and then* their addition. Whereas when you factor out the 11 you are left with a simple addition and a multiplication using the 11's trick.

Another thing is that factoring usually requires the knowledge of another trick. For instance, the first problem required the knowledge of a square (21^2) while the second example involved applying the 11's trick.

The following are examples when factoring would lessen the amount of computations:

Problem Set 1.3.2

- | | |
|------------------------------------|--|
| 1. $8^2 + 24^2 =$ | 11. $2005 \times 5 + 2005 =$ |
| 2. $27^2 + 9^2 =$ | 12. $27 \times 33 - 11 \times 81 =$ |
| 3. $15 \times 12 + 9 \times 30 =$ | 13. $21 \times 38 - 17 \times 21 =$ |
| 4. $28 \times 6 - 12 \times 14 =$ | 14. $40 \times 12 + 20 \times 24 =$ |
| 5. $33^2 + 11^2 =$ | 15. $51^2 + 51 \times 49 =$ |
| 6. $48 \times 22 - 22 \times 78 =$ | 16. $30 \times 11 + 22 \times 15 =$ |
| 7. $3 \cdot 9^2 + 1 \cdot 3^2 =$ | 17. $21^2 + 7^2 =$ |
| 8. $2004 + 2004 \times 4 =$ | 18. $2006 - 2006 \times 6 =$ |
| 9. $32 \times 16 + 16 \times 48 =$ | 19. $12 \times 16 + 8 \times 24 =$ |
| 10. $19^2 + 19 =$ | 20. $\mathbf{1 \cdot 2^2 + 3 \cdot 6^2} =$ |
| | 21. $14 \times 44 - 14 \times 30 =$ |

22. $60 \times 32 - 32 \times 28 =$

23. $45 \times 22 - 44 \times 15 =$

24. $(20 \times 44) - (18 \times 22) =$

25. $49^2 + 49 =$

26. $29^2 + 29 =$

27. $16 \times 66 - 16 \times 50 =$

28. $59^2 + 59 =$

29. $14 \times 38 - 14 \times 52 =$

30. $41 \times 17 - 17 \times 24 =$

31. $17 \times 34 - 51 \times 17 =$

32. $15 \times 36 + 12 \times 45 =$

33. $69^2 + 69 =$

34. $13 \times 77 + 91 \times 11 =$

35. $11^3 - 11^2 =$

36. $12 \times 90 + 72 \times 15 =$

37. $79^2 + 79 =$

38. $54 \times 11 + 99 \times 6 =$

39. $10 \cdot 11 + 11 \cdot 11 + 12 \cdot 11 =$

40. $119^2 + 119 =$

41. $39^2 + 39 =$

42. $18 \times 36 - 18 \times 54 =$

43. $22 \times 75 + 110 \times 15 =$

44. $99 \times 99 + 99 =$

45. $45 \times 16 - 24 \times 30 =$

46. $11^2 - 11^3 =$

47. $25 \times 77 + 25 \times 34 =$

48. $15 \times 18 + 9 \times 30 =$

49. $24 \times 13 + 24 \times 11 =$

50. $129 \times 129 + 129 =$

51. $13 \times 15 + 11 \times 65 =$

52. (*) $33 \times 31 + 31 \times 29 =$

53. $31 \times 44 + 44 \times 44 =$

54. $12^2 + 24^2 =$

55. (*) $73 \times 86 + 77 \times 84 =$

56. (*) $63 \times 119 + 121 \times 17 =$

57. $48 \times 11 + 44 \times 12 =$

58. $109^2 + 109 =$

59. (*) $38 \times 107 + 47 \times 93 =$

60. $64 \times 21 - 42 \times 16 =$

61. (*) $23 \times 34 + 43 \times 32 =$

62. $72 \times 11 + 99 \times 8 =$

63. (*) $43 \times 56 + 47 \times 54 =$

64. $15 \times 75 + 45 \times 25 =$

65. $42 \times 48 + 63 \times 42 =$

66. $14^2 - 28^2 =$

67. (*) $31 \times 117 + 30 \times 213 =$

68. $48 \times 28 + 27 \times 28 =$

69. $34 \times 56 + 55 \times 34 =$

70. (*) $34 \times 45 + 54 \times 43 =$

1.3.3 Sum of Consecutive Squares

Usually when approached with this problem, one of the squares ends in 5 making the squaring of the number relatively trivial. You want to use the approach of factoring to help aid in these problems. For example:

$$35^2 + 36^2 = 35^2 + (35 + 1)^2 = 2 \cdot 35^2 + 2 \cdot 35 + 1^2 = 2 \cdot 1225 + 70 + 1 = \mathbf{2521}$$

This is a brute force technique, however, it is a lot better than squaring both of the numbers then adding them together (which you can get lost very easily doing that).

Here are some more practice problems to familiarize yourself with this procedure.

Problem Set 1.3.3

1. $35^2 + 36^2 =$

4. $25^2 + 26^2 =$

2. $12^2 + 13^2 =$

5. $40^2 + 41^2 =$

3. $15^2 + 16^2 =$

6. $80^2 + 81^2 =$

1.3.4 Sum of Squares: Factoring Method

Usually on the 3rd or 4th column of the test you will have to compute something like: $(30^2 - 2^2) + (30 + 2)^2$ (where the subtracting and additions might be reversed). Instead of memorizing a whole bunch of formulas for each individual case, it is probably just best to view these as factoring problems and using the techniques of FOILING to aid you. So for our example:

$$(30^2 - 2^2) + (30 + 2)^2 = 2 \cdot 30^2 + 2 \cdot 30 \cdot 2 + 2^2 - 2^2 = 1800 + 120 = \mathbf{1920}$$

Usually the number needing to be squared is relatively simple (either ending in 0 or ending in 5), making the computations a lot easier. Other times, another required step of converting a number to something more manageable will be necessary. For example:

$$19^2 + (10^2 - 9^2) = (10 + 9)^2 + (10^2 - 9^2) = 2 \cdot 10^2 + 2 \cdot 10 \cdot 9 + 9^2 - 9^2 = 200 + 180 = \mathbf{380}$$

The following are some more problems to give you practice with this technique:

Problem Set 1.3.4

1. $(11 + 10)^2 + (11^2 - 10^2) =$
2. $(30 + 2)^2 + (30^2 - 2^2) =$
3. $(10 + 9)^2 + (10^2 - 9^2) =$
4. $(30 + 2)^2 - (30^2 - 2^2) =$
5. $24^2 - (20^2 + 4^2) =$
6. $31^2 - (29^2 - 2^2) =$
7. $(30^2 - 2^2) + (30 + 2)^2 =$
8. $81^2 + (80 + 1)(80 - 1) =$
9. $55^2 - (50^2 - 5^2) =$
10. $47^2 + 40^2 - 7^2 =$
11. $(55 + 3)^2 + 55^2 - 3^2 =$
12. $30^2 - (28^2 - 2^2) =$
13. $38^2 + (30 + 8)(30 - 8) =$
14. $42^2 + (40^2 - 2^2) =$
15. $32^2 - (30^2 - 2^2) =$
16. $(28 + 2)^2 + (28^2 - 2^2) =$
17. $22^2 + 20^2 - 2^2 =$
18. $45^2 - (40^2 - 5^2) =$
19. $55^2 - 50^2 + 5^2 =$
20. $(30 + 2)^2 - (30^2 - 2^2) =$
21. $53 \times 53 + 50 \times 50 - 3 \times 3 =$
22. $46^2 - (21^2 - 25^2) =$

1.3.5 Sum of Squares: Special Case

There is a special case of the sum of squares that have repeatedly been tested. In order to apply the trick, these conditions must be met:

1. Arrange the two numbers so that the unit's digit of the first number is one greater than the ten's digit of the second number.
2. Make sure the sum of the ten's digit of the first number and the one's digit of the second number add up to ten.
3. If the above conditions are met, the answer is the sum of the squares of the digits of the first number times 101.

Let's look at an example: $72^2 + 13^2$.

1. The unit's digit of the first number (2) is one greater than the ten's digit of the second number (1).
2. The sum of the ten's digit of the first number (7) and the unit's digit of the second number (3) is 10.

3. The answer will be $(7^2 + 2^2) \times 101 = \mathbf{5353}$.

It is important to arrange the numbers accordingly for this particular trick to work. For example, if you see a problem like: $34^2 + 64^2$, it looks like a difficult problem where this particular trick won't apply. However, if you switch the order of the two numbers you get $34^2 + 64^2 = 64^2 + 34^2 = (6^2 + 4^2) \times 101 = \mathbf{5252}$.

Generally this trick is on the third column, and it is relatively simple to notice when to apply it because if you were having to square the two numbers *and* add them together it would take a long time. That should tip you off immediately that there is a trick that you should apply!

The following are some practice problems with this trick:

Problem Set 1.3.5

1. $93^2 + 21^2 =$

5. $45^2 + 46^2 =$

2. $12^2 + 19^2 =$

6. $36^2 + 57^2 =$

3. $72^2 + 13^2 =$

7. $55^2 + 56^2 =$

4. $82^2 + 12^2 =$

8. $37^2 + 67^2 =$

1.3.6 Difference of Squares

Everybody should know that $x^2 - y^2 = (x - y)(x + y)$, and you can apply this trick when asked to find the difference between squares of numbers. For example:

$$54^2 - 55^2 = (54 - 55)(54 + 55) = \mathbf{-109}$$

This is a pretty basic trick and is easily recognizable on the test.

The following are some more practice to give you a better feel of the problems:

Problems Set 1.3.6

1. $73^2 - 72^2 =$

6. $54^2 - 55^2 =$

2. $36^2 - 34^2 =$

7. $67^2 - 65^2 =$

3. $57^2 - 58^2 =$

8. $88^2 - 87^2 =$

4. $67^2 - 66^2 =$

9. $48^2 - 49^2 =$

5. $69^2 - 67^2 =$

10. $97^2 - 96^2 =$

11. $77^2 - 76^2 =$

12. $54^2 - 53^2 =$

13. $42^2 - 44^2 =$

14. $4.7^2 - 3.3^2 =$

15. $1.3^2 - 2.6^2 =$

16. $65^2 - 64^2 + 63^2 - 62^2 =$

17. $24^2 - 6^2 =$

18. $56^2 - 55^2 + 54^2 - 53^2 =$

19. $76^2 - 74^2 =$

20. $3.5^2 - 6.5^2 =$

21. $22^2 - 23^2 + 24^2 - 25^2 =$

22. $55^2 - 50^2 =$

23. $83^2 - 82^2 + 81^2 - 80^2 =$

24. $55^2 - 52^2 =$

25. $44^2 - 43^2 + 42^2 - 41^2 =$

26. $111^2 - 110^2 + 109^2 - 108^2 =$

27. $11^2 - 22^2 =$

28. $77^2 - 76^2 + 75^2 - 74^2 =$

29. $63^2 - 57^2 =$

30. $56^2 - 55^2 + 54^2 - 53^2 =$

31. $59^2 - 71^2 =$

32. $16^2 - 17^2 + 18^2 - 19^2 =$

33. $41^2 - 42^2 + 43^2 - 44^2 =$

34. $18^2 - 6^2 =$

35. $x^2 + 16^2 = 19^2$, then $x^2 =$

36. $4.5^2 - 1.5^2 =$

37. $21^2 - 20^2 + 19^2 - 18^2 =$

38. $58^2 - 59^2 + 60^2 - 61^2 =$

39. $72^2 - 78^2 =$

40. $24^2 - 22^2 + 20^2 - 18^2 =$

41. $89^2 - 86^2 + 83^2 - 80^2 =$

42. $48^2 - 62^2 =$

43. $74^2 - 76^2 + 78^2 - 80^2 =$

44. $38^2 - 27^2 =$

45. $31^2 - 33^2 + 35^2 - 37^2 =$

46. $48^2 - 44^2 + 40^2 - 36^2 =$

47. $79^2 - 76^2 + 73^2 - 70^2 =$

1.3.7 Multiplying Two Numbers Ending in 5

This is helpful trick for multiplying two numbers ending in 5. Let's look at its derivation, let $n_1 = a5 = 10a + 5$ and $n_2 = b5 = 10b + 5$ then:

$$\begin{aligned} n_1 \times n_2 &= (10a + 5) \cdot (10b + 5) \\ &= 100(ab) + 50(a + b) + 25 \\ &= 100\left(ab + \frac{a + b}{2}\right) + 25 \end{aligned}$$

So what does this mean:

1. If $a + b$ is even then the last two digits are 25.
2. If $a + b$ is odd then the last two digits are 75.
3. The remainder of the answer is just $ab + \lfloor \frac{a + b}{2} \rfloor$, where $\lfloor x \rfloor$ is the greatest integer less than or equal to x .

Let's look at an example in each case:

$45 \times 85 =$	Ones/Tens:	Since $4 + 8$ is even	25
	Rest of Answer:	$4 \times 8 + \frac{4 + 8}{2} = 32 + 6$	38
	Answer:		3825

$35 \times 85 =$	Ones/Tens:	Since $3 + 8$ is odd	75
	Rest of Answer:	$3 \times 8 + \lfloor \frac{3 + 8}{2} \rfloor = 24 + 5$	29
	Answer:		2975

Problem Set 1.3.7

- | | |
|---------------------|---------------------|
| 1. $35 \times 45 =$ | 5. $65 \times 45 =$ |
| 2. $95 \times 45 =$ | 6. $35 \times 85 =$ |
| 3. $35 \times 65 =$ | 7. $65 \times 95 =$ |
| 4. $85 \times 55 =$ | 8. $55 \times 95 =$ |

1.3.8 Multiplying Mixed Numbers

There are two major tricks involving the multiplication of mixed numbers. The first of which isn't a trick at all, only employing the technique of FOILing. Let's illustrate with an example:

$$\begin{aligned} 8\frac{1}{8} \times 24\frac{1}{8} &= \left(8 + \frac{1}{8}\right) \times \left(24 + \frac{1}{8}\right) \\ &= 8 \cdot 24 + (8 + 24) \cdot \frac{1}{8} + \frac{1}{8} \cdot \frac{1}{8} \\ &= 196\frac{1}{64} \end{aligned}$$

For the most part both of the whole numbers in the mixed numbers are usually divisible by the fraction you are multiplying by (in our example both 8 and 24 are divisible by 8), which means you can just write down the fractional part of the answer immediately and then continue with the problem.

The other trick for mixed numbers occur when the sum of the fractional part is 1 and the two whole numbers are the same. For example:

$$\begin{aligned} 9\frac{1}{3} \times 9\frac{2}{3} &= (9 + \frac{1}{3}) \times (9 + \frac{1}{3}) \\ &= 9^2 + (9 \cdot 2 + 9) \cdot \frac{1}{3} + \frac{1}{3} \cdot \frac{2}{3} \\ &= 9^2 + 9 + \frac{2}{9} \\ &= 9(9 + 1) + \frac{2}{9} \\ &= \mathbf{90\frac{2}{9}} \end{aligned}$$

So the trick is:

1. The fractional part of the answer is just the two fractions multiplied together.
2. If the whole part in the problem is n then the whole part of the answer is just $n \cdot (n + 1)$

Here is another example problem to show the procedure:

Fractional Part:	$\frac{2}{5} \cdot \frac{3}{5}$	$\frac{6}{25}$
$7\frac{2}{5} \times 7\frac{3}{5} =$	Whole Part: $7 \cdot (7 + 1)$	56
Answer:		$56\frac{6}{25}$

Although these tricks are great (especially foiling the mixed numbers) sometimes FOILING is very complicated, so the best method is to convert the mixed numbers to improper fractions and see what cancels. For example, you *don't* want to FOIL these mixed numbers:

$$4\frac{7}{12} \times 2\frac{2}{5} = \frac{7}{12} \cdot \frac{2}{5} + 4 \cdot \frac{2}{5} + 2 \cdot \frac{7}{12} + 4 \cdot 2$$

The above is *really* difficult to compute. Instead convert the numbers to improper fractions:

$$4\frac{7}{12} \times 2\frac{2}{5} = \frac{55}{12} \times \frac{12}{5} = \mathbf{11}$$

Usually the best method is to see if you can FOIL the numbers relatively quickly, and if you notice a stumbling block try to convert to improper fractions, then multiply.

Here are more practice problems to help you with these tricks:

Problem Set 1.3.8

1. $4\frac{1}{4} \times 8\frac{1}{4} =$

2. $8\frac{2}{3} \times 8\frac{1}{3} =$

3. $3\frac{4}{5} \times 3\frac{1}{5} =$

4. $4\frac{2}{3} \times 6\frac{1}{4} =$

5. $12\frac{1}{4} \times 8\frac{1}{4} =$

6. $15\frac{1}{6} \times 9\frac{1}{6} =$

7. $6\frac{1}{6} \times 12\frac{1}{6} =$

8. $11\frac{1}{11} \times 22\frac{1}{11} =$

9. $25\frac{2}{5} \times 5\frac{2}{5} =$

10. $5.2 \times 10.2 =$

11. $8\frac{2}{3} \times 4\frac{2}{3} =$

12. $7\frac{1}{7} \times 14\frac{1}{7} =$

13. $5\frac{1}{5} \times 10\frac{1}{5} =$

14. $5\frac{1}{5} \times 25\frac{1}{5} =$

15. $\left(5\frac{2}{5}\right)^2 =$

16. $8\frac{1}{8} \times 16\frac{1}{8} =$

17. $10\frac{5}{6} \times 12\frac{4}{5} =$

18. $3\frac{1}{2} \times 5\frac{6}{7} =$

19. $11 \times 11\frac{10}{11} =$

20. $6\frac{2}{3} \times 9\frac{2}{3} =$

21. $\left(12\frac{2}{3}\right)^2 =$

22. $7\frac{1}{7} \times 49\frac{1}{7} =$

23. $3\frac{3}{4} \times 2\frac{2}{5} =$

24. $4.3 \times 2.1 =$

25. $6 \times 6\frac{5}{6} =$

26. $\left(6\frac{2}{3}\right)^2 =$

27. $15.2 \times 5.2 =$

28. $4\frac{3}{5} \times 4\frac{2}{3} =$

29. $3.125 \times 1.6 =$

30. $2.375 \times 2.4 =$

31. $2\frac{2}{5} \times 5\frac{2}{5} =$

1.3.9 $a \times \frac{a}{b}$ Trick

The following is when you are multiplying an integer times a fraction in the form $a \times \frac{a}{b}$: The derivation of the trick is not of importance, only the result is:

$$a \times \frac{a}{b} = [a + (a - b)] + \frac{(a - b)^2}{b}$$

Let's look at a couple of examples:

$$\begin{aligned}11 \times \frac{11}{13} &= 11 + (11 - 13) + \frac{(11 - 13)^2}{13} \\ &= 11 - 2 + \frac{4}{13} \\ &= \mathbf{9\frac{4}{13}}\end{aligned}$$

It also works for multiplying by fractions larger than 1:

$$\begin{aligned}13 \times \frac{13}{12} &= 13 + (13 - 12) + \frac{(13 - 12)^2}{12} \\ &= 13 + 1 + \frac{1}{12} \\ &= \mathbf{14\frac{1}{12}}\end{aligned}$$

As you can see, when you are multiplying by a fraction less than 1 you will be *subtracting* the difference between the numerator and denominator, while when you are multiplying by a fraction greater than 1 you will be *adding* the difference.

It should be noted that there are exceptions (usually on the fourth column) where applying this trick is relatively difficult, and it is much easier to just convert to improper fractions then subtract. An example of this (which was one of the last problems asked on a test) is:

$$7 \times \frac{7}{15} - 7 = (7 - 8) + \frac{8^2}{15} - 7 = -8 + \frac{64}{15} = -8 + 4 + \frac{4}{15} = -\mathbf{3\frac{11}{15}}$$

The above expression was relatively difficult to compute, however if we converted to improper fractions:

$$7 \times \frac{7}{15} - 7 = \frac{7 \cdot 7}{15} - \frac{7 \cdot 15}{15} = \frac{7 \cdot (7 - 15)}{15} = \frac{-56}{15} = -\mathbf{3\frac{11}{15}}$$

This method is a lot less cumbersome and gets the answer relatively swiftly. However, it should be noted that the majority of times the trick is applicable and should *definitely* be used.

The following are more problems to illustrate this trick:

Problem Set 1.3.9

1. $11 \times \frac{11}{14} =$

6. $29 \times \frac{29}{34} =$

2. $22 \times \frac{22}{25} =$

7. $31 \times \frac{31}{34} =$

3. $19 \times \frac{19}{23} =$

8. $14 \times \frac{14}{17} - 3 =$

4. $27 \times \frac{27}{32} =$

9. $11 \times \frac{11}{14} + 3 =$

5. $16 \times \frac{16}{19} =$

10. $\mathbf{13} \times \frac{\mathbf{13}}{\mathbf{16}} + \mathbf{13} =$

11. $13 \times \frac{13}{17} + 4 =$

12. $13 \times \frac{13}{14} - 13 =$

13. $17 \times \frac{17}{18} - 17 =$

14. $22 \times \frac{22}{25} - 22 =$

15. $14 \times \frac{14}{17} - 14 =$

16. $17 \times 1\frac{17}{21} =$

17. $13 \times \frac{13}{16} - 13 =$

18. $11 \times \frac{11}{12} - 11 =$

19. $7 \times \frac{7}{15} - 7 =$

20. $14 \times \frac{14}{17} - 14 =$

21. $15 \times \frac{15}{17} - 15 =$

22. $35 \times 1\frac{35}{38} =$

23. $13 \times \frac{13}{15} - 13 =$

1.3.10 Combination of Tricks

The following is a practice set of combinations of some of the multiplication tricks already mentioned in the book. Most are approximations which occur on the third or fourth columns of the test.

Problem Set 1.3.10

1. (*) $12 \times 14 \times 16 =$

2. (*) $21 \times 31 \times 41 =$

3. (*) $13 \times 15 \times 17 =$

4. (*) $14 \times 16 \times 28 =$

5. (*) $146 \times 5 \times 154 =$

6. (*) $24 \times 34 \times 44 =$

7. (*) $24 \times 36 \times 48 =$

8. (*) $44 \times 25 \times 11^2 =$

9. (*) $22 \times 25 \times 28 =$

10. (*) $83 \times 87 \times 91 =$

11. (*) $43 \times 47 \times 51 =$

12. (*) $27 \times 29 \times 31 \times 33 =$

13. (*) $23 \times 33 \times 43 =$

14. (*) $29 \times 127 + 31 \times 213 =$

15. (*) $41 \times 44 \times 47 =$

16. (*) $31 \times 42 \times 53 =$

17. (*) $22 \times 44 \times 66 =$

18. (*) $39 \times 40 \times 41 =$

19. (*) $\sqrt[3]{1329} \times \sqrt{171} \times 15 =$
20. (*) $42 \times 48 \times 45 =$
21. (*) $52 \times 55 \times 58 =$
22. (*) $18 \times 20 \times 22 =$
23. (*) $24 \times 34 \times 44 =$
24. (*) $80 \times 82 \times 84 =$
25. (*) $28 \times 30 \times 32 =$
26. (*) $66 \times 68 \times 70 =$
27. (*) $63 \times 65 \times 67 =$
28. (*) $41 \times 43 \div 51 \times 53 =$
29. (*) $67 \times 56 + 65 \times 76 =$
30. (*) $56 \times 45 + 54 \times 65 =$
31. (*) $112 \times 123 + 132 \times 121 =$
32. (*) $29 \times 11 + 31 \times 109 =$
33. (*) $75^2 \div 25^2 \times 50^4 =$
34. (*) $18^3 \times 15^3 \div 9^3 =$
35. (*) $50^5 \div 25^5 \times 5^5 =$
36. (*) $24^3 \times 21^3 \div 4^4 =$
37. (*) $21^3 \times 18^2 \div 9^3 =$
38. (*) $75^4 \div 50^3 \times 25^2 =$
39. $24^2 \times 18^3 \div 6^4 =$
40. (*) $\sqrt[3]{3380} \times \sqrt{223} \times 16 =$

1.4 Dividing Tricks

Most of these tricks concern themselves with finding the remainders when dividing by certain numbers.

1.4.1 Finding a Remainder when Dividing by 4,8, etc...

Everybody knows that to see if a number is divisible by 2 you just have to look at the last digit, and if that is divisible by 2 (i.e. any even number) then the entire number is divisible by 2. Similarly, you can extend this principle to see if any integer is divisible by 4, 8, 16, etc... For divisibility by 4 you look at the last two digits in the number, and if that is divisible by 4, then the entire number is divisible by 4. With 8 it is the last three digits, and so on. Let's look at some examples:

$123456 \div 4$ has what remainder?

$987654 \div 8$ has what remainder?

Look at last two digits: $56 \div 4 = r\mathbf{0}$

Look at last three digits: $654 \div 8 = r\mathbf{6}$

Here are some practice problems to get you familiar with this procedure:

Problem Set 1.4.1

1. $364 \div 4$ has what remainder:
2. **$1324354 \div 4$ has what remainder:**
3. $246531 \div 8$ has what remainder:
4. $81736259 \div 4$ has what remainder:
5. $124680 \div 8$ has what remainder:
6. $214365 \div 8$ has what remainder:
7. Find k so that the five digit number $5318k$ is divisible by 8:

1.4.2 Finding a Remainder when Dividing by 3,9, etc...

In order to find divisibility with 3, you can sum up all the digits and see if that result is divisible by 3. Similarly, you can do the same thing with 9. Let's look at two examples:

$34952 \div 3$ has what remainder?

$112321 \div 9$ has what remainder?

Sum of the Digits: $(3 + 4 + 9 + 5 + 2) = 23$

Sum of the Digits: $(1 + 1 + 2 + 3 + 2 + 1) = 10$

$23 \div 3 = r2$

$10 \div 9 = r1$

For some examples, you can employ faster methods by using modular techniques in order to get the results quicker (see Modular Arithmetic Section). For example, if we were trying to see the remainder of 366699995 when dividing by 3, rather than summing up all the digits (which would be a hassle) and then seeing the remainder when that is divided by 3, you can look at each digit and figure out what its remainder is when dividing by 3 then summing *those*. So for our example:

$366699995 \cong (0 + 0 + 0 + 0 + 0 + 0 + 0 + 0 + 2) \cong 2 \pmod{3}$ therefore it leaves a remainder of **2**.

Here is a set of practice problems:

Problem Set 1.4.2

1. $24680 \div 9$ has a remainder of:
2. $6253178 \div 9$ has a remainder of:
3. $2007 \div 9$ has a remainder of:
4. $13579 \div 9$ has a remainder of:
5. $2468 \div 9$ has a remainder of:
6. Find the largest integer k such that $3k7$ is divisible by 3:

1.4.3 Finding a Remainder when Dividing by 11

Finding the remainder when dividing by 11 is very similar to finding the remainder when dividing by 9 with one catch, you add up alternating digits (beginning with the ones digits) then subtract the sum of the remaining digits. Let's look at an example to illustrate the trick:

13542 \div 11 has what remainder?

Sum of the Alternating Digits: $(2 + 5 + 1) = 8$

Sum of the Remaining Digits: $(4 + 3) = 7$

Remainder: $8 - 7 = \mathbf{1}$

Sometimes adding then subtracting “down the digits” will be easier than finding two explicit sums then subtracting. For example, if we were finding the remainder of $3456789 \div 11$, instead of doing $(9 + 7 + 5 + 3) - (8 + 6 + 4) = 24 - 18 = \mathbf{6}$ it might be easier to do $9 - 8 + 7 - 6 + 5 - 4 + 3 = 1 + 1 + 1 + 3 = \mathbf{6}$. That is what is so great about number sense tricks; there are always methods and approaches to making them faster!

Problem Set 1.4.3

1. 7653 \div 11 has a remainder of:
2. 745321 \div 11 has a remainder of:
3. **142536 \div 11 has a remainder of:**
4. 6253718 \div 11 has a remainder of:
5. 87125643 \div 11 has a remainder of:
6. 325476 \div 11 has a remainder of:
7. Find k so that 23578k is divisible by 11:
8. Find k so that 1482065k5 is divisible by 11:
9. Find k so that 456k89 is divisible by 11:
10. Find k so that 377337k is divisible by 11:

1.4.4 Finding Remainders of Other Integers

A very popular question on recent number sense tests include finding the remainder when dividing by 6 or 12 or some combination using the tricks mentioned above. When dividing seems trivial, sometimes it is best to just long divide to get the remainder (for example $1225 \div 6 = r\mathbf{1}$ from obvious division), however, when this seems tedious, you can use a combination of the two of the tricks mentioned above (depending on the factors of the number you are dividing). Let’s look at an example:

556677 \div 6 has what remainder?

Dividing by 2: $r\mathbf{1}$

Dividing by 3: $(5 + 5 + 6 + 6 + 7 + 7) = 36 \div 3$ $r\mathbf{0}$

So now the task is to find an appropriate remainder (less than 6) such that it is odd (has a remainder of 1 when dividing by 2) and is divisible by 3 (has a remainder of 0 when dividing by 3). From this information, you get $r = \mathbf{3}$. Let’s look at another example to solidify this procedure:

$54259 \div 12$ has what remainder?

Dividing by 4: $59 \div 4$ $r\mathbf{3}$

Dividing by 3: $(5 + 4 + 2 + 5 + 9) = 25 \div 3$ $r\mathbf{1}$

So for this instance, we want an appropriate remainder (less than 12) that has a remainder of 3 when dividing by 4, and a remainder of 1 when dividing by 3. Running through the integers of interest (0 – 11), you get the answer $r = \mathbf{7}$.

The best way of getting faster with this trick is through practice and familiarization of the basic principles. The following are some more practice questions:

Problem Set 1.4.4

1. $2002 \div 6$ has a remainder of:
2. $2006 \div 6$ has a remainder of:
3. $112358 \div 6$ has a remainder of:
4. If $852k$ is divisible by 6 then the largest value for k is:
5. $13579248 \div 6$ has a remainder of:
6. **$322766211 \div 6$ has a remainder of:**
7. $563412 \div 6$ has a remainder of:
8. Find k so that the 4-digit number $567k$ is divisible by 6:
9. **If $86k6$ is divisible by 6 then the largest value for k is:**
10. $423156 \div 12$ has a remainder of:
11. If $555k$ is divisible by 6 then the largest value for k is:
12. **Find $k > 4$ so that the 6-digit number $3576k2$ is divisible by 12:**
13. $735246 \div 18$ has a remainder of:
14. $6253718 \div 12$ has a remainder of:
15. Find k so that the 5-digit number $8475k$ is divisible by 6:

1.4.5 Remainders of Expressions

Questions like $(4^3 - 15 \times 43) \div 6$ has what remainder, are very popular and appear anywhere from the 2^{nd} to the 4^{th} column. This problem has its root in modular arithmetic (See: Modular Arithmetic Section), and the procedure for solving it is simply knowing that “the remainders after algebra is equal to the algebra of the remainders.” So instead of actually finding what $4^3 - 15 \times 43$ is and then dividing by 6, we can figure out what the remainder of each term is when dividing by 6, then do the algebra. So:

$$(4^3 - 15 \times 43) \div 6 \cong (4 - 3 \times 1) \div 6 = r\mathbf{1}$$

It should be noted that if a negative value is computed as the remainder, addition of multiples of the number which you are dividing by are required. Let's look at an example:

$$(15 \times 43 - 34 \times 12) \div 7 \cong (1 \times 1 - 6 \times 5) \div 7 = -29 \Rightarrow -29 + 5 \cdot (7) = r\mathbf{6}$$

So in the above question, after computing the algebra of remainders, we get an unreasonable remainder of -29 . So to make this a reasonable remainder (a positive integer such that $0 \leq r < 7$), we added a multiple of 7 (in this case 35) to get the correct answer.

You can use this concept of "negative remainders" to your benefit as well. For example, if we were trying to see the remainder of $13^8 \div 14$, the long way of doing it would be noticing that $13^2 = 169 \div 14 = r1 \Rightarrow 1^4 \div 14 = r1$ or you could use this concept of negative remainders (or an example of congruencies if you are familiar with that term) to say that $13^8 \div 14 \Rightarrow (-1)^8 \div 14 = r1$.

The following are some practice problems to solidify using the "algebra of remainders" method:

Problem Set 1.4.5

- | | |
|--|--|
| 1. $(31 \times 6 - 17) \div 8$ has a remainder of: | 15. $(34 \times 5 - 6) \div 7$ has a remainder of: |
| 2. $(34 \times 27 + 13) \div 4$ has a remainder of: | 16. $(1 + 2 - 3 \times 4^5) \div 6$ has a remainder of: |
| 3. $(44 \times 34 - 24) \div 4$ has a remainder of: | 17. $(8^2 + 4 \times 6 - 10) \div 3$ has a remainder of: |
| 4. $(33 + 23 \times 13) \div 3$ has a remainder of: | 18. $(12 \times 5 + 18 + 15) \div 8$ has a remainder of: |
| 5. $(23 + 33 \times 43) \div 4$ has a remainder of: | 19. $(7^3 + 8^2 - 9^1) \div 6$ has a remainder of: |
| 6. $(24 \times 34 - 44) \div 7$ has a remainder of: | 20. $(20 + 4 \times 6^2) \div 8$ has a remainder of: |
| 7. $(11^2 + 9 \times 7) \div 5$ has a remainder of: | 21. $(72 \times 64 - 83) \div 7$ has a remainder of: |
| 8. $(15 \times 3 - 6^2) \div 9$ has a remainder of: | 22. $(15 \times 30 - 45) \div 7$ has a remainder of: |
| 9. $(12 \times 9 - 2^3) \div 8$ has a remainder of: | 23. $(6^4 \times 5^3 - 4^2) \div 3$ has a remainder of: |
| 10. $(65 \times 4 - 3^2) \div 10$ has a remainder of: | 24. $(2^4 \times 3^6 - 5^{10}) \div 4$ has a remainder of: |
| 11. $(34 \times 56 - 12) \div 9$ has a remainder of: | 25. $(9^2 - 7 \times 5) \div 4$ has a remainder of: |
| 12. $(65 - 4 \times 3) \div 6$ has a remainder of: | 26. $(8^2 \times 6 - 4) \div 3$ has a remainder of: |
| 13. $(2 \times 3^4 + 5^6) \div 7$ has a remainder of: | 27. $(12 \times 34 - 56) \div 7$ has a remainder of: |
| 14. $(23 - 4 \times 5 + 6) \div 7$ has a remainder of: | |

1.4.6 Dividing by 9 Trick

From a previous section it is explained how a remainder can be found when dividing by 9. However, you can continue this process of adding *select* digits to get the complete answer when dividing by 9. The following is the result when you divide a four digit number $abcd$ by 9 without carries. The details of the proof is omitted, only the result is shown:

$$\begin{array}{rcl}
 & \text{Fractional Part:} & \frac{a + b + c + d}{9} \\
 abcd \div 9 = & \text{Ones:} & a + b + c \\
 & \text{Tens:} & a + b \\
 & \text{Hundreds:} & a
 \end{array}$$

I think the gist of the trick is self explanatory, let's look at a simple example:

$$\begin{array}{rcl}
 & \text{Fractional Part:} & \frac{3 + 2 + 1 + 1}{9} & \mathbf{\frac{7}{9}} \\
 & \text{Ones:} & 3 + 2 + 1 & \mathbf{6} \\
 3211 \div 9 = & \text{Tens:} & 3 + 2 & \mathbf{5} \\
 & \text{Hundreds:} & 3 & \mathbf{3} \\
 & \text{Answer:} & & \mathbf{356\frac{7}{9}}
 \end{array}$$

Here is a little bit more complicated of a problem involving a larger number being divided as well as incorporating carries:

$$\begin{array}{rcl}
 & \text{Fractional Part:} & \frac{3 + 2 + 2 + 5 + 7}{9} & \mathbf{2\frac{1}{9}} \\
 & \text{Ones:} & 3 + 2 + 2 + 5 + 2 & \mathbf{14} \\
 32257 \div 9 = & \text{Tens:} & 3 + 2 + 2 + 1 & \mathbf{8} \\
 & \text{Hundreds:} & 3 + 2 & \mathbf{5} \\
 & \text{Thousands:} & 3 & \mathbf{3} \\
 & \text{Answer:} & & \mathbf{3584\frac{1}{9}}
 \end{array}$$

Here are some problems to give you more practice with this trick:

Problem Set 1.4.6

1. $354 \div 9 =$

5. $456 \div 9 =$

2. $503 \div 9 =$

6. $1234 \div 9 =$

3. $2003 \div 9 =$

7. $12345 \div 9 =$

4. $321 \div 9 =$

8. $2475 \div 45 =$

1.4.7 Converting $\frac{a}{40}$ and $\frac{b}{80}$, etc... to Decimals

The following isn't necessarily a trick but more of a procedure I like to follow when I am approached with converting $\frac{a}{40}$ and $\frac{b}{80}$ into decimals (usually on the first column of problems). So for $\frac{a}{40}$ I treat it as:

$$\frac{a}{40} = \frac{a}{40} \times \frac{\frac{1}{4}}{\frac{1}{4}} = \frac{a}{10}$$

So the technique is to divide the numerator by 4 then shift the decimal point over. Similarly, for $\frac{b}{80}$ you want to divide by 8 and shift the decimal point over. Let's look at a couple of examples:

$$\frac{43}{40} = 1 + \frac{3}{40} = 1 + \frac{.75}{10} = \mathbf{1.075}$$

$$\frac{27}{80} \Rightarrow \frac{27}{8} = 3.375 \Rightarrow \frac{3.375}{10} = \mathbf{.3375}$$

Here are some practice problems of this type:

Problem Set 1.4.7

1. $\frac{1}{40} = \text{-----}\%$

7. $.0125 = \text{-----}\%$ (frac.)

2. $\frac{3}{40} = \text{-----}\%$

8. 48 is -----% greater than 40.

3. $\frac{7}{40} = \text{-----}\%$

9. $\frac{7}{40} = \text{-----}\%$

4. $\frac{21}{40} = \text{-----}\%$

10. 32 is what % of 80.-----

5. $\frac{43}{40} = \text{-----}$ (dec.)

11. $\frac{11}{40} = \text{-----}\%$

6. $\frac{3}{(2^3)(5^1)} = \text{-----}$ (dec.)

12. $\frac{3^2}{(2^3)(5^2)} = \text{-----}$ (dec.)

13. 72 is what % of 400.

17. 27.5% = (frac.)

14. $\frac{5}{(2^3)(5^2)} = \dots\dots\dots$ (dec.)

18. $\frac{4^3}{(2^3)(5^2)} = \dots\dots\dots$ (dec.)

15. $4\frac{7}{20} = \dots\dots\dots\%$

19. 1.6 is% of 20.

16. $\frac{5}{80} = \dots\dots\dots\%$

20. $\frac{3^4}{(2^4)(5^4)} = \dots\dots\dots$ (dec.)

1.5 Adding and Subtracting

The following are tricks where adding/subtracting are required to solve the problems.

1.5.1 Subtracting Reverses

A common first column problem involves subtracting two numbers whose digits are reverses of each other (like $715 - 517$ or $6002 - 2006$). Let the first number $n_1 = abc = 100a + 10b + c$ so the second number with the digits reversed would be $n_2 = cba = 100c + 10b + a$ so:

$$\begin{aligned} n_1 - n_2 &= (100a + 10b + c) - (100c + 10b + a) \\ &= 100(a - c) + (c - a) \\ &= 100(a - c) - (a - c) \end{aligned}$$

So the gist of the trick is:

1. Take the difference between the most significant digit and the least significant digit and multiply it by 100 if it is a three-digit number, or if it is a four digit number multiply by 1000 (however, it only works for 4-digit numbers and above if the middle digits are 0's, for example $7002 - 2007$ the method works but $7012 - 2107$ it *doesn't* work).
2. Then subtract from that result the difference between the digits.

Let's look at an example:

812 - 218 =	Step 1:	$(8 - 2) \times 100$	600
	Step 2:	$600 - 6$	594
	Answer:		594

It also works for when the subtraction is a negative number, but you need to be careful:

105 - 501 =	Step 1:	$(1 - 5) \times 100$	-400
	Step 2:	$-400 - (1 - 5)$	-396
	Answer:		-396

Like I said, you have to be careful with negative signs, a better (and highly recommended approach outlined in the next section) is to say: $105 - 501 = -(501 - 105) = -396$. By negating and reversing the numbers, you deal with positive numbers which are naturally more manageable. After you find the solution, you negate the result because of the sign switch.

Problem Set 1.5.1

1. $654 - 456 =$

6. $5002 - 2005 =$

2. $256 - 652 =$

7. $2006 - 6002 =$

3. $4002 - 2004 =$

8. $2003 - 3002 =$

4. $702 - 207 =$

9. $678 - 876 =$

5. $453 - 354 =$

10. $2007 - 7002 =$

1.5.2 Switching Numbers and Negating on Subtraction

Far too common, students make a mistake when subtracting two fractions whose result is a negative answer. An example of this is $4\frac{5}{6} - 10\frac{11}{12}$. Most of the time, it is incredibly easier to switch the order of the subtraction then negating the answer. Taking the above problem as an example:

$$\begin{aligned} 4\frac{5}{6} - 10\frac{11}{12} &= -(10\frac{11}{12} - 4\frac{5}{6}) \\ &= -(10\frac{11}{12} - 4\frac{10}{12}) \\ &= -(6\frac{1}{12}) \end{aligned}$$

Here is another example to illustrate the same point:

$$\begin{aligned} 2\frac{5}{6} - 4\frac{2}{3} &= -(4\frac{2}{3} - 2\frac{5}{6}) \\ &= -(4\frac{4}{6} - 2\frac{5}{6}) \\ &= -(1\frac{5}{6}) \end{aligned}$$

Problems Set 1.5.2

1. $2\frac{2}{3} - 3\frac{5}{6} =$

6. $2\frac{3}{8} - 3\frac{1}{4} =$

2. $4\frac{2}{3} - 6\frac{3}{5} =$

7. $2\frac{3}{4} - 6\frac{7}{8} =$

3. $1\frac{5}{9} - 3\frac{5}{9} =$

8. $3\frac{4}{5} - 8\frac{9}{10} =$

4. $2\frac{3}{4} - 4\frac{3}{5} =$

9. $3\frac{4}{9} - 5\frac{1}{3} =$

5. $1\frac{3}{7} - 3 =$

10. $5\frac{6}{7} - 12\frac{13}{14} =$

$$11. 3\frac{1}{6} - 6\frac{1}{3} =$$

$$14. 4\frac{5}{6} - 10\frac{11}{12} =$$

$$12. 2\frac{5}{6} - 4\frac{2}{3} =$$

$$15. 2\frac{3}{5} - 7\frac{1}{10} =$$

$$13. 4\frac{7}{8} - 12\frac{23}{24} =$$

$$16. 1\frac{4}{5} - 3\frac{2}{5} =$$

$$\mathbf{1.5.3} \quad \frac{a}{b \cdot (b+1)} + \frac{a}{(b+1) \cdot (b+2)} + \cdots$$

The best way to illustrate this trick is by example:

$$\begin{aligned} \frac{1}{6} + \frac{1}{12} + \frac{1}{20} + \frac{1}{30} &= \frac{1}{2 \cdot 3} + \frac{1}{3 \cdot 4} + \frac{1}{4 \cdot 5} + \frac{1}{5 \cdot 6} \\ &= \frac{1+1+1+1}{2 \cdot 6} \\ &= \frac{4}{12} = \frac{1}{3} \end{aligned}$$

So the strategy when you see a series in the form of $\frac{a}{b \cdot (b+1)} + \frac{a}{(b+1) \cdot (b+2)} + \cdots$ is to add up all the numerators and then divide it by the smallest factor in the denominators multiplied by the largest factor in the denominators. Let's look at another series:

$$\begin{aligned} \frac{1}{42} + \frac{1}{56} + \frac{1}{72} + \frac{1}{90} + \frac{1}{110} &= \frac{1}{6 \cdot 7} + \frac{1}{7 \cdot 8} + \frac{1}{8 \cdot 9} + \frac{1}{9 \cdot 10} + \frac{1}{10 \cdot 11} \\ &= \frac{1+1+1+1+1}{6 \cdot 11} \\ &= \frac{5}{66} \end{aligned}$$

Problems Set 1.5.3

$$1. \frac{1}{12} + \frac{1}{20} + \frac{1}{30} + \frac{1}{42} =$$

$$3. \frac{1}{30} + \frac{1}{42} + \frac{1}{56} =$$

$$2. \frac{1}{72} + \frac{1}{90} + \frac{1}{110} + \frac{1}{132} =$$

$$4. \frac{7}{30} + \frac{7}{20} + \frac{7}{12} =$$

1.5.4 $\frac{a}{b} + \frac{b}{a}$ Trick

Let's look at when we add the two fractions $\frac{a}{b} + \frac{b}{a}$:

$$\begin{aligned} \frac{a}{b} + \frac{b}{a} &= \frac{a^2 + b^2}{ab} \\ &= \frac{2ab}{ab} - \frac{2ab}{ab} + \frac{a^2 + b^2}{ab} \\ &= 2 + \frac{(a-b)^2}{ab} \end{aligned}$$

Here is an example:

$$\frac{5}{7} + \frac{7}{5} = 2 + \frac{(7-5)^2}{7 \cdot 5} = \mathbf{2\frac{4}{35}}$$

There are some variations to this trick. For example:

$$\frac{11}{13} + \frac{2}{11} = \frac{11}{13} + \frac{13}{11} - \frac{11}{11} = 2 + \frac{2^2}{143} - 1 = \mathbf{1\frac{4}{143}}$$

This is a popular variation that is used especially on the last column of the test because the trick is there but not as obvious.

The following are some practice problems to help you master this trick:

Problems Set 1.5.4

1. $\frac{12}{13} + \frac{13}{12} =$

2. $\frac{5}{6} + \frac{6}{5} =$

3. $\frac{15}{19} + \frac{19}{15} =$

4. $\frac{3}{5} + \frac{5}{3} - 2 =$

5. $\frac{7}{5} + \frac{5}{7} - 1 =$

6. $\frac{11}{13} + \frac{2}{11} =$

7. $\frac{7}{13} + \frac{6}{7} =$

8. $\frac{5}{6} + 1\frac{1}{5} - 2 =$

9. $\frac{13}{15} + \frac{2}{13} =$

10. $\frac{5}{8} + \frac{8}{5} - \frac{9}{40} =$

11. $\frac{3}{5} + \frac{5}{3} + \frac{11}{15} =$

12. $\frac{5}{7} + \frac{7}{5} - 3 =$

13. $\frac{15}{17} + \frac{2}{15} =$

14. $\frac{11}{15} + \frac{4}{11} =$

15. $\frac{11}{13} + \frac{2}{11} =$

16. $\frac{14}{15} + \frac{1}{14} =$

17. $1\frac{12}{13} + 1\frac{1}{12} =$

18. $\left(\frac{5}{7} + \frac{7}{5}\right) \div 2 =$

19. $\frac{11}{12} + \frac{1}{11} =$

20. $\frac{15}{22} + \frac{7}{15} - 1 =$

21. $\frac{11}{14} + \frac{3}{11} - 2 =$

1.5.5 $\frac{a}{b} - \frac{na-1}{nb+1}$

The following deals with subtracting fractions in the form $\frac{a}{b} - \frac{na-1}{nb+1}$. Most of these problems are on the 3rd of 4th columns, and they are relatively easy to pick out because of how absurd the problem would be if you didn't know the formula:

$$\frac{a}{b} - \frac{na-1}{nb+1} = \frac{(a+b)}{b \cdot (nb+1)}$$

So the numerator of the answer is just the sum of the numerator and denominator of the first number (i.e. the number who's numerator and denominators are small values) while the denominator of the answer is just the multiplication of the two denominators. Here is an example:

$$\frac{6}{7} - \frac{29}{36} = \frac{6+7}{7 \cdot 36} = \frac{\mathbf{13}}{\mathbf{252}}$$

Like I said it is easy to notice when to do this problem because, if you didn't know the formula, it would be relatively difficult to solve swiftly.

There is one variation to the formula which is:

$$\frac{a}{b} - \frac{na+1}{nb-1} = \frac{-(a+b)}{b \cdot (nb-1)}$$

When approached with these problems, it is best to take stock of which type it is. The easiest way of noticing which formula to apply is observing whether the denominator of the more "complicated" number is one *greater* or one *less* than a multiple of the denominator of the "simple" number. Let's look at another example:

$$\frac{7}{11} - \frac{43}{65} = \frac{-(7+11)}{11 \cdot 65} = \frac{\mathbf{-18}}{\mathbf{715}}$$

So on the above question, notice that 65 is one less a multiple of 11, so you know to apply the second formula.

Here are some practice problems to help you out:

Problems Set 1.5.5

1. $\frac{4}{9} - \frac{11}{28} =$

7. $\frac{3}{8} - \frac{26}{73} =$

2. $\frac{2}{7} - \frac{7}{29} =$

8. $\frac{4}{5} - \frac{\mathbf{67}}{\mathbf{86}} =$

3. $\frac{4}{13} - \frac{11}{40} =$

9. $\frac{8}{3} - \frac{41}{14} =$

4. $\frac{7}{15} - \frac{27}{61} =$

10. $\frac{8}{9} - \frac{87}{100} =$

5. $\frac{8}{11} - \frac{31}{45} =$

11. $\frac{\mathbf{67}}{\mathbf{81}} - \frac{\mathbf{17}}{\mathbf{20}} =$

6. $\frac{8}{11} - \frac{\mathbf{87}}{\mathbf{122}} =$

12. $\frac{3}{8} - \frac{14}{41} =$

$$13. \frac{7}{15} - \frac{15}{29} =$$

$$14. \frac{5}{8} - \frac{24}{41} =$$

$$15. \frac{8}{9} - \frac{31}{37} =$$

$$16. \frac{10}{11} - \frac{39}{45} =$$

$$17. \frac{11}{16} - \frac{32}{49} =$$

$$18. \frac{8}{11} - \frac{87}{122} =$$

$$19. \frac{4}{7} - \frac{35}{64} =$$

$$20. \frac{9}{46} - \frac{2}{9} =$$

$$21. \frac{3}{8} - \frac{14}{41} =$$

$$22. \frac{7}{11} - \frac{55}{89} =$$

2 Memorizations

2.1 Important Numbers

2.1.1 Squares

In order for faster speed in taking the test, squares up to 25 should definitely be memorized (however, memorization of squares up to 50 would be highly recommended). In the event that memorization can't be achieved, remember the tricks discussed in Section 1 of the book as well as the method of foiling. The following table should aid in memorization:

$11^2 = 121$	$12^2 = 144$	$13^2 = 169$	$14^2 = 196$
$15^2 = 225$	$16^2 = 256$	$17^2 = 289$	$18^2 = 324$
$19^2 = 361$	$20^2 = 400$	$21^2 = 441$	$22^2 = 484$
$23^2 = 529$	$24^2 = 576$	$25^2 = 625$	$26^2 = 676$
$27^2 = 729$	$28^2 = 784$	$29^2 = 841$	$30^2 = 900$
$31^2 = 961$	$32^2 = 1024$	$33^2 = 1089$	$34^2 = 1156$
$35^2 = 1225$	$36^2 = 1296$	$37^2 = 1369$	$38^2 = 1444$
$39^2 = 1521$	$40^2 = 1600$	$41^2 = 1681$	$42^2 = 1764$
$43^2 = 1849$	$44^2 = 1936$	$45^2 = 2025$	$46^2 = 2116$
$47^2 = 2209$	$48^2 = 2304$	$49^2 = 2401$	$50^2 = 2500$

On the next page you will find practice problems concerning squares. Avoid FOILING when possible so that you can work on having automatic responses on most of the questions.

Problems Set 2.1.1

- $28^2 =$
- $3 \cdot 2^2 =$
- $29 \times 29 =$
- $16 \times 16 =$
- $31^2 =$
- If 2.2 cm=1 inch, then 2.2 in equals how many cm.:
- $34 \times 34 =$
- $17 \times 17 =$
- $23 \times 23 =$
- $19^2 =$
- $18 \times 18 =$
- 24% of 24 is:
- $23^2 =$
- $32^2 =$
- $22^2 =$
- $14 \times 14 =$
- $21^2 =$
- $24^2 =$
- 31% of 31 is:
- What is 27% of 27 :
- $28^2 =$
- $34^2 =$
- $26^2 =$
- $17^2 =$
- $33 \times 33 =$
- Find $x < 0$ when $x^2 = 729$:
- (*) $\sqrt{1090} \times 31 =$
- (*) $\sqrt{291} \times 23 =$
- $\sqrt{-196} \times \sqrt{-256} =$
- $\frac{3}{4}$ of 24% of 1.8 :
- (*) $509 \times \sqrt{905} =$
- (*) $\sqrt{327} \times \sqrt{397} \times \sqrt{487} =$
- (*) $14^4 =$
- (*) $\sqrt{362} \times \sqrt{440} =$
- $959 \times \sqrt{960} =$
- (*) $13^4 =$
- (*) $\sqrt{451} \times 451 =$
- (*) $\sqrt{574} \times \sqrt{577} \times \sqrt{580} =$
- (*) $17^4 =$

40. (*) $\sqrt{1025} \times \sqrt{63} =$

42. (*) $\sqrt{1030} \times 2^5 =$

41. (*) $28 \times 56 \times 14 \div 42 =$

43. (*) $21^4 =$

2.1.2 Cubes

The following cubes should be memorized:

$5^3 = 125$

$6^3 = 216$

$7^3 = 343$

$8^3 = 512$

$9^3 = 729$

$10^3 = 1000$

$11^3 = 1331$

$12^3 = 1728$

$13^3 = 2197$

$14^3 = 2744$

$15^3 = 3375$

$16^3 = 4096$

$17^3 = 4913$

$18^3 = 5832$

$19^3 = 6859$

$20^3 = 8000$

Again, only FOIL when necessary on the practice problems on the next page.

Problem Set 2.1.2

- $(1728)^{\frac{1}{3}} =$
- $11^3 =$
- $14 \times 14 \times 14 =$
- $(-343)^{\frac{1}{3}} =$
- $12^3 =$
- $16^3 =$
- $\sqrt[3]{1728} \div \sqrt{36} =$
- $11^4 \div 11 =$
- $(-12)^3 =$
- $(2197)^{\frac{1}{3}} =$
- $(-729)^{\frac{1}{3}} =$
- $8^3 =$
- $15^3 =$
- $12 \times 12 \times 12 =$
- $(125 \div 64)^{\frac{1}{3}} =$
- $13^3 =$
- $7 \times 7 \times 7 =$
- $-1331^{\frac{1}{3}} =$
- $6 \times 6 \times 6 =$
- $15 \times 15 \times 15 =$
- $(*) \sqrt[3]{1730} \times 145 =$
- $(27 \div 216)^{\frac{1}{3}} =$
- If $x = 7$ then $(x + 3)(x^2 - 3x + 9) =$
- $\sqrt{676} \div \sqrt[3]{-2197} =$
- $(1.728)^{\frac{1}{3}} =$
- $8^3 \times 5^3 =$
- $11^5 \div 121 =$
- $\sqrt[3]{1.331} =$
- $(*) 89 \times 90 \times 91 =$
- $\sqrt[3]{.729} =$
- $(*) (121)^3 =$
- $3^4 - 6^3 + 9^2 =$
- $\sqrt[3]{1728} \div \sqrt{576} =$
- $\sqrt{225} \times \sqrt[3]{3375} =$
- $8^3 - 9^3 =$
- $(*) 13^3 \times 3^4 =$
- $2^3 \times 5^3 \times 7^3 =$
- $(*) 119 \times 120 \times 121 =$
- $(*) 14^3 \times 4^5 =$
- $8^4 =$

2.1.3 Powers of 2, 3, 5

Memorizing powers of certain integers like 2, 3, 5, etc... can be beneficial in solving a variety of problems ranging from approximation problems to logarithm problems. In some instances, powers of integers can be calculated based on other means than memorization. For example, $7^4 = (7^2)^2 = 49^2 = 2401$ However, the following powers should be memorized for quick calculation:

$2^3 = 8$	$3^3 = 27$	$5^3 = 125$
$2^4 = 16$	$3^4 = 81$	$5^4 = 625$
$2^5 = 32$	$3^5 = 243$	$5^5 = 3125$
$2^6 = 64$	$3^6 = 729$	
$2^7 = 128$	$3^7 = 2187$	
$2^8 = 256$		
$2^9 = 512$		
$2^{10} = 1024$		

The following are problems concerning higher powers of certain integers.

Problem Set 2.1.3

1. $5^3 + 3^3 + 2^3 =$

2. $2^3 - 3^3 - 4^3 =$

3. $(\sqrt{64} - \sqrt{36})^5 =$

4. $5^x = 125, x^5 =$

5. $4^3 - 5^3 =$

6. $2^{x+1} = 32, x - 1 =$

7. $2^3 + 3^3 + 5^3 =$

8. $5^3 - 3^3 =$

9. $\sqrt[3]{125 \times 512} =$

10. $2^3 + 3^3 + 4^3 - 5^3 =$

11. $x^3 = 64$, so $3^x =$

12. $4^5 \times 5^5 =$

13. $27^2 =$

14. If $x^5 = -32$, then $5^x =$

15. $2^5 \times 5^3 =$

16. $8^4 \times 5^4 =$

17. (*) $5^5 + 4^4 + 3^3 + 2^2 + 1^1 =$

18. $2^6 \times 5^4 =$

19. $5^{x-1} = 3125, x + 1 =$

20. $2^3 - 3^3 - 5^3 =$

21. $\frac{3^4}{2^3 \cdot 5^3} =$

22. $6^3 + 4^3 + 2^3 =$

23. $3^4 + 4^3 = 5 \cdot x. x =$

24. (*) $5^1 + 4^2 + 3^3 + 2^4 + 1^5 =$

25. $9^x = 243, x =$

26. $8^3 \times 5^3 =$

27. $2^3 \times 8^3 \times 5^3 =$

28. $2^5 \times 3^4 \times 5^2 =$

29. $2^4 \times 7^2 \times 5^3 =$

30. $4^2 \times 5^2 \times 6^2 =$

31. $2^5 \times 3^3 \times 5^2 =$

32. $2^3 \times 3^4 \times 5^5 =$

33. $(3^3 - 2^3 + 1^3) \times 5^3 =$

34. $2^5 \times 3^4 \times 5^2 =$

35. $2^5 \times 3^4 \times 5^5 =$

36. $2^3 \times 3^2 \times 4^2 \times 5^3 =$

2.1.4 Important Fractions

The following fractions should be memorized for reasons stated in Section 1.2.5. In addition, early problems on the test involved converting these fractions to decimals and percentages, so if they were memorized, time would be saved. Omitted are the “obvious” fractions ($\frac{1}{4}, \frac{1}{3}, \frac{1}{5}, \text{etc.}$).

Fraction	%	Fraction	%	Fraction	%/Decimal
$\frac{1}{6}$	$16\frac{2}{3}\%$	$\frac{1}{7}$	$14\frac{2}{7}\%$	$\frac{1}{8}$	$12\frac{1}{2}\% = .125$
$\frac{5}{6}$	$83\frac{1}{3}\%$	$\frac{2}{7}$	$28\frac{4}{7}\%$	$\frac{3}{8}$	$37\frac{1}{2}\% = .375$
		$\frac{3}{7}$	$42\frac{6}{7}\%$	$\frac{5}{8}$	$62\frac{1}{2}\% = .625$
		$\frac{4}{7}$	$57\frac{1}{7}\%$	$\frac{7}{8}$	$87\frac{1}{2}\% = .875$
		$\frac{5}{7}$	$71\frac{3}{7}\%$		
		$\frac{6}{7}$	$85\frac{5}{7}\%$		

Fraction	%	Fraction	%	Fraction	%	Fraction	%
$\frac{1}{9}$	$11\frac{1}{9}\%$	$\frac{1}{11}$	$9\frac{1}{11}\%$	$\frac{1}{12}$	$8\frac{1}{3}\%$	$\frac{1}{16}$	$6\frac{1}{4}\%$
$\frac{2}{9}$	$22\frac{2}{9}\%$	$\frac{2}{11}$	$18\frac{2}{11}\%$	$\frac{5}{12}$	$41\frac{2}{3}\%$	$\frac{3}{16}$	$18\frac{3}{4}\%$
$\frac{3}{9}$	$33\frac{3}{9}\%$	$\frac{3}{11}$	$27\frac{3}{11}\%$	$\frac{7}{12}$	$58\frac{1}{3}\%$	$\frac{5}{16}$	$31\frac{1}{4}\%$
$\frac{4}{9}$	$44\frac{4}{9}\%$	$\frac{4}{11}$	$36\frac{4}{11}\%$	$\frac{11}{12}$	$91\frac{2}{3}\%$	$\frac{7}{16}$	$43\frac{3}{4}\%$
$\frac{5}{9}$	$55\frac{5}{9}\%$	$\frac{5}{11}$	$45\frac{5}{11}\%$			$\frac{9}{16}$	$56\frac{1}{4}\%$
$\frac{6}{9}$	$66\frac{6}{9}\%$	$\frac{6}{11}$	$54\frac{6}{11}\%$			$\frac{11}{16}$	$68\frac{3}{4}\%$
$\frac{7}{9}$	$77\frac{7}{9}\%$	$\frac{7}{11}$	$63\frac{7}{11}\%$			$\frac{13}{16}$	$81\frac{1}{4}\%$
$\frac{8}{9}$	$88\frac{8}{9}\%$	$\frac{8}{11}$	$72\frac{8}{11}\%$			$\frac{15}{16}$	$93\frac{3}{4}\%$
		$\frac{9}{11}$	$81\frac{9}{11}\%$				
		$\frac{10}{11}$	$90\frac{10}{11}\%$				

Fraction	%	Fraction	%
$\frac{1}{13}$	$7\frac{9}{13}\%$	$\frac{1}{14}$	$7\frac{1}{7}\%$
$\frac{2}{13}$	$15\frac{5}{13}\%$	$\frac{3}{14}$	$21\frac{3}{7}\%$
$\frac{3}{13}$	$23\frac{1}{13}\%$	$\frac{5}{14}$	$35\frac{5}{7}\%$
$\frac{4}{13}$	$30\frac{10}{13}\%$	$\frac{9}{14}$	$64\frac{2}{7}\%$
$\frac{5}{13}$	$38\frac{6}{13}\%$	$\frac{11}{14}$	$78\frac{4}{7}\%$
$\frac{6}{13}$	$46\frac{2}{13}\%$	$\frac{13}{14}$	$92\frac{6}{7}\%$
$\frac{7}{13}$	$53\frac{11}{13}\%$		
$\frac{8}{13}$	$61\frac{7}{13}\%$		
$\frac{9}{13}$	$69\frac{3}{13}\%$		
$\frac{10}{13}$	$76\frac{12}{13}\%$		
$\frac{11}{13}$	$84\frac{8}{13}\%$		
$\frac{12}{13}$	$92\frac{4}{13}\%$		

To aid in memorization, it would first help to memorize the first fractions in each column. From, here the others can be quickly derived by multiplying the initial fraction by the required integer to get the desired results. For example, if you only had $\frac{1}{11}$ memorized as $9\frac{1}{11}\%$, but you need to know what $\frac{5}{11}$ is, then you could simply multiply by 5:

$$5 \times \frac{1}{11} = 5 \times \left(9\frac{1}{11}\%\right) = 45\frac{5}{11}\%$$

Although memorization of *all* fractions is ideal, this method will result in correctly answering the question, albeit a lot slower. On the next page is a set of practice problems concerning fractions.

Problem Set 2.1.4

1. $12\frac{1}{2}\%$ = _____ (frac.)
2. $\frac{11}{5}$ = _____%
3. Which is larger $\frac{5}{9}$ or .56 :
4. **Which is larger $\frac{5}{8}$ or .622 :**
5. $\frac{17}{8}$ = _____ (dec.)
6. $.777\dots - .333\dots + .555\dots =$
7. $\frac{3}{5}$ = _____%
8. $\frac{1}{8}$ = _____ (dec.)
9. **Which is smaller $\frac{9}{11}$ or .81 :**
10. $\frac{1}{16}$ = _____%
11. $.125 - .375 - .625 =$
12. $\frac{11}{4}$ = _____%
13. Which is larger $\frac{5}{9}$ or .555 or 55% :
14. $.1666\dots - .333\dots + .8333\dots =$
15. The reciprocal of -1.0625 is:
16. Which is larger .46 or $\frac{5}{11}$:
17. $.111\dots - .333\dots - .666\dots =$
18. 37.5% = _____ (frac.)
19. Which is smaller $\frac{9}{11}$ or .8 :
20. $\frac{3}{7}$ = _____%
21. $\frac{7}{9}$ = _____%
22. $.08333\dots + .1666\dots + .25 =$
23. Which is smaller $\frac{7}{11}$ or .56 :
24. **Which is larger $\frac{9}{11}$ or 81% :**
25. $.1666\dots + .333\dots + .8333\dots =$
26. $\frac{7}{16}$ = _____% (dec.)
27. $32 \div .181818\dots =$
28. $\frac{2}{7}$ = _____%
29. Which is larger $-.375$ or $\frac{-5}{12}$:
30. **$.333\dots - .666\dots - .999\dots =$**
31. $\frac{1}{14}$ = _____%
32. $.0625 + .125 + .25 =$
33. $55\frac{5}{9}\%$ of 27 is:
34. 12.5% of 24 is:
35. Which is larger $-.27$ or $\frac{-2}{7}$:
36. $55 \div .454545\dots =$
37. $.111\dots - .1666\dots - .333\dots =$

38. $\frac{5}{16} = \text{-----}\%$ (dec.)

54. $64\frac{2}{7}\% = \text{-----}$ (frac.)

39. $363 \div .272727 \dots =$

55. $1.21 \div .09090 \dots =$

40. $21\frac{3}{7}\% = \text{-----}$ (frac.)

56. $1\frac{7}{8} = \text{-----}\%$ (frac.)

41. $88 \times .090909 \dots =$

57. $6.25\% = \text{-----}$ (frac.)

42. $4\frac{4}{5} \div .444 \dots =$

58. $\frac{17}{14} = \text{-----}\%$

43. $\frac{3}{14} = \text{-----}\%$

59. $42\frac{6}{7}\% = \text{-----}$ (frac.)

44. $35\frac{5}{7}\% = \text{-----}$ (frac.)

60. $3\frac{3}{4}\% = \text{-----}$ (frac.)

45. $72 \times .083333 \dots =$

61. $1\frac{1}{10}\% = \text{-----}$ (frac.)

46. $78\frac{4}{7}\% = \text{-----}$ (frac.)

62. $92\frac{6}{7}\% = \text{-----}$ (frac.)

47. $911 \div .090909 \dots =$

63. $7\frac{1}{7}\% = \text{-----}$ (frac.)

48. $\frac{1}{12} = \text{-----}\%$

64. 75 is 3.125% of:

49. $\frac{11}{14} = \text{-----}\%$

65. $6\frac{7}{8}\% = \text{-----}$ (dec.)

50. 50 is 6.25% of:

66. $\frac{13}{14} = \text{-----}\%$

51. $242 \div .181818 \dots =$

67. $3\frac{1}{13}\% = \text{-----}$ (frac.)

52. $16\frac{2}{3}\% \times 482 =$

68. $\frac{15}{14} = \text{-----}\%$

53. $75 \div .5555 \dots =$

69. $21\frac{3}{7}\% = \text{-----}$ (frac.)

2.1.5 Special Integers

The following integers have important properties which are exploited regularly on the number sense test. They are:

$$\begin{array}{llllll}
 \mathbf{999} & 999 = 27 \times 37 & \mathbf{77} & 77 = \frac{1001}{13} & \mathbf{3367} & 3367 = \frac{10101}{3} & \mathbf{1430} & 1430 = \frac{10010}{7} \\
 \mathbf{1073} & 1073 = 29 \times 37 & \mathbf{154} & 154 = \frac{2002}{13} & \mathbf{1443} & 1443 = \frac{10101}{7} & & \\
 & & \vdots & & & & & \\
 & & \mathbf{693} & 693 = \frac{9009}{13} & & & &
 \end{array}$$

The following are some examples showing how to use these special numbers:

999 Trick:

$$\begin{aligned}
 333 \times \frac{1}{27} \times \frac{1}{37} &= \frac{1}{3} \times \mathbf{999} \times \frac{1}{27} \times \frac{1}{37} \\
 &= \frac{1}{3} \times \frac{27 \cdot 37}{27 \cdot 37} \\
 &= \frac{1}{3}
 \end{aligned}$$

1001 Trick:

$$\begin{aligned}
 385 \times 13 &= 77 \times 5 \times 13 \\
 &= \frac{\mathbf{1001}}{13} \times 5 \times 13 \\
 &= 1001 \times 5 \\
 &= \mathbf{5005}
 \end{aligned}$$

10101 Trick:

$$\begin{aligned}
 1443 \times 56 &= \frac{\mathbf{10101}}{7} \times 56 \\
 &= 10101 \times \frac{56}{7} \\
 &= 10101 \times 8 \\
 &= \mathbf{80808}
 \end{aligned}$$

Problem Set 2.1.5

1. $572 \times 21 =$

2. $\frac{2}{37} \times 999 =$

3. $33.67 \times 15 =$

4. $715 \times 35 =$

5. $3367 \times 21 =$

6. $1073 \div 29 =$

7. $715 \times 28 =$

8. $429 \times 35 =$

9. $63 \times 429 =$

10. $1073 \div 37 =$

11. $444 \times \frac{5}{37} =$

12. $63 \times 572 =$

13. $143 \times 49 = 1001 \times \text{-----}$

14. $29 \times 37 =$

15. $42 \times 715 =$

16. $715 \times 98 =$

17. $27 \times 37 =$

18. $715 \times 77 =$

19. **$105 \times 715 =$**

20. **$429 \times 357 =$**

21. $14 \times 715 =$

22. $42 \times 429 =$

23. $21 \times 336.7 =$

24. $36 \times 3.367 =$

25. $715 \times 49 =$

26. $33.67 \times 27 =$

27. $707 \times 715 =$

28. $429 \times 21 =$

29. **$336.7 \times 3.3 =$**

30. **$707 \times 429 =$**

31. $385 \times 13 =$

32. $111 \times \frac{7}{27} =$

33. $539 \times 13 =$

34. $666 \times \frac{2}{37} =$

35. (*) $\frac{5}{37} \times 5548 =$

36. $333 \times \frac{1}{27} \times \frac{1}{37} =$

37. $462 \times 13 =$

38. $999 \times \frac{7}{27} \times \frac{7}{37} =$

39. $6006 \div 462 =$

40. $444 \times \frac{4}{37} =$

55. $888 \times \frac{4}{37} =$

41. $770 \times 13 =$

56. $666 \times \frac{1}{27} =$

42. $888 \times \frac{4}{37} =$

57. $777 \times \frac{7}{37} =$

43. $666 \times \frac{16}{27} \times \frac{24}{37} =$

58. $444 \times \frac{2}{27} =$

44. $143 \times 77 =$

59. $999 \times \frac{3}{37} =$

45. **$143 \times 63 =$**

60. $666 \times \frac{3}{27} =$

46. **$888 \times \frac{16}{27} \times \frac{24}{37} =$**

61. $888 \times \frac{24}{27} =$

47. $84 \times 429 =$

62. $999 \times \frac{1}{27} =$

48. $143 \times 49 =$

63. **$143 \times 13 \times 7 =$**

49. $444 \times \frac{5}{37} =$

64. **$666 \times \frac{18}{37} =$**

50. $222 \times \frac{1}{27} =$

65. **$999 \times \frac{5}{27} =$**

51. $63 \times 143 =$

66. $1001 \times 25 = 143 \times \text{-----}$

52. $555 \times \frac{6}{37} =$

67. $3 \times 11 \times 13 \times 21 =$

53. $444 \times \frac{1}{27} =$

68. $3 \times 5 \times 7 \times 11 \times 13 =$

54. $143 \times 77 =$

2.1.6 Roman Numerals

The following are the roman numerals commonly tested on the exam:

$$I = 1 \quad V = 5 \quad X = 10 \quad L = 50$$

$$C = 100 \quad D = 500 \quad M = 1000$$

Knowing the above table and also the fact that you arrange the numerals in order from greatest to least ($M \rightarrow I$) with the exception of one rule: you can't put four of the same numerals consecutively. For example,

to express 42 in roman numerals it would *not* be 42 =XXXXII, it would be 42 =XLII. To circumvent the problem of putting four of the same numerals consecutively, you use a method of “subtraction.” Anytime a numeral of lesser value is placed in front of a numeral of greater value, you subtract from the larger numeral the small numeral. So in our case 40 is represented by XL=50 – 10 = 40. When converting numbers, it best to think of the number as a sum of ones, tens, hundreds, etc... units). A good example of what I mean is to express 199 in roman numerals. The way you want to look at it is 199 = 100 + 90 + 9 then express each one as a roman numeral. So 100 = C, 90 = XC, and 9 = IX, so 199 = **CXCIX**.

Problem Set 2.1.6

- | | |
|---------------------------------|-------------------------------|
| 1. MMXLII= | 18. MCXI+DLV= |
| 2. XLIV= | 19. MMV–DCXLI= |
| 3. MMIII= | 20. MMLIX–LIII= |
| 4. CXCIX= | 21. MCXI–DLV= |
| 5. MDCLXVI= | 22. CMIX–CDIV= |
| 6. CDXLIV= | 23. MDXLV–XV= |
| 7. CCLXXVII= | 24. DCII÷IX= |
| 8. MCDLIX = | 25. CCCLXXIV÷XI= |
| 9. CMXCIX= | 26. CDI×V= |
| 10. MMCCXXII= | 27. CCLXXX÷XIV= |
| 11. CXI–CC= | 28. MMV÷V= |
| 12. MD+DC= | 29. XXVII×CXI= |
| 13. CM+XC+IX= | 30. MI × XI = |
| 14. DC–LX–VI= | 31. MMVII×XXV= |
| 15. XIII + MMIV = | 32. MCCLX÷XV= |
| 16. MIII+MIV= | 33. MMVI×XI= |
| 17. MC+DL+XIV= | 34. CDIV ÷ XL = |

2.1.7 Platonic Solids

The following is a list of important characteristics of Platonic Solids which are popularly asked on the test:

Platonic Solid	Face Polygons	# of Faces	# of Vertices	# of Edges
Tetrahedron	Triangles	4	4	6
Cube	Squares	6	8	12
Octahedron	Triangles	8	6	12
Dodecahedron	Pentagons	12	20	30
Icosahedron	Triangles	20	12	30

If you ever forget one of the characteristics of the solids but remember the other two, you can always use Euler's formula of: $Faces + Vertices - Edges = 2$ to get the missing value.

The following is, albeit abridged, problem set concerning Platonic Solids. For best practice, cover up the above to table!

Problem Set 2.1.7

1. A dodecahedron has _____ vertices.
2. An icosahedron has _____ congruent faces.
3. The area of the base of a tetrahedron is 4 ft^2 .
The total surface area is _____ ft^2 .
4. A decahedron has _____ congruent regions.
5. A tetrahedron has _____ vertices.
6. An octahedron has _____ edges.
7. A hexahedron has _____ faces.
8. A dodecahedron is a platonic solid with 30 edges and _____ vertices.
9. An octahedron has _____ vertices.
10. An icosahedron is a platonic solid with 30 edges and _____ vertices.
11. **A dodecahedron is a platonic solid with 30 edges and _____ vertices.**

2.1.8 π and e Approximations

Using the standard approximations of: $\pi \approx 3.1$, $e \approx 2.7$, and $e^2 \approx 7.4$ lead to the beneficial results of:

$$\pi^2 \approx 10, e^3 \approx 20, \text{ and } \pi \cdot e \approx 8.5$$

Knowing these values, we can approximate various powers of e and π relatively simple and within the require margin of error of $\pm 5\%$. The following is an example where these approximations are useful:

$$\begin{aligned}
(e \times \pi)^4 &= e^4 \times \pi^4 \\
&= e \cdot e^3 \cdot (\pi^2)^2 \\
&\approx e \cdot 20 \cdot 100 \\
&\approx e \cdot 2000 \\
&\approx \mathbf{5400}
\end{aligned}$$

The following are more practice problems concerning these approximations:

Problem Set 2.1.8

- | | |
|--------------------------------------|---|
| 1. (*) $2\pi^4 =$ | 9. (*) $(e + 1.3)^5 =$ |
| 2. (*) $e^2 \times \pi^4 =$ | 10. (*) $[(\pi - .2)(+.3)]^3 =$ |
| 3. (*) $e^4 =$ | 11. (*) $(\pi + \mathbf{1.9})^3(\mathbf{e} + \mathbf{2.3})^3 =$ |
| 4. (*) $\pi^5 =$ | 12. (*) $(4e)^3 =$ |
| 5. (*) $(e \times \pi)^4 =$ | 13. (*) $\mathbf{e^4}\pi^4 =$ |
| 6. (*) $\pi^5 + e^4 =$ | 14. (*) $\pi^e e^\pi =$ |
| 7. (*) $\pi^3 \times \mathbf{e^4} =$ | 15. (*) $(3\pi + 2e)^4 =$ |
| 8. (*) $(3\pi)^4 =$ | 16. (*) $\pi^\pi \mathbf{e^e} =$ |

2.1.9 Distance Conversions

The following are important conversion factors for distances:

$$1 \text{ mile} = 5280 \text{ ft.}$$

$$1 \text{ mile} = 1760 \text{ yd.}$$

$$1 \text{ mile/hr} = \frac{22}{15} \text{ ft/s}$$

$$1 \text{ ft/min} = \frac{1}{5} \text{ in/s}$$

$$1 \text{ ft/hr} = \frac{1}{5} \text{ in/min}$$

$$1 \text{ mile/hr} = \frac{88}{5} \text{ in/s}$$

$$1 \text{ inch} = 2.54 \text{ cm.}$$

Problem Set 2.1.9

1. 15 miles per hour=_____ feet per second.
2. 3.5 yards=_____ inches.
3. .375 of a foot=_____ in.
4. 48 inches per second=_____ ft/min.
5. 7.5 mph=_____ inches per second.
6. $12\frac{1}{2}\%$ of a mile=_____ yards.
7. 25% of a mile=_____ yards.
8. $\frac{1}{3}$ of a mile = _____ feet.
9. $\frac{3}{4}$ of 3 yards=_____ inches.
10. $\frac{2}{3}$ of a mile=_____ ft.
11. 10 feet=_____ yards.
12. $83\frac{1}{3}\%$ of a foot=_____ inches.
13. 30 mph=_____ ft/sec.
14. 30 feet per minute=_____ feet per second.
15. 36 in/s =_____ inches per minute.
16. 480 inches per minute=_____ in/s.
17. 45 mph = _____ ft/s.
18. 33 ft/s=_____ mph.
19. **7.5 mph=_____ ft/s.**

2.1.10 Conversion between Distance → Area, Volume

Students find linear conversions relatively simple (for example 1ft.= 12in.), however when asked to find how many cubic inches are in cubic feet, they want to revert back to the linear conversion, which is incorrect ($1\text{ft.}^3 \neq 12\text{in.}^3$). When converting between linear distance to areas, and volumes you must square or cube the conversion factor appropriately. So in our example, we know that:

$$1\text{ft.} = 12\text{in.} \Rightarrow 1\text{ft.}^3 = (12)^3\text{in.}^3 = \mathbf{1728\text{in.}^3}$$

Another example converting ft.^2 to yd.^2 is:

$$1\text{yd.} = 3\text{ft.} \Rightarrow 1\text{yd.}^2 = (3)^2\text{ft.}^2 = \mathbf{9\text{ft.}^2}$$

Problem Set 2.1.10

1. 3 cubic yards=_____ ft.^3
2. 1 cubic foot=_____ cubic inches.
3. 9 square yards=_____ square feet.
4. 432 square inches=_____ ft.^2
5. 3 square yards=_____ square feet.
6. 243 cubic feet=_____ cubic yards.
7. 3 cubic feet=_____ cubic inches.
8. 4320 cubic inches=_____ cubic feet.

9. 1 square meter=_____ square centimeters.

10. 12 square feet=_____ square yards.

11. 216 square inches=_____ square feet.

12. 1728 cubic inches=_____ cubic feet.

13. $1\frac{1}{3}$ cubic yards=_____ cubic feet.

14. 2 cubic feet=_____ cubic inches.

15. 5 square decameters=_____square meters.

2.1.11 Fluid and Weight Conversions

The following are important fluid conversions. Although some conversions can be made from others (for example, the amount of cups in a gallon doesn't need to be explicitly stated, but it would be helpful to have it memorized so you don't have to multiply how many quarts in a gallon, how many pints in a quart, and how many cups in a pint), it is recommended that everything in the table should be memorized:

1 gallon = 4 quarts

1 quart = 2 pints

1 pint = 2 cups

1 gallon = 16 cups

1 gallon = 128 oz.

1 cup = 8 oz.

1 tbsp. = .5 oz.

1 tsp. = $\frac{1}{6}$ oz.

1 gallon = 231 in³

1 pound = 16 oz.

1 ton = 2000 lbs.

Problem Set 2.1.11

1. 1 quart=_____ cups.

2. 1 quart=_____ ounces.

3. 3 pints=_____ ounces.

4. 3 gallons=_____ cubic inches.

5. $\frac{2}{3}$ gallon=_____ cubic inches.

6. $1\frac{1}{3}$ gallon= _____ cubic inches.

7. 75% of 1 gallon=_____ ounces.

8. 256 ounces=_____ pounds.

9. **750 pounds=_____% of a ton.**

10. 75% of a gallon=_____ pints.

11. $12\frac{1}{2}$ % of a pint=_____ ounces.

12. 4 pints is what % of a gallon:

13. 2 quarts is what % of a pint:

14. 6 tablespoons is _____% of a cup.

15. 9 cups is what % of a quart:
16. A quart is what % of a cup:
17. 2541 cubic inches=_____ gallons.
18. 3 pints is what % of a cup:
19. 3 pints is what % of a gallon:
20. **5 gallons=_____cubic inches.**
21. 32 ounces=_____ pints.
22. 3.5 pints=_____ quarts.
23. 2.5 pints=_____ cups.
24. 37.5% of a gallon is _____ pints.
25. 62.5% of a gallon is _____ quarts.
26. **87.5% of a gallon is _____ ounces.**
27. 16 ounces is what part of a gallon:
28. 1 gallon=_____ cubic inches.
29. $\frac{3}{11}$ of a gallon=_____ cubic inches.
30. $\frac{3}{8}$ of a quart=_____ ounces.
31. $\frac{7}{11}$ of a gallon=_____ cubic inches.
32. 3 quarts and 2 pints=_____ ounces.
33. **7 quarts and 6 pints=_____ gallons.**

2.1.12 Celsius to Fahrenheit Conversions

These types of problems used to *always* be on the number sense tests in the early 1990's, but have since been noticeably absent until recently. Here are the conversion factors:

$$\text{Fahrenheit} \rightarrow \text{Celsius: } C = \frac{5}{9}(F - 32)$$

$$\text{Celsius} \rightarrow \text{Fahrenheit: } F = \frac{9}{5}C + 32$$

A couple of important degrees which pop-up frequently is that $32^{\circ}\text{F} = 0^{\circ}\text{C}$, $212^{\circ}\text{F} = 100^{\circ}\text{C}$, and $-40^{\circ}\text{F} = -40^{\circ}\text{C}$.

Do the following conversions:

Problem Set 2.1.12

1. $25^{\circ}\text{C} = \text{_____}^{\circ}\text{F}$
2. $-40^{\circ}\text{C} = \text{_____}^{\circ}\text{F}$
3. $98.6^{\circ}\text{F} = \text{_____}^{\circ}\text{C}$

2.2 Formulas

The following are handy formulas which, when mastered, will lead to solving a large handful of problems.

2.2.1 Sum of Series

The following are special series whose sums should be memorized:

Sum of the First m Integers

$$\sum_{n=1}^m n = 1 + 2 + 3 + \dots + m = \frac{m \cdot (m + 1)}{2}$$

Example:

$$1 + 2 + 3 + \dots + 11 = \frac{11 \cdot 12}{2} = \mathbf{66}$$

Sum of the First m Odd Integers

$$\sum_{n=1}^m 2n - 1 = 1 + 3 + 5 + \dots + (2m - 1) = \left(\frac{(2m - 1) + 1}{2} \right)^2 = m^2$$

Example:

$$1 + 3 + 5 + \dots + 15 = \left(\frac{15 + 1}{2} \right)^2 = 8^2 = \mathbf{64}$$

Sum of the First m Even Numbers

$$\sum_{n=1}^m 2n = 2 + 4 + 6 + \dots + 2m = m \cdot (m + 1)$$

Example:

$$2 + 4 + 6 + \dots + 22 = \frac{22}{2} \cdot \left(\frac{22}{2} + 1 \right) = 11 \cdot 12 = \mathbf{132}$$

Sum of First m Squares

$$\sum_{n=1}^m n^2 = 1^2 + 2^2 + \dots + m^2 = \frac{m \cdot (m + 1) \cdot (2m + 1)}{6}$$

Example:

$$1^2 + 2^2 + \dots + 10^2 = \frac{10 \cdot (10 + 1) \cdot (2 \cdot 10 + 1)}{6} = 35 \cdot 11 = \mathbf{385}$$

Sum of the First m Cubes

$$\sum_{n=1}^m n^3 = 1^3 + 2^3 + \dots + m^3 = \left(\frac{m \cdot (m + 1)}{2} \right)^2$$

Example:

$$1^3 + 2^3 + 3^3 + \dots + 10^3 = \left(\frac{10 \cdot 11}{2} \right)^2 = 55^2 = \mathbf{3025}$$

Sum of the First m Alternating Squares

$$\sum_{n=1}^m (-1)^{n+1} n^2 = 1^2 - 2^2 + 3^2 - \dots \pm m^2 = \pm \frac{m \cdot (m + 1)}{2}$$

Examples:

$$1^2 - 2^2 + 3^2 - \dots + 9^2 = \frac{9 \cdot 10}{2} = \mathbf{45}$$

$$1^2 - 2^2 + 3^2 - \dots - 12^2 = -\frac{12 \cdot 13}{2} = \mathbf{-78}$$

Sum of a General Arithmetic Series

$$\sum_{i=1}^m a_i = a_1 + a_2 + a_3 + \cdots + a_m = \frac{(a_1 + a_m) \cdot m}{2}$$

To find the number of terms: $m = \frac{a_m - a_1}{d} + 1$

Where d is the common difference.

Example:

$$8 + 11 + 14 + \cdots + 35 =$$

$$m = \frac{35 - 8}{3} + 1 = 10$$

$$\text{So } \sum = \frac{(8 + 35) \cdot 10}{2} = 43 \cdot 5 = \mathbf{215}$$

Sum of an Infinite Geometric Series

$$\sum_{n=0}^{\infty} a_1 \cdot (d)^n = a_1(1 + d + d^2 + \cdots) = \frac{a_1}{1 - d}$$

Where d is the common ratio with $|d| < 1$ and a_1 is the first term in the series.

Examples:

$$3 + 1 + \frac{1}{3} + \cdots = \frac{3}{1 - \frac{1}{3}} = \frac{3}{\frac{2}{3}} = \frac{\mathbf{9}}{\mathbf{2}}$$

$$4 - 2 + 1 - \frac{1}{2} + \cdots = \frac{4}{1 - \left(\frac{-1}{2}\right)} = \frac{4}{\frac{3}{2}} = \frac{\mathbf{8}}{\mathbf{3}}$$

Special Cases: Factoring

Sometimes simple factoring can lead to an easier calculation. The following are some examples:

$$\begin{aligned} 3 + 6 + 9 + \cdots + 33 &= 3 \cdot (1 + 2 + \cdots + 11) \\ &= 3 \left(\frac{11 \cdot 12}{2} \right) \\ &= 18 \cdot 11 = \mathbf{198} \end{aligned}$$

$$\begin{aligned} 11 + 33 + 55 + \cdots + 99 &= 11 \cdot (1 + 3 + 5 + \cdots + 9) \\ &= 11 \cdot \left(\frac{1 + 9}{2} \right)^2 \\ &= 11 \cdot 25 = \mathbf{275} \end{aligned}$$

Another important question involving sum of integers are word problems which state something similar to: The sum of three consecutive odd numbers is 129, what is the largest of the numbers?

In order to solve these problems it is best to know what you are adding. You can represent the sum of the three odd numbers by: $(n - 2) + n + (n + 2) = 129$. From this you can see that if you divide the number by 3, you will get that the *middle* integer is 43, thus making the largest integer $43 + 2 = \mathbf{45}$.

Here is another example problem: The sum of four consecutive even numbers is 140, what is the smallest?

For this one you can represent the sum by $(n - 2) + (n) + (n + 2) + (n + 4) = 140$, so dividing the number by 4 will get you the integer *between* the second and third even number. So $140 \div 4 = 35$, so the two middle integers are 34 and 36, making the smallest integer $\mathbf{32}$.

So from this we have learned that you can divide the sum by the number of consecutive integers you are adding, and if the number of terms are odd, you get the middle integer, and if the number of terms are

even, you get the number between the two middle integers.

The following are some more practice problems concerning the sum of series:

Problem Set 2.2.1

1. $2 + 4 + 6 + 8 + \cdots + 22 =$
2. $1 + 2 + 3 + 4 + \cdots + 21 =$
3. $1 + 3 + 5 + 7 + \cdots + 25 =$
4. The 25^{th} term of $3, 8, 13, 18, \dots$:
5. $6 + 4 + \frac{8}{3} + \frac{16}{9} + \cdots =$
6. $2 + 4 + 6 + 8 + \cdots + 30 =$
7. $1 + 3 + 5 + 7 + \cdots + 19 =$
8. $\frac{3}{5} - \frac{3}{10} + \frac{3}{20} - \cdots =$
9. The 20^{th} term of $1, 6, 11, 16, \dots$:
10. $22 + 20 + 18 + 16 + \cdots + 2 =$
11. $1 + 3 + 5 + \cdots + 17 =$
12. $2 + 4 + 6 + \cdots + 44 =$
13. $1 + \frac{1}{3} + \frac{1}{9} + \frac{1}{27} + \cdots =$
14. $1^3 + 2^3 + 3^3 + \cdots + 6^3 =$
15. $6 + 12 + 18 + \cdots + 66 =$
16. $3 + 5 + 7 + 9 + \cdots + 31 =$
17. $2 + 1 + \frac{1}{2} + \frac{1}{4} + \cdots =$
18. $-\frac{3}{2} + \frac{1}{2} - \frac{1}{6} + \frac{1}{18} - \cdots =$
19. $3 + 5 + 7 + 9 + \cdots + 23 =$
20. $\frac{4}{7} + \frac{8}{49} + \frac{16}{343} + \cdots =$
21. $1 + 4 + 7 + \cdots + 25 =$
22. $4 + 1 + \frac{1}{4} + \frac{1}{16} + \cdots =$
23. $2 + \frac{2}{5} + \frac{2}{25} + \cdots =$
24. $3 + 9 + 15 + 21 + \cdots + 33 =$
25. $7 + 14 + 21 + 28 + \cdots + 77 =$
26. The 11^{th} term in the arithmetic sequence $12, 9.5, 7, 4.5 \dots$ is:
27. $4 + 8 + 12 + \cdots + 44 =$
28. $8 + 16 + 24 + 32 + \cdots + 88 =$
29. $5^1 - 5^0 + 5^{-1} - 5^{-2} + \cdots =$
30. $(x) + (x+2) + (x+4) = 147$, then $(x) + (x+4) =$
31. $6 + 12 + 18 + 24 + \cdots + 36 =$
32. $3 + 8 + 13 + 18 + \cdots + 43 =$
33. $1^2 + 2^2 + 3^2 + 4^2 + 5^2 + 6^2 =$
34. $5 + 1 + \frac{1}{5} + \frac{1}{25} + \cdots =$

$$35. \frac{2}{3} + \frac{1}{2} + \frac{3}{8} + \frac{9}{32} + \dots =$$

$$36. 3 + 5 + 7 + 9 + \dots + 31 =$$

$$37. 7 + 14 + 21 + 28 + 35 + 42 =$$

$$38. 8 + 10 + 12 + \dots + 20 =$$

$$39. 10 + 15 + 20 + 25 + \dots + 105 =$$

$$40. 8 + 4 + 2 + 1 + \dots =$$

$$41. 4 + 8 + 12 + 16 + \dots + 44 =$$

$$42. (*) 1^3 + 2^3 + 3^3 + \dots + 6^3 =$$

$$43. \mathbf{6 + 12 + 18 + 24 + \dots + 66 =}$$

$$44. 2 + 6 + 10 + \dots + 42 =$$

$$45. 1^3 - 2^3 + 3^3 - 4^3 + 5^3 =$$

$$46. 3 + 1\frac{1}{2} + \frac{3}{4} + \dots =$$

$$47. 14 + 28 + 42 + 56 + 70 + 84 =$$

$$48. \mathbf{121 + 110 + 99 + \dots + 11 =}$$

$$49. 2 + 9 + 16 + 23 + \dots + 44 =$$

$$50. 13 + 26 + 39 + 52 + 65 + 78 =$$

$$51. \mathbf{36 + 32 + 28 + \dots + 12 =}$$

$$52. 88 + 80 + 72 + \dots + 8 =$$

53. The sum of 3 consecutive odd integers is 105.
The largest integer:

$$54. 4^1 - 4^0 + 4^{-1} - 4^{-2} + \dots =$$

$$55. (*) (1 + 2 + 3 + \dots + 29)^2 =$$

$$56. (*) 1^3 + 2^3 + 3^3 + \dots + 11^3 =$$

$$57. \frac{1}{5} + \frac{2}{5} + \frac{3}{5} + \dots + 1\frac{4}{5} + 2 =$$

$$58. (6^3 + 4^3 + 2^3) - (5^3 + 3^3 + 1^3) =$$

$$59. 3 - 1 - \frac{1}{3} - \frac{1}{9} - \frac{1}{27} - \dots =$$

$$60. \frac{1}{3} + \frac{2}{3} + 1 + 1\frac{1}{3} + \dots + 2\frac{1}{3} =$$

$$61. 3^3 - 4^3 - 2^3 + 5^3 =$$

$$62. \mathbf{6 - 1 - \frac{1}{6} - \frac{1}{36} - \dots =}$$

$$63. 2 + 5 + 8 + \dots + 20 =$$

$$64. (*) 1^3 + 2^3 + 3^3 + \dots + 13^3 =$$

$$65. \frac{3}{4} + \frac{9}{16} + \frac{27}{64} + \dots =$$

$$66. \frac{1}{4} + \frac{3}{4} + \frac{5}{4} + \dots + \frac{15}{4} =$$

$$67. (*) (3 + 6 + 9 + \dots + 30)^2 =$$

$$68. (*) 1^3 + 2^3 + 3^3 + \dots + 8^3 =$$

2.2.2 Fibonacci Numbers

It would be best to have the Fibonacci numbers memorized up to F_{15} because they crop up every now and then on the number sense test. In case you are unaware, the fibonacci sequence follows the recursive relationship of $F_{n-2} + F_{n-1} = F_n$. The following is a helpful table:

$$\begin{array}{cccc}
F_1 = 1 & F_2 = 1 & F_3 = 2 & F_4 = 3 \\
F_5 = 5 & F_6 = 8 & F_7 = 13 & F_8 = 21 \\
F_9 = 34 & F_{10} = 55 & F_{11} = 89 & F_{12} = 144 \\
F_{13} = 233 & F_{14} = 377 & F_{15} = 610 &
\end{array}$$

The most helpful formula to memorize concerning Fibonacci Numbers is the the sum of the first n Fibonacci Numbers is equal to $F_{n+2} - 1$.

A common problem asked on the latter parts of the number sense test is:

Find the sum of the first eight terms of the Fibonacci sequence 2, 5, 7, 12, 19, ...

Now there are two methods of approach for doing this. The first requires knowledge of large Fibonacci numbers:

Method 1:

The sum of the first n -terms of a general Fibonacci sequence $a, b, a + b, a + 2b, 2a + 3b, \dots$ is $\sum = a \cdot (F_{n+2} - 1) + d \cdot (F_{n+1} - 1)$. Where $d = (b - a)$.

So for our example:

$$\sum = 2 \cdot (F_{10} - 1) + (5 - 2) \cdot (F_9 - 1) = 2 \cdot 54 + 3 \cdot 33 = 108 + 99 = \mathbf{207}$$

Method 2:

The other method of doing this sum requires memorization of a formula for each particular sum. The following is a list of the sums of a general Fibonacci sequence $a, b, a + b, a + 2b, 2a + 3b, \dots$ for 1-12 terms (the number of terms which have been on the exam):

n	Fibonacci Number	Sum of First F_n Numbers	Formula
1	a	a	$a = F_1$
2	b	$a + b$	$a + b = F_3$
3	$a + b$	$2a + 2b$	$2(a + b) = 2 \cdot F_3$
4	$a + 2b$	$3a + 4b$	$4(a + b) - a = 4 \cdot F_3 - a$
5	$2a + 3b$	$5a + 7b$	$7(a + b) - 2a = 7 \cdot F_3 - 2a$
6	$3a + 5b$	$8a + 12b$	$4(2a + 3b) = 4 \cdot F_5$
7	$5a + 8b$	$13a + 20b$	$4(3a + 5b) + a = 4 \cdot F_6 + a$
8	$8a + 13b$	$21a + 33b$	$7(3a + 5b) - 2b = 7 \cdot F_6 - 2b$
9	$13a + 21b$	$34a + 54b$	$7(5a + 8b) - (a + 2b) = 7 \cdot F_7 - F_4$
10	$21a + 34b$	$55a + 88b$	$11(5a + 8b) = 11 \cdot F_7$
11	$34a + 55b$	$89a + 143b$	$11(8a + 13b) + a = 11 \cdot F_8 + a$
12	$55a + 89b$	$144a + 232b$	$18(8a + 13b) - b = 18 \cdot F_8 - b$

So in our case, we are summing the first 8 terms, which is just $7 \cdot F_6 - 2b$, where F_6 represents the sixth term in the sequence of 2, 5, 7, 12, 19, ... (which is 31), so $7 \cdot 31 - 2 \cdot 5 = 217 - 10 = \mathbf{207}$.

So in solving it this way you have to calculate the 6th term in the sequence as well as knowing the formula. Usually it will be required to calculate a middle term in the sequence, and then apply the formula.

These type of questions are usually computationally intense, so it is recommended to skip them and come back to work on them after the completion of all other problems. The following are some more practice problems:

Problem Set 2.2.2

- | | |
|---|---|
| 1. The sum of the first 11 terms of the Fibonacci Sequence
2, 4, 6, 10, 16, 26, ...: | 4, 5, 9, 14, 23, ...: |
| 2. The sum of the first 9 terms of the Fibonacci Sequence
3, 5, 8, 13, 21, ...: | 5. The sum of the first 11 terms of the Fibonacci Sequence
1, 5, 6, 11, 17, 28, ...: |
| 3. The sum of the first 9 terms of the Fibonacci Sequence
4, 7, 11, 18, 29, ...: | 6. The sum of the first 12 terms of the Fibonacci Sequence
1, 2, 3, 5, 8, 13, 21, ...: |
| 4. The sum of the first 10 terms of the Fibonacci Sequence | 7. The sum of the first 11 terms of the Fibonacci Sequence
2, 5, 7, 12, 19, 31, ...: |

- | | |
|--|---|
| <p>8. The sum of the first 9 terms of the Fibonacci Sequence
3, 8, 11, 19, ...:</p> | <p>13. The sum of the first 9 terms of the Fibonacci Sequence
1, 1, 2, 3, 5, ...:</p> |
| <p>9. The sum of the first 9 terms of the Fibonacci Sequence
2, 4, 6, 10, 16, ...:</p> | <p>14. The sum of the first 9 terms of the Fibonacci Sequence
-3, 2, -1, 1, 0, ...:</p> |
| <p>10. The sum of the first 9 terms of the Fibonacci Sequence
1, 5, 6, 11, 17, ...:</p> | <p>15. The sum of the first 9 terms of the Fibonacci Sequence
1, 3, 4, 7, 11, ...:</p> |
| <p>11. The sum of the first 9 terms of the Fibonacci Sequence
3, 5, 8, 13, 21, ...:</p> | <p>16. $1 + 1 + 2 + 3 + 5 + 8 + \dots + 55 =$</p> |
| <p>12. The sum of the first 9 terms of the Fibonacci Sequence
-3, 4, 1, 5, 6, ...:</p> | <p>17. $1 + 3 + 4 + 7 + 11 + 18 + \dots + 123 =$</p> |
| | <p>18. $3 + 6 + 9 + 15 + 24 + \dots + 267 =$</p> |
| | <p>19. $4 + 6 + 10 + 16 + 26 + \dots + 288 =$</p> |

2.2.3 Integral Divisors

The following are formulas dealing with integral divisors. On all the formulas, it is necessary to prime factorize the number of interest such that: $n = p_1^{e_1} \cdot p_2^{e_2} \cdot p_3^{e_3} \cdots p_n^{e_n}$.

Number of Prime Integral Divisors

Number of prime integral divisors can be found by simply prime factorizing the number, and count how many distinct prime numbers you have in it's representation.

Example:

Find the number of prime integral divisors of 120.

$$120 = 5 \cdot 2^3 \cdot 3 \Rightarrow \# \text{ of prime divisors} = (1 + 1 + 1) = \mathbf{3}$$

Number of Integral Divisors

$$\text{Number of Integral Divisors} = (e_1 + 1) \cdot (e_2 + 1) \cdot (e_3 + 1) \cdots (e_n + 1)$$

Example:

Find the number of integral divisors of 48.

$$48 = 2^4 \cdot 3 \Rightarrow (4 + 1) \cdot (1 + 1) = \mathbf{10}$$

Sum of the Integral Divisors

$$\Sigma = \frac{p_1^{e_1+1} - 1}{p_1 - 1} \cdot \frac{p_2^{e_2+1} - 1}{p_2 - 1} \cdots \frac{p_n^{e_n+1} - 1}{p_n - 1}$$

Example:

Find the sum of the integral divisors of 36.

$$36 = 2^2 \cdot 3^2$$

$$\sum = \frac{2^3 - 1}{2 - 1} \cdot \frac{3^3 - 1}{3 - 1} = \frac{7}{1} \cdot \frac{26}{2} = 7 \cdot 13 = \mathbf{91}$$

Number of Relatively Prime Integers less than N

$$\text{Number of Relatively Prime} = (p_1 - 1) \cdot (p_2 - 1) \cdots (p_n - 1) \cdot (p_1^{e_1 - 1}) \cdot (p_2^{e_2 - 1}) \cdots (p_n^{e_n - 1})$$

or

$$\text{Number of Relatively Prime} = \frac{p_1 - 1}{p_1} \cdot \frac{p_2 - 1}{p_2} \cdots \frac{p_n - 1}{p_n} \times n$$

Both techniques are relatively (no pun intended) quick and you should do whichever you feel comfortable with. Here is an example to display both method:

Example:

Find the number of relatively prime integers less than 20.

$$20 = 2^2 \cdot 5$$

$$\# \text{ of Relatively Prim Integers} = (2 - 1) \cdot (5 - 1) \cdot (2^{2-1}) \cdot (5^{1-1}) = 4 \cdot 2 = \mathbf{8}$$

or

$$\# \text{ of Relatively Prim Integers} = \frac{1}{2} \cdot \frac{4}{5} \times 20 = \mathbf{8}$$

Sum of Relatively Prime Integers less than N

$$\sum = (\# \text{ of Relatively Prime Integers}) \times \frac{n}{2}$$

Example:

Find the sum of the relatively prime integers less than 24.

$$24 = 2^3 \cdot 3$$

$$\# \text{ of Relatively Prim Integers} = \frac{1}{2} \cdot \frac{2}{3} \times 24 = 8$$

$$\sum = 8 \times \frac{24}{2} = 8 \cdot 12 = \mathbf{96}$$

We should introduce a distinction between proper and improper integral divisors here. A proper integral divisor is any positive integral divisor of the number excluding the number itself. So for example, the number 14 has 4 total integral divisors (1, 2, 7, 14), but only 3 proper integral divisors (1, 2, 7). Some number sense questions will ask for the sum of proper integral divisors or the number of proper integral divisors of number. When those are asked, you need to be aware to *exclude* the number itself from those calculations. For example, the sum of the proper integral divisors of $22 = 3 \times 12 - 22 = 36 - 22 = \mathbf{14}$.

In addition, on the questions asking for the number of co-prime (or relatively prime) within a range of values, it is best to calculate the total number of relatively prime integers and then start excluding ones that are out of range. For example, to calculate the number of integers greater than 3 which are co-prime to 20 you would find the number of co-prime integers less than 20 which is $(2 - 1)(5 - 1)(2^{2-1})(5^{1-1}) = 8$ then you can exclude the numbers 1 and 3. So the number of integers greater than three which are co-prime to 20 would be $8 - 2 = 6$. The quickest way of finding whether or not an integer is co-prime to another integer, is to put it in fraction form and see if the fraction is reducible. For example, 3 is co-prime to 20 because $\frac{3}{20}$ is irreducible.

With integral divisor problems it is best to get a lot of practice so that better efficiency can be reached. The following are some sample practice problems:

Problem Set 2.2.3

1. 30 has how many positive prime integral divisors:
2. 36 has how many positive integral divisors:
3. The sum of the prime factors of 42 is:
4. The number of prime factors of 210 is:
5. The number of positive integral divisors of 80 is:
6. The number of positive integral divisors of $2^4 \times 5$ is:
7. The sum of the distinct prime factors of 75 total:
8. The number of positive integral divisors of 96 is:
9. **The number of positive integral divisors of 100 is:**
10. The sum of the positive integral divisors 48 is:
11. The sum of the proper positive integral divisors of 24 is:
12. The sum of the positive integral divisors of 28 is:
13. The number of positive integral divisors of $6^1 \times 3^2 \times 2^3$:
14. The sum of the proper positive integral divisors of 30 is:
15. How many positive integral divisors does 81 have:
16. How many positive integral divisors does 144 have:
17. The sum of the positive integral divisors $3 \times 5 \times 7$ is:
18. The number of positive integral divisors of $6^5 \times 4^3 \times 2^1$:
19. The sum of the positive integral divisors of 20 is:
20. The number of positive integral divisors of 24 is:
21. The sum of the positive integral divisors of 28 is:
22. The number of positive integral divisors of $2^3 \times 3^4 \times 4^5$:
23. The number of positive integral divisors of 64 is:
24. **The sum of the proper positive integral divisors of 36 is:**
25. The number of positive integral divisors of $2^4 \times 3^6 \times 5^1$ is:
26. The number of positive integral divisors of $5^3 \times 3^2 \times 2^1$:
27. How many positive integers less than 90 are relatively prime to 90:
28. Sum of the proper positive integral divisors of 18 is:
29. The sum of the positive integers less than 18 that are relatively prime to 18:
30. The number of positive integral divisors of $12 \times 3^3 \times 2^4$:

- | | |
|--|--|
| <p>31. How many positive integers less than 16×25 are relatively prime to 16×25:</p> <p>32. How many integers between 30 and 3 are relatively prime to 30:</p> <p>33. How many positive integers divide 48:</p> <p>34. How many positive integer less than 9×8 are relatively prime to 9×8:</p> | <p>35. How many integers between 1 and 20 are relatively prime to 20:</p> <p>36. The number of positive integral divisors of $50 \times 5^4 \times 2^3$:</p> <p>37. How many positive integers divide 64:</p> <p>38. The sum of the positive integral divisors of 48:</p> |
|--|--|

2.2.4 Number of Diagonals of a Polygon

The formula for the number of diagonals in a polygon is derived by noticing that from each of the n vertices in an n -gon, you can draw $(n - 3)$ diagonals creating $n \cdot (n - 3)$ diagonals, however each diagonal would be drawn twice, so the total number of diagonals is:

$$\# \text{ of Diagonals} = \frac{n \cdot (n - 3)}{2}$$

As an example lets look at the number of diagonals in a hexagon:

$$\# \text{ of Diagonals in a Hexagon} = \frac{6 \cdot 3}{2} = 9$$

. Here are some problems for you to practice this formula:

Problem Set 2.2.4

- | | |
|---|--|
| <p>1. The number of diagonals a 5-sided regular polygon has:</p> <p>2. If a regular polygon has 27 distinct diagonals, then it has how many sides:</p> <p>3. A pentagon has how many diagonals:</p> <p>4. A nonagon has how many diagonals:</p> | <p>5. An octagon has how many diagonals:</p> <p>6. A decagon has how many diagonals:</p> <p>7. A rectangle has how many diagonals:</p> <p>8. A septagon has how many diagonals:</p> |
|---|--|

2.2.5 Exterior/Interior Angles

When finding the exterior, interior, or the sum of exterior or interior angles of a regular n -gon, you can use the following formulas:

Sum of Exterior Angles:	360°
Exterior Angle:	$\frac{360^\circ}{n}$
Interior Angle:	$180^\circ - \frac{360^\circ}{n} = \frac{180^\circ(n-2)}{n}$
Sum of Interior Angles:	$n \cdot \frac{180^\circ(n-2)}{n} = 180(n-2)$

If you were to only remember one of the above formulas, let it be that the sum of the exterior angles of every regular polygon be equal to 360° . From there you can derive the rest relatively swiftly (although it is *highly* recommended that you have all formulas memorized).

Example: Find the sum of the interior angles of an octagon.

Solution: $\sum = 180(8 - 2) = \mathbf{1080}$.

In order to find the interior angle from the exterior angle, you used the fact that they are supplements. Both supplements and complements of angles appear on the number sense test every now and then, so here are their definitions:

Complement of $\theta = 90^\circ - \theta$

Supplement of $\theta = 180^\circ - \theta$

Here are some practice problems on both exterior/interior angles as well as supplement/complement:

Problem Set 2.2.5

- | | |
|---|--|
| 1. A regular nonagon has an interior angle of: | 5. The measure of an interior angle of a regular hexagon measures: |
| 2. An interior angle of a regular pentagon has a measure of: | 6. The sum of the angles in a regular decagon is: |
| 3. The supplement of an interior angle of a regular octagon measures: | 7. The supplement of a 47° angle is: |
| 4. The angles in a regular octagon total: | 8. The sum of the interior angles of a regular pentagon is: |

2.2.6 Triangular, Pentagonal, etc... Numbers

We are all familiar with the concept of square numbers $1, 4, 9, 16, \dots, n^2$ and have a vague idea of how they can be viewed geometrically (n^2 can be represented by n rows of dots by n columns of dots). This same concept of translating “dots to numbers” can extend to any regular polygon. For example, the idea of a triangular number is the amount of dots which can be arranged into an equilateral triangle ($1, 3, 6, \dots$). The following are formulas for these “geometric” numbers:

$$\text{Triangular: } T_n = \frac{n(n+1)}{2}$$

$$\text{Square: } S_n = \frac{n(2n-0)}{2} \\ = n^2$$

$$\text{Pentagonal: } P_n = \frac{n(3n-1)}{2}$$

$$\text{Hexagonal: } H_n = \frac{n(4n-2)}{2}$$

$$\text{Heptagonal: } E_n = \frac{n(5n-3)}{2}$$

$$\text{Octagonal: } O_n = \frac{n(6n-4)}{2}$$

$$\text{M-Gonal: } M_n = \frac{n[(M-2)n - (M-4)]}{2}$$

As one can see, only the last formula is necessary for memorization (all the others can be derived from that one).

Some other useful formulas:

$$\text{Sum of Consecutive Triangular Numbers: } T_{n-1} + T_n = n^2$$

$$\text{Sum of First } m \text{ Triangular Numbers: } \sum_{n=1}^m T_n = T_1 + T_2 + \dots + T_m = \frac{m(m+1)(m+2)}{6}$$

$$\text{Sum of the Same Triangular and Pentagonal Numbers: } T_n + P_n = 2n^2$$

Examples:

1. The 6th Triangular Number?

$$\frac{6(6+1)}{2} = \mathbf{21}$$

2. The 4th Octagonal Number?

$$\frac{4(6 \cdot 4 - 4)}{2} = \frac{4 \cdot 20}{2} = \mathbf{40}$$

3. The 5th Pentagonal Number?

$$\frac{5(3 \cdot 5 - 1)}{2} = \frac{5 \cdot 14}{2} = \mathbf{35}$$

4. The Sum of the 6th and 7th Triangular Numbers?

$$7^2 = \mathbf{49}$$

Problem Set 2.2.6:

1. The 7th pentagonal number:

6. The 7th septagonal number is:

2. The 4th octagonal number:

7. **The 5th pentagonal number is:**

3. The 5th pentagonal number:

8. The 6th pentagonal number is:

4. The 8th octagonal number:

9. The 5th hexagonal number is:

5. **The 12th hexagonal number:**

10. The 11th triangular number is:

11. The 12th triangular number is: the 6th triangular numbers:
12. **The 6th hexagonal number is:** 14. The sum of the 3rd triangular and the 3rd pentagonal numbers:
13. The sum of the 5th triangular and

2.2.7 Finding Sides of a Triangle

A popular triangle question gives two sides of a triangle and asks for the minimum/maximum value for the other side conforming to the restriction that the triangle is right, acute or obtuse. The two sets of formulas which will aid in solving these questions are:

Triangle Inequality

$$a + b > c$$

Right Triangle: $a^2 + b^2 = c^2$

Variations on the Pythagorean Theorem Acute Triangle: $a^2 + b^2 > c^2$

Obtuse Triangle: $a^2 + b^2 < c^2$

If you don't have the Pythagorean relationships for acute/obtuse triangle memorized, the easiest way to think about the relationship on the fly is remembering that an equilateral triangle is acute so $a^2 + a^2 > a^2$.

Let's look at some examples:

An acute triangle has integer sides of 4, x , and 9. What is the largest value of x ?

Solution: Using the Pythagorean relationship we know: $4^2 + 9^2 > x^2$ or $97 > x^2$. Knowing this and the fact that x is an integer, we know that the largest value of x is **9**.

An acute triangle has integer sides of 4, x , and 9. What is the smallest value of x ?

Solution: For this we use the triangle inequality. We want 9 to be the largest side (so x would have to be less than 9), so apply the inequality knowing this: $4 + x > 9$ which leads to the smallest integer value of x is **6**

An obtuse triangle has integer sides of 7, x , and 8. What is the smallest value of x ?

Solution: For this, we want the largest value in the obtuse triangle to be 8 then apply the Triangle Inequality: $7 + x > 8$ with x being an integer. This makes the smallest value of x to be **2**.

An obtuse triangle has integer sides of 7, x , and 8. What is the largest value of x ?

Solution: Here, x is restricted by the Triangle Inequality (if we used the Pythagorean Theorem for obtuse, we would get an unbounded result for x : $7^2 + 8^2 < x^2$ makes x unbounded). So we know from that equation: $7 + 8 > x$ so the largest integer value for x is **14**.

Another important type of triangle problem involves being given one side of a right triangle and having to compute the other sides. For example, the sides of a right triangle are integers, one of its sides is 9, what is the hypotenuse?

Where this gets its foundation is from the Pythagorean Theorem which states that $a^2 + b^2 = c^2$. If the smallest side is given (call it a , then we can express $a^2 = c^2 - b^2 = (c - b)(c + b)$). Now is where the trick comes into play. The goal becomes to find two numbers that when subtracted together from each other multiplied with them added to each other is the smallest side squared. When the smallest side squared gives an odd number (in our case 81 is odd), the goal is reduced considerably by thinking of taking consecutive integers (so $c - b = 1$) and $c + b = a^2$. The easiest way to find two consecutive integers whose sum is a third number is to divide, the third number by 2, and the integers straddle that mixed number. So in our case $9^2 = 81 \div 2 = 40.5$ so $b = 40$ and $c = 41$, and we're done. Let's look at another example:

Example: The sides of a right triangle are integers, one of its sides is 11, what is the other side?

Solution: $11^2 = 121$ which is odd, so $121 \div 2 = 60.5$ so the other side is **60**.

Very seldom do they give you a side whose square is even. In that case let's look at the result:

Example: The sides of a right triangle are integers, one of its sides is 10, what is the hypotenuse?

Example: The easiest way of solving these problems is divide the number they give you by a certain amount to get an odd number, then perform the usual procedure on that odd number (outlined above), then when you get the results multiply each side by the number you originally divided by. Let's look at what happens in our example. So to get an odd number we must divide 10 by 2 to get 5. Now to find the other side/hypotenuse with smallest side given is 5 you do: $5^2 \div 2 = 12.5 \Rightarrow b = 12$ and $c = 13$. Now to get the correct side/hypotenuse lengths, we must multiply by what we originally divided by (2) so $b = 12 \cdot 2 = 24$ and $c = 13 \cdot 2 = 26$. As you can see there are a couple of more steps to this procedure, and you have to remember what you divided by at the beginning so you can multiply the side/hypotenuse by that same amount at the end.

There are some variations to this, say they tell you that the hypotenuse is 61 and ask for the smallest side. Since half of the smallest side squares is roughly the hypotenuse, you will be looking for squares who are near $61 \cdot 2 = 122$, so you know that $s = 11$.

In addition, there are some algebraic applications that frequently ask the same thing. For example, if it is given that $x^2 - y^2 = 53$ and asks you to solve for y . You do the same procedure: $(x + y)(x - y) = 53$, since 53 is odd, you are concerned with consecutive numbers adding up to 53, so $53 \div 2 = 26.5 \Rightarrow x = 27$ and $y = 26$.

Getting practice with these problems are critical so that you can immediately know which formula to apply and which procedure to follow. Complete the following:

Problem Set 2.2.7

- | | |
|---|--|
| 1. An obtuse triangle has integral sides of 3, x , and 7. The largest value for x is: | 3. x, y are positive integers with $x^2 - y^2 = 53$
Then $y =$ |
| 2. The sides of a right triangle are integers. If one leg is 9 then the other leg is: | 4. A right triangle with integer sides has a hypotenuse of 113. The smallest leg is: |

5. An acute triangle has integer side lengths of 4,7,and x. The smallest value for x is:
6. An acute triangle has integer side lengths of 4,7,and x. The largest value for x is:
7. **x,y are integers with $x^2 - y^2 = -67$ then x is:**
8. **An obtuse triangle has integer side lengths of x,7, and 11. The smallest value of x is:**
9. $a^2 + b^2 = 113^2$ where $0 < a < b$ and a, b are integers. Then $a =$
10. The sides of a right triangle are x,7,and 11. If $x < 7$ and $x = a\sqrt{2}$ then $a =$
11. An acute triangle has integer sides of 2,7,and x.
The largest value of x is:
12. An obtuse triangle has integer sides of 6,x,and 11. The smallest value of x is:
13. An acute triangle has integer sides of 7, 11,and x. The smallest value of x is:
14. **An obtuse triangle has integer sides of 8,15, and x. The smallest value of x is:**
15. The sides of a right triangle are integral. If one leg is 13, find the length of the other leg:
16. A right triangle has integer side lengths of 7,x,and 25. Its area is:

2.2.8 Equilateral Triangle Formulas

Area of an Equilateral Triangle when knowing the side-length s :

$$\text{Area} = \frac{s^2 \cdot \sqrt{3}}{4}$$

Area of an Equilateral Triangle when knowing the height h :

$$\text{Area} = \frac{h^2 \cdot \sqrt{3}}{3}$$

Finding the height when given the side length s :

$$\text{Height} = \frac{s \cdot \sqrt{3}}{2}$$

Example:

An equilateral triangle's perimeter is 12. It's area is $4k \cdot \sqrt{3}$. What is k?

$$s = \frac{12}{3} = 4 \text{ so } A = \frac{4^2 \cdot \sqrt{3}}{4} = 4\sqrt{3} \Rightarrow k = 1$$

Example:

An equilateral triangle has a height of 4, what is its side length?

$$h = 4 = \frac{\sqrt{3} \cdot s}{2} \Rightarrow s = \frac{4 \cdot 2}{\sqrt{3}} = \frac{8 \cdot \sqrt{3}}{3}$$

Here are some practice problems for this formula:

Problem Set 2.2.8

- The sides of an equilateral triangle are $2\sqrt{3}$ cm, then its height is:
- The area of an equilateral triangle is $9\sqrt{3}\text{cm}^2$, then its side length is:
- If the area of an equilateral triangle is $3\sqrt{3}\text{ft}^2$ then its side length is:
- The height of an equilateral triangle is 12 in. Its area is $4k\sqrt{3}$, $k =$
- The perimeter of an equilateral triangle is 12 cm. Its area is $k\sqrt{3}\text{cm}^2$. $k =$
- Find the perimeter of an equilateral triangle whose area is $9\sqrt{3}\text{cm}^2$:
- The area of an equilateral triangle is $3\sqrt{3}\text{in}^2$. Its height is:
- An equilateral triangle has an area of $27\sqrt{3}\text{cm}^2$. Its height is:

2.2.9 Formulas of Solids

Usually basic formulas for spheres, cubes, cones, and cylinders are fair game for the number sense test. In order to solve these problems, memorize the following table:

Type of Solid	Volume	Surface Area
Cube	s^3	$6s^2$
Sphere	$\frac{4}{3}\pi r^3$	$4\pi r^2$
Cone	$\frac{1}{3}\pi r^2 h$	$\pi r l + \pi r^2$
Cylinder	$\pi r^2 h$	$2\pi r h$

In the above formulas, s is the side-length, r is the radius, h is the height, and l is the slant height. In addition to knowing the above formulas, a couple of other ones are:

Face Diagonal of a Cube = $s\sqrt{2}$
 Body Diagonal of a Cube = $s\sqrt{3}$

Problem Set 2.2.9

- Find the surface area of a square who's side length is 11in. :
- Find the surface area of a sphere who's radius is 6in. :
- If the radius of a sphere is tripled, then the volume is multiplied by:
- The total surface area of a cub with an edge of 4 inches is:

5. A cube has a volume of 512cm^3 . The area of the base is:
6. A cube has a surface area of 216cm^2 . The volume of the cube is:
7. If the total surface area of a cube is 384cm^2 then the volume of the cube is:
8. Find the volume of a cube with an edge of 12 cm.:
9. A tin can has a diameter of 8 and a height of 14. The volume is $k\pi$, $k =$

2.2.10 Combinations and Permutations

For most this is just a refresher on the definitions of ${}_nC_k$ and ${}_nP_k$:

$${}_nC_k = \frac{n!}{k! \cdot (n-k)!}$$

$${}_nP_k = \frac{n!}{(n-k)!}$$

Here is an example:

$${}_7C_4 = \frac{7!}{4!(7-4)!} = \frac{7 \cdot 6 \cdot 5}{3 \cdot 2} = \mathbf{35}$$

With combinations and permutations (and factorials in general) you want to look at ways of canceling factors from the factorial to ease in calculation. In addition, the following is a list of the factorials which should be memorized for quick access:

$3! = 6$	$4! = 24$	$5! = 120$	$6! = 720$
$7! = 5040$	$8! = 40320$	$9! = 362880$	$10! = 3628800$

Another often tested principle on Combinations is that:

$${}_nC_k = {}_nC_{n-k}$$

The above will show up in the form of questions like this:

Problem: ${}_5C_2 = {}_5C_k$. $k = ?$

Solution: Using the above formula, you know that $k = 5 - 2 = \mathbf{3}$.

Another often tested question on Combinations and Permutations is when you divide one by another:

$$\frac{{}_nC_k}{{}_nP_k} = \frac{1}{k!}$$

$$\frac{{}_nP_k}{{}_nC_k} = k!$$

Problem Set 2.2.10

1. ${}_5P_3 =$

12. ${}_4P_2 \div {}_4C_2 =$

2. ${}_5C_3 =$

13. ${}_6P_3 \div {}_6C_3 =$

3. ${}_6C_3 =$

14. ${}_7P_4 \div {}_7C_3 =$

4. ${}_7C_4 =$

15. ${}_8C_5 \div {}_8P_5 =$

5. ${}_7P_4 =$

16. ${}_9P_3 \div {}_9C_3 =$

6. ${}_6P_2 =$

17. ${}_4P_3 \div {}_3P_2 =$

7. ${}_8C_6 =$

18. ${}_4C_3 \times {}_3C_2 =$

8. ${}_5C_2 =$

19. ${}_5P_3 \times {}_4P_2 =$

9. ${}_8P_3 =$

20. ${}_6C_3 \div {}_6P_3 =$

10. ${}_8C_3 =$

21. ${}_6C_1 + {}_4P_1 =$

11. ${}_9C_2 =$

22. $({}_5C_2)({}_5P_2) =$

2.2.11 Trigonometric Values


Trigonometry problems have been increasingly popular for writers of the number sense test. Not only are they testing the basics of sines, cosines, and tangents of special angles (30° , 45° , 60° , 90° , and variations in each quadrant) but also the trigonometric reciprocals (cosecant, secant, and cotangent).

First, let's look at the special angles in the first quadrant where all values of the trigonometric functions are positive. In the table, each trigonometric function is paired below with its reciprocal:

Trig Function	0°	30°	45°	60°	90°
sin	0	$\frac{1}{2}$	$\frac{\sqrt{2}}{2}$	$\frac{\sqrt{3}}{2}$	1
csc	Undefined	2	$\sqrt{2}$	$\frac{\sqrt{3}\cdot 2}{3}$	1
cos	1	$\frac{\sqrt{3}}{2}$	$\frac{\sqrt{2}}{2}$	$\frac{1}{2}$	0
sec	1	$\frac{\sqrt{3}\cdot 2}{3}$	$\sqrt{2}$	2	Undefined
tan	0	$\frac{\sqrt{3}}{3}$	1	$\sqrt{3}$	Undefined
cot	Undefined	$\sqrt{3}$	1	$\frac{\sqrt{3}}{3}$	0

All of those can be derived using the memorable “SOHCAHTOA” technique to special right triangles (it assumes that one can do this, so it is omitted in this text. If help is needed, see any elementary geometry book.). In addition, it is clear that the values at the reciprocal trigonometric function is just the multiplicative inverse (that’s why they are called *reciprocal* trigonometric functions!).

Now to find the values of trigonometric functions in any quadrant it is essential to remember two things. The first is you need to get the sign straight of the values depending on what quadrant you are in. The following plot and mnemonic device will help with getting the sign correct:



The above corresponds to which trigonometric functions (and their reciprocals) are positive in which quadrants. Now if you forget this, you can take the first letter of each function in their respected quadrants and remember the mnemonic device of “**All Students Take Calculus**” to remember where each function is positive.

The second challenge to overcome in computing each Trigonometric Function at any angle is to learn how to reference each angle to its first quadrant angle, so that the chart above could be used. The following chart will help you find the appropriate reference angle depending on what quadrant you are in. Assume that you are given an angle θ which resides in each of the quadrants mentioned. The following would be it’s reference angle:

	Quadrant I	Quadrant II	Quadrant III	Quadrant IV
Reference Angle:	θ	$180^\circ - \theta$	$\theta - 180^\circ$	$360^\circ - \theta$

So now we have enough information to compute any trigonometric function at any angle. Let's look at a couple of problems:

Problem: $\sin(210^\circ)$

Solution: Now you know the angle is in Quadrant-III, so the result will be negative (only cosine is positive in Q-III). Now to find the reference angle is just $\theta - 180^\circ = 210^\circ - 180^\circ = 30^\circ$. So the $\sin(30^\circ)$ from the table is $\frac{1}{2}$ so the answer is: $\sin(210^\circ) = -\frac{1}{2}$.

Problem: $\cot(135^\circ)$

Solution: So the \cot/\tan function is negative in Q-II. To find the reference angle, it is simply $180^\circ - \theta = 180^\circ - 135^\circ = 45^\circ$. Now the $\cot(45^\circ) = 1$ (from the table) so the answer is: $\cot(135^\circ) = -1$.

Problem: $\cos(-30^\circ)$

Solution: So an angle of $-30^\circ = 330^\circ$ which is in Q-IV where cosine is positive. Now to find the reference angle you just do $360^\circ - \theta = 360^\circ - 330^\circ = 30^\circ$, and $\cos(30^\circ) = \frac{\sqrt{3}}{2}$. So the answer is just $\cos(-30^\circ) = \frac{\sqrt{3}}{2}$.

It should be noted that all of these problems have been working with degrees. Students should familiarize themselves with using radians as well using the conversion rate of: $\pi = 180^\circ$. So an angle (given in radians) of $\frac{\pi}{6} = \frac{180^\circ}{6} = 30^\circ$.

It is great for all students to practice solving these types of problems. The following are some practice problems. If more are needed, just consult any elementary geometric textbook or pre-calculus textbook.

Problem Set 2.2.11

1. $\sin(-30^\circ) =$

10. $\cos(\sec^{-1} 3) =$

2. $\cos \theta = .375$ then $\sec \theta =$

11. $\frac{5\pi}{8} = \text{-----}^\circ$

3. $\sin(3\pi) =$

12. $\frac{\pi}{5} = \text{-----}^\circ$

4. $\tan(225^\circ) =$

13. $\cos(\sin^{-1} 1) =$

5. $\sin(\sin^{-1} \frac{1}{2}) =$

14. $\tan(-45^\circ) =$

6. $\sin \theta = -.1$ then $\csc \theta =$

15. $\sin(-\pi) =$

7. $\sin \frac{11\pi}{6} =$

16. $\cos(-300^\circ) =$

8. $\cos(-5\pi) =$

17. $\sin^{-1}(\sin 1) =$

9. $\frac{\pi}{18} = \text{-----}^\circ$

18. $\csc(-150^\circ) =$

19. $\sec(120^\circ) =$
20. $\tan(-225^\circ) =$
21. $\frac{3\pi}{5} = \text{-----}^\circ$
22. $\tan(-45^\circ) =$
23. $\tan(315^\circ) =$
24. If $0^\circ < x < 90^\circ$ and $\tan x = \cot x, x =$
25. $280^\circ = k\pi$ then $k =$
26. $\tan \frac{5\pi}{4} =$
27. $\cos \theta = .08333\dots$ then $\sec \theta =$
28. $\sin(5\pi) + \cos(5\pi) =$
29. $\sec(60^\circ) =$
30. $12^\circ = \frac{\pi}{k}, k =$
31. $\cos \theta = -.25$ then $\sec \theta =$
32. $\tan^2 60^\circ =$
33. $1.25\pi = \text{-----}^\circ$
34. $\cot^2 60^\circ =$
35. $\sin[\cos^{-1}(\frac{\sqrt{2}}{2})] =$
36. $\cos(-3\pi) - \sin(-3\pi) =$
37. $\cos(\frac{-4\pi}{3}) + \sin(\frac{-5\pi}{6}) =$
38. $2 \sin 120^\circ \cos 30^\circ =$
39. $\cos(240^\circ) - \sin(150^\circ) =$
40. $\sin(\cos^{-1} \frac{\sqrt{3}}{2}) =$
41. $\sin(\cos^{-1} 1) =$
42. If $\csc \theta = -3$, where $270^\circ < \theta < 300^\circ$, then $\sin \theta =$
43. $\sin(\frac{-7\pi}{6}) - \cos(\frac{-2\pi}{3}) =$
44. $\sec \theta = -3$, θ is in **QIII**, then $\cos \theta =$
45. $\cos \frac{5\pi}{6} \times \sin \frac{2\pi}{3} =$
46. $\sin \frac{3\pi}{4} \times \cos \frac{5\pi}{4} =$
47. $\sin 30^\circ + \cos 60^\circ = \tan x$
 $0^\circ \leq x \leq 90^\circ, x =$
48. $\cos\left(\sin^{-1} \frac{\sqrt{3}}{2}\right) =$
49. $\sin(\frac{-\pi}{3}) \times \sin(\frac{\pi}{3}) =$
50. $\cos(120^\circ) \times \cos(120^\circ) =$
51. $216^\circ = k\pi, k =$
52. $\cos(\frac{-2\pi}{3}) \times \cos(\frac{4\pi}{3}) =$
53. $\tan(30^\circ) \times \cot(60^\circ) =$
54. $\cos(\frac{-\pi}{3}) \times \cos(\frac{\pi}{3}) =$
55. $\sin \frac{\pi}{6} + \cos \frac{\pi}{3} = \tan \frac{\pi}{k}$
then k =
56. $\cos^{-1} .8 + \cos^{-1} .6 = k\pi$ then $k =$
57. $\sin(300^\circ) \times \cos(330^\circ) =$
58. $\sin(\frac{-\pi}{6}) \times \cos(\frac{\pi}{3}) =$
59. $630^\circ = k\pi, k =$

2.2.12 Trigonometric Formulas

Recently, questions involving trigonometric functions have encompassed some basic trigonometric identities. The most popular ones tested are included here:

The Fundamental Identity

$\sin^2 + \cos^2 = 1$ with subsequent variants:
 $1 + \cot^2 = \csc^2$
 $\tan^2 + 1 = \sec^2$

Sum to Difference Formulas

$\sin(a \pm b) = \sin(a) \cos(b) \pm \sin(b) \cos(a)$
 $\cos(a \pm b) = \cos(a) \cos(b) \mp \sin(a) \sin(b)$

Double Angle Formulas

$\sin(2a) = 2 \sin(a) \cos(a)$
 $\cos(2a) = \cos^2(a) - \sin^2(a)$ with variants:
 $\cos(2a) = 1 - 2 \sin^2(a)$
 $\cos(2a) = 2 \cos^2(a) - 1$

Sine \rightarrow Cosine

$\sin(90^\circ - \theta) = \cos(\theta)$

Most of the time, using trigonometric identities will not only aid in speed but will also be necessary. Take for example this example:

$$\sin(10^\circ) \cos(20^\circ) + \sin(20^\circ) \cos(10^\circ)$$

Without using the sum to difference formula, this would be impossible to calculate, however after using the formula you get:

$$\sin(10^\circ) \cos(20^\circ) + \sin(20^\circ) \cos(10^\circ) = \sin(10^\circ + 20^\circ) = \sin(30^\circ) = \frac{1}{2}$$

The following are some practice problems using these identities:

Problem Set 2.2.12

1. $\cos^2 30^\circ + \sin^2 30^\circ =$

2. $\cos^2 30^\circ - \sin^2 30^\circ =$

3. $2 \sin 15^\circ \cos 15^\circ =$

4. $2 \sin 30^\circ \sin 30^\circ - 1 =$

5. $1 - \sin^2 30^\circ =$

6. $\cos 22^\circ = \sin \theta, 0^\circ < \theta < 90^\circ, \theta =$

7. $[2 \sin \frac{\pi}{3} \cos \frac{\pi}{3}]^2 =$

8. $2 \sin 15^\circ \cos 15^\circ - 1 =$

9. $3 \csc^2 45^\circ - 3 \cot^2 45^\circ =$

10. $\cos^2 30^\circ - \sin^2 30^\circ =$

11. $\sin 105^\circ \cos 105^\circ =$

12. $\sin 38^\circ = \cos \theta, 270^\circ < \theta < 360^\circ, \theta =$

13. $\sin 30^\circ \cos 60^\circ - \sin 60^\circ \cos 30^\circ =$

14. $2 \cos^2 \frac{\pi}{6} - 1 =$

15. $(1 - \sin 60^\circ)(1 + \sin 60^\circ) =$

16. $2 \sin 15^\circ \sin 75^\circ =$

17. $(\sin \frac{\pi}{3} - \cos \frac{\pi}{3})(\sin \frac{\pi}{3} + \cos \frac{\pi}{3}) =$

18. If $\sin(A) = \frac{3}{5}$, then $\cos(2A) =$

19. $1 - 2 \sin^2 \frac{\pi}{6} =$

20. $\cos 75^\circ \sin 75^\circ =$

21. $\sin 15^\circ \cos 45^\circ - \sin 45^\circ \cos 15^\circ =$

22. $2 - 4 \sin^2 30^\circ =$

23. $\cos 95^\circ \cos 25^\circ - \sin 95^\circ \sin 25^\circ =$

24. $\sin \frac{\pi}{6} + \cos \frac{\pi}{3} =$

25. $\cos 15^\circ \sin 45^\circ - \cos 45^\circ \sin 15^\circ =$

26. $(\sin \frac{\pi}{6} - \cos \frac{\pi}{6})(\sin \frac{\pi}{6} + \cos \frac{\pi}{6}) =$

27. $2 \tan^2 \theta - 2 \sec^2 \theta =$

2.2.13 Graphs of Sines/Cosines

Popular questions for the last column involve determining amplitudes, periods, phase shifts, and vertical shifts for plots of sines/cosines. If you haven't been introduced this in a pre-calculus class, use the following as a rough primer:

The general equation for any sine/cosine plot is:

$$y = A \sin[B(x - C)] + D$$

Amplitude: $|A|$

Period: $\frac{2\pi}{B}$

Phase Shift: C

Vertical Shift: D (Up if > 0 , Down if < 0)

Example: Find the period of $y = 3 \sin(\pi x - 2) + 8$.

Solution: We need the coefficient in front of x to be 1, so we need to factor out π , making the graph: $y = 3 \sin[\pi(x - \frac{2}{\pi})] + 8$. Now we can apply the above table to see that the period = $\frac{2\pi}{\pi} = 2$. The other characteristics of the graph is that the amplitude = 3, the phase shift = $\frac{2}{\pi}$, and it is vertically shifted by 8 units.

Here are some more practice problems:

Problem Set 2.2.13

1. What is the amplitude of $y = 4 \cos(2x) + 1$:
2. The graph of $y = 2 - 3 \cos[2(x - 5)]$ has a horizontal displacement of:
3. The graph of $y = 2 - 2 \cos[3(x - 5)]$ has a vertical shift of:
4. The amplitude of $y = 2 - 3 \cos[4(x + 5)]$ is:
5. The period of $y = 5 \cos[\frac{1}{4}(x + 3\pi)] + 2$ is $k\pi, k =$
6. The phase shift of $y = 5 \cos[4(x + 3)] - 2$ is:
7. The amplitude of $y = 2 - 5 \cos[4(x - 3)]$ is:
8. The vertical displacement of $y = 5 \cos[4(x + 3)] - 2$ is:
9. The phase shift of $f(x) = 2 \sin(3x - \frac{\pi}{2})$ is $k\pi, k =$
10. **The period of $y = 2 - 3 \cos(4\pi x + 2\pi)$ is:**
11. The period of $y = 2 + 3 \sin(\frac{x}{5})$ is:
12. The graph of $y = 1 - 2 \cos(3x + 4)$ has an amplitude of:

2.2.14 Vertex of a Parabola

This question was much more popular on tests from the 90's, but it is being resurrected on some of the more recent tests. When approached with a parabola in the form of $f(x) = Ax^2 + Bx + x$, the coordinate of the vertex is:

$$(h, k) = (\frac{-B}{2A}, f(\frac{-B}{2A})).$$

Example: Find the y-coordinate of the vertex of the parabola whose equation is $y = 3x^2 - 12x + 16$.

Solution: $x = \frac{-(-12)}{2 \cdot 3} = 2 \Rightarrow y = 3 \cdot 2^2 - 12 \cdot 2 + 16 = 4.$

It should be noted that if the parabola is in the form $x = ay^2 + by + c$, then the vertex is: $(h, k) = (f(\frac{-b}{2a}), \frac{-b}{2a})$. (due to a rotation of axis).

The following are some practice problems:

Problem Set 2.2.14

1. The vertex of the parabola $y = 2x^2 + 8x - 1$ is $(h, k), k =$
2. The vertex of $y = x^2 - 2x - 4$ is $(h, k), k =$
3. If $g(x) = 2 - x - x^2$, then the axis of symmetry is $x =$

2.2.15 Discriminant and Roots

A very popular question is, when given a quadratic equation, determining the value of an undefined coefficient so that the roots are distinct/equal/complex. Take the following question:

Find the value for k such that the quadratic $3x^2 - x - 2k = 0$ has equal roots.

Well we know from the quadratic equation that the roots of a general polynomial $ax^2 + bx + c = 0$ can be determined from:

$$r_{1,2} = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

So we know from this that:

$$\text{Distinct Roots:} \quad b^2 - 4ac > 0$$

$$\text{Equal Roots:} \quad b^2 - 4ac = 0$$

$$\text{Complex Conjugate Roots:} \quad b^2 - 4ac < 0$$

So in our case we need to find the value of k such that the discriminant ($b^2 - 4ac$) is equal to zero.

$$b^2 - 4ac = 1^2 - 4 \cdot 3 \cdot (-2k) = 0 \Rightarrow k = \frac{-1}{4 \cdot 3 \cdot 2} = \frac{-1}{24}$$

The following are some more practice problems:

Problem Set 2.2.15

1. For $2x^2 - 4x - k = 0$ to have 2 equal roots, the smallest value of k is:
2. For $3x^2 - x - 2k = 0$ to have equal roots k has to be:
3. For $3x^2 - 2x + 1 - k = 0$ to have equal roots, k has to be:
4. The discriminant of $2x^2 - 3x = 1$ is:
5. For what value of k does $3x^2 + 4x + k = 0$ have equal roots:
6. For $x^2 - 2x - 3k = 0$ to have one real solution k has to be:

3 Miscellaneous Topics

3.1 Random Assortment of Problems

3.1.1 GCD/GCF and LCM

How finding the Greatest Common Divisor (or GCD) is taught in classes usually involves prime factorizing the two numbers and then comparing powers of exponents. However, this is not the most efficient way of doing it during a number sense competition. One of the quickest way of doing it is by employing Euclid's Algorithm who's method won't be proven here (if explanation is necessary, just google to find the proof). The following outlines the procedure:

1. Arrange the numbers so that $n_1 < n_2$ then find the remainder when n_2 is divided by n_1 and call it r_1 .
2. Now divide n_1 by r_1 and get a remainder of r_2 .
3. Continue the procedure until any of the remainders are 0 and the number you are dividing by is the GCD or when you notice what the GCD of any pair of numbers is.

Let's illustrate with an example:

$GCD(36, 60) =$ Well, when 60 is divided by 36 it leaves a remainder of 24. So:
 $GCD(36, 60) = GCD(24, 36)$. Continuing the procedure, when 36 is divided by 24 it leaves a remainder of 12. So:
 $GCD(36, 60) = GCD(24, 36) = GCD(12, 24)$ Which from here you can tell the GCD is **12**. You could also have stopped after the first step when you notice that the $GCD(24, 36)$ is 12, and you wouldn't have to continue the procedure.

Here is another example:

$GCD(108, 140) = GCD(32, 108) = GCD(12, 32) = GCD(8, 12) = GCD(4, 8) = 4$
If at any point in that process you notice what the GCD of the two numbers is by observation, you can cut down on the amount of steps in computation.

For computing the LCM between two numbers a and b , I use the formula:

$$LCM(a, b) = \frac{a \times b}{GCD(a, b)}$$

So to find what the LCM is, we must first compute the GCD. Using a prior example, let's calculate the $LCM(36, 60)$:

$$LCM(36, 60) = \frac{36 \times 60}{12} = 3 \times 60 = \mathbf{180}$$

The procedure is simple enough, let's do one more example by finding the LCM of 44 and 84:
 $GCD(44, 84) = GCD(40, 44) = GCD(4, 40) = 4$ therefore

$$LCM(44, 84) = \frac{44 \times 84}{4} = 11 \times 84 = \mathbf{924}$$

It should be noted that there are some questions concerning the GCD of more than two numbers (usually not ever more than three). The following outlines the procedure which should be followed:

1. Find the GCD of two of the numbers.
2. Find the LCM of those two numbers by using the GCD and the above formula.
3. Calculate the GCD of the LCM of those two numbers and the third number.

It should be noted that usually one of the numbers is a multiple of another, thus leaving less required calculations (because the LCM between two numbers which are multiples of each other is just the larger of the two numbers).

The following are some more practice problems for finding GCDs and LCMs using this method:

Problem Set 3.1.1

1. The GCF of 35 and 63 is:
2. The LCM of 64 and 20 is:
3. The LCM of 27 and 36 is:
4. The GCF of 48 and 72 is:
5. The GCD of 27 and 36 is:
6. The LCM of 63 and 45 is:
7. The GCD of 132 and 156 is:
8. **The LCM of 57 and 95 is:**
9. The GCD of 52 and 91 is:
10. The LCM of 52 and 28 is:
11. The GCD of 48 and 54 is:
12. The GCD of 54 and 36 is:
13. The LCM of 27 and 36 is:
14. The LCM of 108 and 81 is:
15. The GCD of 28 and 52 is:
16. The LCM of 51 and 34 is:
17. **The LCM of $2^3 \times 3^2$ and $2^2 \times 3^3$ is:**
18. The LCM of 28 and 42 is:
19. The LCM of 54 and 48 is:
20. The GCF of 84 and 70 is:
21. **The GCF of 132 and 187 is:**
22. The LCM of 48 and 72 is:
23. The GCF of 51, 68, and 85 is:
24. The GCF(24,44)-LCM(24,44)=
25. The LCM of 16, 20, and 32 is:
26. The GCD(15,28) times LCM(15,28) is:
27. The LCM of 12, 18, and 20 is:
28. The LCM of 14, 21, and 42 is:
29. The LCM of 8, 18, and 32 is:
30. The GCD(15,21)+LCM(15,21)=
31. The GCF of 44, 66, and 88 is:
32. The product of the GCF and LCM of 21 and 33 is:
33. The LCM of 16, 32, and 48 is:
34. The GCD(18,33)+LCM(18,33)=

35. The LCM of 14, 28, and 48 is:
36. **The LCM(21,84)-GCF(21,84)=**
37. The LCM of 24, 36, and 48 is:
38. The GCD(16,20)-LCM(16,20)=
39. The GCF of 42, 28, and 56 is:
40. **The product of the GCF and LCM of 24 and 30 is:**
41. The LCM of 36, 24, and 20 is:
42. **The LCM of 28, 42, and 56 is:**

3.1.2 Perfect, Abundant, and Deficient Numbers

For this section let's begin with the definitions of each type.

A perfect number has the sum of the proper divisors equal to itself. The first three perfect numbers are 6 ($1 + 2 + 3 = 6$), 28 ($1 + 2 + 4 + 7 + 14 = 28$), and 496 ($1 + 2 + 4 + 8 + 16 + 31 + 62 + 124 + 248 = 496$). Notice that there are really *only* two perfect numbers that would be reasonable to test on a number sense test (6 and 28 should be memorized as being perfect).

An abundant number has the sum of the proper divisors greater than itself. Examples of an abundant number is 12 ($1 + 2 + 3 + 4 + 6 = 16 > 12$) and 18 ($1 + 2 + 3 + 6 + 9 = 21 > 18$). An interesting property of abundant numbers is that any multiple of a perfect or abundant number is abundant. Knowing this is *very* beneficial to the number sense test.

As you can assume through the process of elimination, a deficient number has the sum of the proper divisors less than itself. Examples of these include any prime number (because they have only one proper divisor which is 1), 10 ($1 + 2 + 5 = 8 < 10$), and 14 ($1 + 2 + 7 = 10 < 14$) just to name a few. An interesting property is that any power of a prime is deficient (this is often tested on the number sense test).

3.1.3 Sum and Product of Coefficients in Binomial Expansion

From the binomial expansion we know that:

$$\begin{aligned} (ax + by)^n &= \sum_{k=0}^n \binom{n}{k} (ax)^{n-k} (by)^k \\ &= \binom{n}{0} a^n \cdot x^n + \binom{n}{1} a^{n-1} b^1 \cdot x^{n-1} y^1 + \dots + \binom{n}{n} b^n y^n \end{aligned}$$

From here we can see that the sum of the coefficients of the expansion is:

$$\sum_{k=0}^n \binom{n}{k} a^{n-k} b^k$$

Where we can retrieve these sums by setting $x = 1$ and $y = 1$. So the Sum of the coefficients is just $(a + b)^n$!

Here is an example to clear things up:

Example: Find the Sum of the Coefficients of $(x + y)^6$.

Solution: Well let $x = 1$ and $y = 1$ which lead to the Sum of the Coefficients = $(1 + 1)^6 = \mathbf{64}$.

An interesting side note on this is when asked to find the Sum of the Coefficients of $(x - y)^n$ it will always be 0 because by letting $x = 1$ and $y = 1$ you get the Sum of the Coefficients = $(1 - 1)^n = \mathbf{0}$.

As for the product of the coefficients, there are no easy way to compute them. The best method is to memorize some of the first entries of the Pascal triangle:

$$\begin{array}{c} 1 \\ 1 \ 1 \\ 1 \ 2 \ 1 \\ 1 \ 3 \ 3 \ 1 \\ 1 \ 4 \ 6 \ 4 \ 1 \\ 1 \ 5 \ 10 \ 10 \ 5 \ 1 \\ 1 \ 6 \ 15 \ 20 \ 15 \ 6 \ 1 \end{array}$$

Here are some more practice to get acquainted with both the sum and product of coefficients:

Problem Set 3.1.3

1. The sum of the coefficients in the expansion of $(5x - 9y)^3$ is:
2. The sum of the coefficients in the expansion of $(5x + 7y)^3$ is:
3. The sum of the coefficients in the expansion of $(x - y)^3$ is:
4. The sum of the coefficients in the expansion of $(a + b)^3$ is:
5. The sum of the coefficients in the expansion of $(x + y)^6$ is:
6. The sum of the coefficients in the expansion of $(x + y)^2$ is:
7. **The sum of the coefficients in the expansion of $(a + b)^5$ is:**
8. The sum of the coefficients in the expansion of $(a - b)^4$ is:
9. The sum of the coefficients in the expansion of $(3x - y)^4$ is:
10. The product of all the coefficients in the expansion $(x + y)^4$ is:
11. The product of the coefficients in the expansion of $(2a + 2b)^2$ is:
12. The product of the coefficients in the expansion of $(a + b)^3$ is:
13. The product of the coefficients in the expansion of $(a - b)^4$ is:
14. The product of the coefficients in the expansion of $(3a + 3b)^2$ is:
15. **The product of the coefficients in the expansion of $(a + b)^5$ is:**
16. The product of the coefficients in the expansion of $(a - b)^2$ is:

17. The product of the coefficients in the expansion of $(4a - 3b)^2$ is:
18. The sum of the coefficients in the expansion of $(x^2 - 6x + 9)^2$ is:

19. **The product of the coefficients in the expansion of $(4x + 5)^2$ is:**

3.1.4 Sum/Product of the Roots

Define a polynomial by $p_n = a_n x^n + a_{n-1} x^{n-1} + a_{n-2} x^{n-2} \cdots a_1 x^1 + a_0 = 0$. The three most popular questions associated with the number sense test concerning roots of polynomials are: sum of the roots, sum of the roots taken two at a time, and product of the roots. For the polynomial p_n these values are defined by:

Sum of the roots: $\frac{-a_{n-1}}{a_n}$

Sum of the roots taken two at a time: $\frac{a_{n-2}}{a_n}$

Product of the roots: If n is even $\frac{a_0}{a_n}$

If n is odd $\frac{-a_0}{a_n}$

Let's see what this means for our generic quadratics/cubics: $p_2 = ax^2 + bx + c = 0$ and $p_3 = ax^3 + bx^2 + cx = 0$

Sum of the roots: $\frac{-b}{a}$

$p_2 = ax^2 + bx + c = 0$

Product of the roots: $\frac{c}{a}$

Sum of the roots: $\frac{-b}{a}$

$p_3 = ax^3 + bx^2 + cx = 0$

Product of the roots taken two at a time: $\frac{c}{a}$

Product of the roots: $\frac{-d}{a}$

Since the quadratic only has two roots, the sum of the roots taken two at a time happens to be the product of the roots. You can extend the same procedure for polynomials of any degree, keeping in mind the alternating signs for the product of the roots. The following are practice problems:

Problem Set 3.1.4

1. The sum of the roots of $2x^2 - 3x + 1 = 0$ is:

3. The sum of the roots of $3x^3 - 2x^2 + x - 4 = 0$ is:

2. The sum of the roots of $(x - 4)(x - 5) = 0$ is:

4. The product of the roots of $x^2 + 3x = 7$ is:

5. The sum of the roots of $x^2 - 9 = 0$ is:
6. The sum of the roots of $4x^2 + 3x = 2$ is:
7. The sum of the roots of $(2x - 3)^2 = 0$ is:
8. The product of the roots of $5x^3 - 8x^2 + 2x + 3 = 0$ is:
9. The product of the roots of $4x^3 - 3x^2 + 2x - 1 = 0$ is:
10. The sum of the roots of $3x^3 + 2x^2 = 9$ is:
11. **The sum of the roots of $x^3 - 13x = 12$ is:**
12. Let R,S,T be the roots of $2x^3 + 4x = 5$. Then $R \times S \times T =$
13. The product of the roots of $5x^3 + 4x - 3 = 0$ is:
14. The sum of the roots of $(3x - 2)(2x + 1) = 0$ is:
15. The sum of the product of the roots taken two at a time of $2x^3 + 4x^2 - 6x = 8$ is:
16. The sum of the roots of $2x^3 + 4x^2 - 3x + 5 = 0$ is:
17. The product of the roots of $(2x - 1)(3x + 2)(4x - 3) = 0$ is:
18. Let R,S,T be the roots of $2x^3 + 4x = 5$. Then $RS + RT + ST =$
19. **The equation $2x^3 - bx^2 + cx = d$ has roots r,s,t and $rst=3.5$, then $d =$**
20. The sum of the roots of $3x^2 - bx + c = 0$ is -12 then $b =$
21. If r,s, and t are the roots of the equation $2x^3 - 4x^2 + 6x = 8$ then $rs + rt + st =$
22. The sum of the roots of $4x^3 + 3x^2 - 2x - 1 = 0$ is:
23. The product of the roots of $4x^3 - 3x^2 + 2x + 1 = 0$ is:
24. The sum of the roots of $5x^3 + 4x - 3 = 0$ is:
25. **The equation $2x^3 - bx^2 + cx = d$ has roots r,s,t. If $r + s + t = -2$ then $b =$**

3.1.5 Finding Units Digit of x^n

This is a common problem on the number sense test which seems considerably difficult, however there is a shortcut method. Without delving too much into the modular arithmetic required, you can think of this problem as exploiting patterns. For example, let's find the units digit of 3^{47} , knowing:

3^1	3	Units Digit: 3
3^2	9	Units Digit: 9
3^3	27	Units Digit: 7
3^4	81	Units Digit: 1
3^5	243	Units Digit: 3
3^6	729	Units Digit: 9
3^7	2187	Units Digit: 7
3^8	6561	Units Digit: 1

From this you can see it repeats every 4th power.

So in order to see what is the units digit you can divide the power in question by 4 then see what the remainder r is. Then to find the appropriate units digit, you look at the units digit of 3^r . For example, the units digit for 3^5 could be found by saying $5 \div 4$ has a remainder of 1 so, the units digit of 3^5 corresponds to that of 3^1 which is **3**. So to reiterate, the procedure is:

1. For low values of n , compute what the units digit of x^n is.
2. Find out how many unique integers there are before repetition (call it m).
3. Find the remainder when dividing the large n value of interest by m (call it r)
4. Find the units digit of x^r , and that's your answer.

So for our example of 3^{47} :

$47 \div 4$ has a remainder of 3

3^3 has the units digit of **7**

Other popular numbers of interest are:

Numbers	Repeating Units Digits	Number of Unique Digits
Anything ending in 2	2, 4, 8, 6	4
Anything ending in 3	3, 9, 7, 1	4
Anything ending in 4	4, 6	2
Anything ending in 5	5	1
Anything ending in 6	6	1
Anything ending in 7	7, 9, 3, 1	4
Anything ending in 8	8, 4, 2, 6	4
Anything ending in 9	9, 1	2

Using the above table, we can calculate the units digit of any number raised to any power relatively simple. To show this, find the units digit of 27^{63} :

From the table, we know it repeats every 4^{th} power, so: $63 \div 4 \Rightarrow r = 3$
 $r = 3$ corresponds to 7^3 which ends in a **3**

This procedure is also helpful with raising the imaginary number i to any power. Remember from Algebra:

i^1	i
i^2	-1
i^3	$-i$
i^4	1
i^5	i
i^6	-1
i^7	$-i$
i^8	1

So, after noticing that it repeats after every 4^{th} power, we can compute for example i^{114} .

$$114 \div 4 \text{ has a remainder of } 2 \Rightarrow i^2 = -1$$

The following are examples of these types of problems:

Problem Set 3.1.5

- Find the units digit of 19^7 :
- Find the units digit of 17^6 :
- Find the units digit of 8^8 :
- Find the units digit of 7^7 :
- Find the units digit of 13^{13} :
- Find the units digit of 17^5 :
- $i^{78} =$
- $i^{66} =$
- Find the units digit of 16^5 :

3.1.6 Exponent Rules

These problems are usually on the third column, and if you know the basics of exponential rules they are easy to figure out. The rules to remember are as followed:

$$x^a \cdot x^b = x^{a+b} \quad \frac{x^a}{x^b} = x^{a-b} \quad (x^a)^b = x^{ab}$$

The following are problems concerning each type:

Product Rule: Let $3^x = 70.1$, then $3^{x+2} = ?$

Solution: $3^{x+2} = 3^x \cdot 3^2 = 70.1 \cdot 9 = \mathbf{630.9}$

Quotient Rule: Let $5^x = 2$, represent 5^{x-2} as a decimal.

Solution: $5^{x-2} = \frac{5^x}{5^2} = \frac{2}{25} = \mathbf{.08}$

Power Rule: Let $4^x = 1.1$ then $2^{6x} = ?$

Solution: $4^x = 2^{2x} = 1.1 \Rightarrow 2^{6x} = (2^{2x})^3 = 1.1^3 = \mathbf{1.331}$

The following are some more problems about exponent rules:

Problem Set 3.1.6

1. $6^x = 34$, then $6^{x+2} =$
2. $3^x = 70.1$, then $3^{x+2} =$
3. $4^{x+1} = 2$, then $4^{x-1} =$
4. $6^x = 72$, then $6^{x-2} =$
5. $7^x = 14$, then $7^{x-2} =$
6. $4^x = .125$, then $4^{2x} =$
7. $8^x = 17$, then $8^{2x} =$
8. $2^x = 14.6$, then $2^{x+1} =$
9. $4^x = 32$, then $x =$
10. $9^x = 108$, then $3^{2x+1} =$
11. $6^{2x} = 36$, then $6^{3x} =$
12. $8^x = 256$, then $x =$
13. $27^x = 81$, $x =$
14. $2^8 \div 4^3$ has a remainder of:
15. $9^x = 27^{x+2}$, $x =$
16. $n^4 = 49$, then $n^6 =$
17. $16^x = 169$, then $4^x =$
18. $5^{3x} = 25^{2+x}$, $x =$
19. $n^6 = 1728$, then $n^4 =$
20. $4^x \div 16^x = 4^{-2}$, $x =$
21. $\sqrt{a^5} \times \sqrt[5]{a^2} = \sqrt[n]{a^{29}}$, $n =$
22. $6^8 \div 8$ has a remainder of:
23. $\sqrt[3]{a^4} + \sqrt[4]{a^3} = \sqrt[12]{a^n}$, $n =$

3.1.7 Log Rules

Logarithms are usually tested on the third and fourth columns of the test, however, if logarithm rules are fully understood these can be some of the simplest problems on the test. The following is a collection of log rules which are actively tested:

Definition:	$\log_a b = x$	$a^x = b$
Power Rule:	$\log_a b^n$	$n \log_a b$
Addition of Logs:	$\log_a b + \log_a c$	$\log_a (bc)$
Subtraction of Logs:	$\log_a b - \log_a c$	$\log_a \left(\frac{b}{c}\right)$
Change of Bases:	$\log_a b$	$\frac{\log b}{\log a}$

In the above table $\log_{10} a$ is represented as $\log a$. The following are some sample problems illustrating how each one of the rules might be tested:

Example: Find $\log_4 .0625$.

Solution: Applying the definition we know that $4^x = .0625 = \frac{1}{16}$. Therefore, our answer is $x = -2$

Example: Find $\log_8 16$.

Solution: Again, applying the definition, $8^x = 16$, which can be changed to $2^{3x} = 2^4 \Rightarrow x = \frac{4}{3}$.

Example: Find $\log_{12} 16 + \log_{12} 36 - \log_{12} 4$.

Solution: We know from the addition/subtraction of logs that the above expression can be written as $\log_{12} \frac{16 \cdot 36}{4} = \log_{12} 16 \cdot 9 = \log_{12} 144 \Rightarrow 12^x = 144 \Rightarrow x = 2$.

Example: Find $\log_5 8 \div \log_{25} 16$

Solution: These are probably the most challenging logarithm problems you will see on the exam. They involved changing bases and performing the power rule. Let's look at what happens when we change bases:

$$\log_5 8 \div \log_{25} 16 = \frac{\log 8}{\log 5} \div \frac{\log 16}{\log 25} = \frac{\log 2^3}{\log 5} \times \frac{\log 5^2}{\log 2^4} = \frac{3 \cdot \log 2}{\log 5} \times \frac{2 \cdot \log 5}{4 \cdot \log 2} = 3 \times \frac{1}{2} = \frac{3}{2}$$

In addition to the above problems, there are some approximations of logarithms which pop up. For those, there are some quantities which would be nice to have memorized to compute a more accurate approximations. Those are:

$$\log_{10} 2 \approx .3 \quad \log_{10} 5 \approx .7$$

$$\ln 2 \approx .7 \quad \ln 10 \approx 2.3$$

Where $\ln x = \log_e x$.

The following is example of how approximations of logs can be calculated:

$$200 \log 200 = 200 \log(2 \cdot 100) = 200 \cdot (\log 2 + \log 100) \approx 200 \cdot (.3 + 2) = \mathbf{460}$$

The following are some more practice problems:

Problem Set 3.1.7

- $-2\log_3 x = 4, x =$
- $\log_{12} 2 + \log_{12} 8 + \log_{12} 9 =$
- $\log_3 40 - \log_3 8 + \log_3 1.8 =$
- $\log_x 216 = 3, x =$
- $f(x) = \log_3 x - 4, f(3) =$
- $\log_8 16 =$
- $\log_3 x = 4, \sqrt{x} =$
- $\log_x 343 = 3, x =$
- If $\log .25 = 3$, then $\log 4 =$
- $(\log_5 6)(\log_6 5) =$
- $\log_3 216 \div \log_3 6 =$
- $\log_3 32 - \log_3 16 + \log_3 1.5 =$
- $\log_2 64 \div \log_2 4 =$
- $\log_4 32 + \log_4 2 - \log_4 16 =$
- $\log_5 625 \times \log_5 25 \div \log_5 125 =$
- $\log_4 8 \times \log_8 4 =$
- $\log_4 256 \div \log_4 16 \times \log_4 64 =$
- $\log_8 k = \frac{1}{3}, k =$
- $\log_5 M = 2, \sqrt{M} =$
- $4\log_9 k = 2, k =$
- $\log_4 8 = N$ then $2N =$
- $\log_9 3 = W$ then $3W =$
- $\log_k 32 = 5, k =$
- $\log_3[\log_2(\log_2 256)] =$
- $\log_4 .5 = k, k =$
- $\log_5[\log_4(\log_3 81)] =$
- $\log_{16} 8 = w, w =$
- $\log_9 k = 2.5, k =$
- $\log_2[\log_3(\log_2 512)] =$
- $\log_b .5 = -.5, b =$
- $\log_b 8 = 3, b =$
- $\log_3[\log_4(\log_5 625)] =$
- $\log_4 8 = k, k =$
- $\log_4[\log_3(\log_5 125)] =$
- $\log_4 .125 = k, k =$
- $\log_8(3x - 2) = 2, x =$
- $\log_4[\log_2(\log_6 36)] =$
- $\log_4 x = 3, \sqrt{x} =$
- $\log_5 x^2 = 4, \sqrt{x} =$

40. (*) $300 \log 600 =$

46. $\log_k 1728 = 3, k =$

41. $\log_4 x = -.5, x =$

47. $\log_4 x = 3, \sqrt{x} =$

42. $3 \log_2 x = 6, \sqrt{x} =$

48. $\log_2(\log_{10} 100) =$

43. $\log_2 x = 9, \sqrt[3]{x} =$

49. $\log_x 64 = 1.5, x =$

44. $\log_x 64 = 3, x^{-2} =$

50. $\log_8(\log_4 16) =$

45. $\log_9 x = 2, \sqrt{x} =$

51. $\log_9(\log_3 27) =$

3.1.8 Square Root Problems

A common question involves the multiplication of two square roots together to solve for (usually) an integer value. For example:

$$\begin{aligned}\sqrt{12} \times \sqrt{27} &= \sqrt{12} \times \sqrt{3} \times \sqrt{9} \\ &= \sqrt{36} \times \sqrt{9} \\ &= 6 \times 3 = \mathbf{18}\end{aligned}$$

Usually the best approach is to figure out what you can take away from one of the square roots and multiply the other one by it. So from the above example, notice that we can take a 3 away from the 37 to multiply the 12 with, leading to just $\sqrt{36} \times \sqrt{9}$ which are easy square roots to calculate. With this method, there are really no “tricks” involved, just a procedure that should be practiced in order to master it. The following are some more problems:

Problem Set 3.1.8

1. $\sqrt{75} \times \sqrt{27} =$

7. $\sqrt{44 \times 11} =$

2. $\sqrt{75} \times \sqrt{48} =$

8. $\sqrt{96 \times 24} =$

3. $\sqrt{44} \times \sqrt{99} =$

9. $\sqrt{72 \times 18} =$

4. $\sqrt{39} \times \sqrt{156} =$

10. $\sqrt{45} \div \sqrt{80} =$

5. $\sqrt{27} \times \sqrt{48} =$

11. $\sqrt{28} \div \sqrt{63} =$

6. $\sqrt{98 \times 8} =$

12. $\sqrt[3]{125 \times 512} =$

3.1.9 Finding Approximations of Square Roots

Seeing a problem like approximating $\sqrt{1234567}$ is very common in the middle of the test. The basic trick is you want to “take out” factors of 100 under the radical. Let’s look at the above example after noticing that we can roughly approximate (within the margin of error) $\sqrt{1234567} \approx \sqrt{1230000}$. Now:

$$\sqrt{1230000} = \sqrt{123 \cdot 100 \cdot 100} = 10 \cdot 10\sqrt{123}$$

Now we are left with a much simpler approximation of the $100 \cdot \sqrt{123} \approx 100 \cdot 11 = \mathbf{1100}$.

You can follow the same procedure for cubed roots as well, only you need to find factors of 1000 under the radical to take out. Let’s look at the example of $\sqrt[3]{1795953}$ after making the early approximation of $\sqrt[3]{1795953} \approx \sqrt[3]{1795000}$

$$\sqrt[3]{1795000} = \sqrt[3]{1795 \cdot 1000} = 10 \cdot \sqrt[3]{1795}$$

Well we should have memorized that $12^3 = 1728$ so we can form a rough approximation:

$$10 \cdot \sqrt[3]{1795} = 10 \cdot 12.1 = \mathbf{121}$$

So the trick is if you are approximating the n^{th} root of some number, you “factor out” sets of the n -digits and then approximate a much smaller value, then move the decimal place over accordingly.

Now in some instances you are asked to find the *exact* value of the cubed root. For example: $\sqrt[3]{830584}$. Now the procedure would be as followed:

1. Figure out how many digits you are going to have by noticing how many three-digit “sets” there are. Most will only be two digit numbers, however this is not guaranteed.
2. To find out the units digit, look at the units digit of the number given and think about what number cubed would give that result.
3. After that, you want to disregard the last three digits, and look at the remaining number and find out what number cubed is the first integer *less* than that value.

So to use the procedure give above for the problem of $\sqrt[3]{830584}$:

1. Well you have two, three-digit “sets” (the sets being 584 and 830). This means that we are looking for a two-digit number in our answer.
2. The last digit is 4, so what number cubed ends in a 4? The answer is that $4^3 = 64$ so the last digit of the answer is **4**.
3. Now we disregard the first set of three (584) and look at the remaining numbers (830). So what number cubed is *less* than 830. Well we know $10^3 = 1000$ and $9^3 = 729$ so **9** is the largest integer so that when cubed is less than 830. So that is the tens digit.
4. The answer is **94**.

The following are problems so that you can practice this procedure of finding approximate and exact values of square and cubed roots.

Problem Set 3.1.9

1. (*) $\sqrt{15376} =$
2. $\sqrt[3]{830584} =$
3. (*) $\sqrt{23456} =$
4. (*) $\sqrt{32905} =$
5. (*) $\sqrt{6543210} =$
6. $\sqrt[3]{658503} =$
7. (*) $\sqrt{6213457} =$
8. (*) $\sqrt{173468} =$
9. (*) $\sqrt{6420135} =$
10. (*) $\sqrt{872143} =$
11. (*) $\sqrt{272727} =$
12. (*) $\sqrt{38527} =$
13. (*) $\sqrt{32323} =$
14. (*) $\sqrt{18220} =$
15. (*) $\sqrt{25252} =$
16. (*) $\sqrt{265278} =$
17. (*) $\sqrt{81818} =$
18. (*) $\sqrt{262626} =$
19. (*) $\sqrt{765432} =$
20. (*) $\sqrt{80808} =$
21. (*) $\sqrt{97531} =$
22. (*) $\sqrt{86420} =$
23. (*) $\sqrt{8844} \times \sqrt{6633} =$
24. (*) $\sqrt[3]{217777} \times \sqrt{3777} \times 57 =$
25. (*) $\sqrt[3]{26789} \times \sqrt{911} \times 31 =$
26. (*) $\sqrt[3]{215346} \times \sqrt{3690} \times 57 =$
27. (*) $\sqrt[3]{2006} \times 6002 =$
28. (*) $\sqrt[3]{63489} \times \sqrt{1611} \times 41 =$
29. (*) $\sqrt[4]{14643} \times \sqrt[3]{1329} \times \sqrt{120} =$

3.1.10 Complex Numbers

The following is a review of Algebra-I concerning complex numbers. Recall that $i = \sqrt{-1}$. Here are important definitions concerning the imaginary number $a + bi$:

Complex Conjugate: $a - bi$

Complex Modulus: $\sqrt{a^2 + b^2}$

Complex Argument: $\arctan \frac{b}{a}$

The only questions that are usually asked on the number sense test is multiplying two complex numbers and rationalizing a complex number. Let's look at examples of both:

Multiplication: $(a + bi) \cdot (c + di) = (ac - bd) + (ad + bc)i$

Example: $(3 - 2i) \cdot (4 + i) = a + bi, a + b = ?$

Solution: $a = 3 \cdot 4 + 2 \cdot 1 = 14$ and $b = 3 \cdot 1 + (-2) \cdot 4 = -5$. So $a + b = 14 - 5 = \mathbf{9}$.

Rationalizing: $(a + bi)^{-1} = \frac{a - bi}{a^2 + b^2}$

Example: $(3 - 4i)^{-1} = a + bi, a - b = ?$

Solution: $(3 - 4i)^{-1} = \frac{3 + 4i}{3^2 + 4^2} \Rightarrow a = \frac{3}{25}$ and $b = \frac{4}{25}$. So $a - b = \frac{3}{25} - \frac{4}{25} = -\frac{1}{25}$.

The following are some more practice problems about Complex Numbers:

Problem Set 3.1.10

1. $(4 - i)^2 = a + bi, a =$

13. $(2 - 5i)(3 - 4i) = a + bi, a - b =$

2. $(6 - 5i)(6 + 5i) =$

14. $(4 - 3i)(2 - i) = a + bi, a - b =$

3. The conjugate of $(4i - 6)$ is $a + bi, a =$

15. $(2 + 7i)(2 - 7i) = a + bi, a - b =$

4. $(5 + i)^2 = a + bi, a =$

16. $(2 + 3i)(4 + 5i) = a + bi, a =$

5. $(9 - 3i)(3 + 9i) = a + bi, a =$

17. $(3 + 4i)^2 = a + bi, a =$

6. $(8 + 3i)(3 - 8i) = a + bi, a =$

18. The modulus of $14 + 48i$ is:

7. $(2 + 3i) \div (2i) = a + bi, a =$

19. $(2 - 5i)^2 = a + bi, a + b =$

8. $(3 - 4i)(3 + 4i) =$

20. $(5 + 4i)(3 + 2i) = a + bi, a =$

9. $(24 - 32i)(24 + 32i) =$

21. $(0 + 4i)^2 = a + bi, b =$

10. $(5 + 12i)^2 = a + bi, a + b =$

22. $(4 + 5i)(4 - 5i) =$

11. $(3 - 5i)(2 - 5i) = a + bi, a + b =$

23. The modulus of $(11 + 60i)^2$ is:

12. $(2 - 5i)(3 + 5i) = a + bi, a =$

24. $(0 - 3i)^5 = a + bi, b =$

25. $(3 - 5i)(2 + i) = a + bi, a + b =$

26. $(4 - 2i)(3 - i) = a + bi$, $a + b =$

27. $(1 + i)^9 =$

28. $(2 + 3i) \div (3 - 2i) = a + bi$, $b =$

29. $(2 - 3i) \div (3 - 2i) = a + bi$, $a =$

30. $(2i)^6 =$

31. $(3 + 4i) \div (5i) = a + bi$, $a + b =$

32. The modulus of $(24 + 7i)^2$ is:

33. $(3i - 2) \div (3i + 2) = a + bi$, $b =$

34. The modulus of $(5 + 12i)^2$ is:

3.1.11 Function Inverses

Usually on the last column you are guaranteed to have to compute the inverse of a function at a particular value. The easiest way to do this is to not *explicitly* solve for the inverse and plug in the point but rather, compute the inverse at that point as you go. For example if you are given a function $f(x) = \frac{3}{2}x - 2$ and you want to calculate $f^{-1}(x)$ at the point $x = 3$, you *don't* want to do the standard procedure for finding inverses (switch the x and y variables and solve for y) which would be:

$$x = \frac{3}{2}y - 2 \Rightarrow y = (x + 2) \cdot \frac{2}{3} \text{ at } x=3: \Rightarrow y = (3 + 2) \cdot \frac{2}{3} = \frac{10}{3}$$

Not only do you solve for the function, you have to remember the function while you're plugging in numbers. An easier way is just switch the x and y variables, then plug in the value for x , then compute y . That way you aren't solving for the inverse function for *all* points, but rather the inverse at that particular point. Let's see how doing that procedure would look like:

$$x = \frac{3}{2}y - 2 \Rightarrow 3 = \frac{3}{2}y - 2 \Rightarrow y = (3 + 2) \cdot \frac{2}{3} = \frac{10}{3}$$

Although this might not seem like much, it does help in saving some time.

Another important thing to remember when computing inverses is a special case when the function is in the form:

$$f(x) = \frac{ax + b}{cx + d} \Rightarrow f^{-1}(x) = \frac{-dx + b}{cx - a}$$

This was a very popular trick awhile back, but slowly it's appearance has been dwindling, however that does not mean a resurgence is unlikely. The important thing to remember is to line up the x 's on the numerator and denominator so it is in the require form. Here is an example problem to show you the trick:

$$\begin{aligned} \text{Find } f^{-1}(2) \text{ where } f(x) &= \frac{2x + 3}{4 + 5x}. \\ f(x) = \frac{2x + 3}{4 + 5x} &= \frac{2x + 3}{5x + 4} \Rightarrow f^{-1}(x) = \frac{-4x + 3}{5x - 2} \\ \text{So } f^{-1}(2) &= \frac{-4 \cdot 2 + 3}{5 \cdot 2 - 2} = \frac{-5}{8}. \end{aligned}$$

Here are some problems to give you some practice:

Problem Set 3.1.11

1. $f(x) = 3x + 2$, $f^{-1}(-2) =$
2. $f(x) = \frac{4x}{5}$, $f^{-1}(2) =$
3. $f(x) = 2 - 3x$, $f^{-1}(1) =$
4. $f(x) = x^2 - 1$ and $x > 0$, $f^{-1}(8) =$
5. $f(x) = 5 + 3x$, $f^{-1}(-2) =$
6. $f(x) = 4 - 3x$, $f^{-1}(2) =$
7. $f(x) = \frac{8}{3+x}$, $f^{-1}(2) =$
8. $f(x) = \frac{3-2x}{4}$, $f^{-1}(-1) =$
9. $f(x) = \frac{x^3}{3} + 3$, $f^{-1}(-6) =$
10. $\mathbf{f(x) = 2 - \frac{3x}{4}}$, $\mathbf{f^{-1}(5) =}$
11. $f(x) = 2x + 1$, $f^{-1}(3) =$
12. $g(x) = 3x + 2$, $g^{-1}(-1) =$
13. $h(x) = 2x - 3$, $h^{-1}(-1) =$
14. $f(x) = 2(x + 3)$, $f^{-1}(-4) =$
15. $f(x) = 2 - 3x$, $f^{-1}(4) =$
16. $h(x) = 5x - 3$, $h^{-1}(2) =$
17. $\mathbf{h(x) = 5 - 3x}$, $\mathbf{h^{-1}(-2) =}$
18. $f(x) = 2x + 2$, $f^{-1}(-2) =$
19. $f(x) = 3x - 3$, $f^{-1}(-3) =$
20. $f(x) = 4 - 4x$, $f^{-1}(-4) =$
21. $f(x) = \frac{3x-1}{x-3}$, $f^{-1}(1) =$
22. $\mathbf{f(x) = \frac{2x+1}{x-2}}$, $\mathbf{f^{-1}(3) =}$
23. $f(x) = \frac{3x-1}{x-3}$, $f^{-1}(-1) =$
24. $\mathbf{f(x) = \frac{1-3x}{x+3}}$, $\mathbf{f^{-1}(-2) =}$

3.1.12 Patterns

There is really no good trick to give you a quick answer to most pattern problems (especially the ones on the latter stages of the test). However, it is best to try to think of common things associated between the term number and the term itself. For example, you might want to keep in mind: squares, cubes, factorials, and fibonacci numbers. Let's look at some example problems:

Problem: Find the next term of 1, 5, 13, 25, 41, ...

Solution I: So for this, notice that you are adding to each term 4, 8, 12, 16 respectively. So each time you are incrementing the addition by 4 so, the next term will simply be 16 + 4 added to 41 which is **61**.

Solution II: Another way of looking at this is to notice that $1 = 1^2 + 0^2$, $5 = 2^2 + 1^2$, $13 = 3^2 + 2^2$, $25 = 4^2 + 3^2$, $41 = 5^2 + 4^2$, so the next term is equal to $6^2 + 5^2 = \mathbf{61}$

Problem: Find the next term of 0, 7, 26, 63, ...

Solution: For this one, notice that each term is one less than a cube: $0 = 1^3 - 1$, $7 = 2^3 - 1$, $26 = 3^3 - 1$, $63 = 4^3 - 1$, so the next term would be equal to $5^3 - 1 = \mathbf{124}$.

Here are some more problems to give you good practice with patterns:

Problem Set 3.1.12

1. Find the next term of 48, 32, 24, 20, 18, ...:
2. Find the next term of 1, 4, 11, 26, 57, ...:
3. Find the next term of 1, 8, 21, 40, ...:
4. Find the next term of 0, 1, 5, 14, 30, 55, ...:
5. **Find the next term of:**
2, 9, 28, 65, 126, ...:
6. The next term of 1, 2, 6, 24, 120, ... is:
7. The next term of 2, 2, 4, 6, 10, 16, ... is:
8. Find the 9th term of 1, 2, 4, 8, ...:
9. **Find the 10th term of:**
2, 6, 12, 20, 30, ...:
10. Find the 100th term of
2, 6, 10, 14, 18, ...:
11. The 10th term of 2, 5, 10, 17, 26 ... is:
12. The next term of 1, 4, 10, 19, 31, ... is:
13. The 8th term of 2, 9, 28, 65, 126, ... is:
14. The 8th term of 0, 7, 26, 63, 124, ... is:
15. The next term of 1, 5, 6, 11, 17, 28, ... is:
16. Find the next term of .0324, .054, .09, .15, ...:

3.1.13 Probability and Odds

Usually these problems involve applying the definitions of Odds and Probability which are:

$$\text{Probability} = \frac{\text{Desired Outcomes}}{\text{Total Outcomes}}$$

$$\text{Odds} = \frac{\text{Desired Outcomes}}{\text{Undesirable Outcomes}}$$

So the *probability* of rolling snake-eyes on a dice would be $\frac{1}{36}$ while the *odds* of doing this would be $\frac{1}{35}$. Usually the problems involving odds and probability on the number sense tests are relatively simple where desired outcomes can be computed by counting. The following are some practice problems so you can be familiar with the types of problems asked:

Problem Set 3.1.13

1. The odds of drawing a king from a 52-card deck is:
2. If 2 dice are tossed, what is the probability of getting a sum of 11:
3. A bag has a 3 red, 6 white, and 9 blue marbles. What is the probability of drawing a red one:
4. Three coins are tossed. Find the odds of getting 3 tails:

5. The odds of losing are 4-to-9. The probability of winning is:
6. The probability of winning is $\frac{5}{9}$.
The odds of losing is:
7. The odds of losing is $\frac{7}{13}$.
The probability of winning is:
8. If three dice are tossed once, what is the probability of getting three 5's:
9. If all of the letters in the words "NUMBER SENSE" are put in a box, what are the odds of drawing an 'E':
10. **The probability of success if $\frac{8}{17}$.
The odds of failure is:**
11. **If all of the letters in the words "STATE MEET" were put in a box, what is the probability of drawing an 'E':**
12. A pair of dice is thrown, the odds that the sum is a multiple of 5 is:
13. The probability of losing is $44\frac{4}{9}\%$.
The odds of winning is:
14. **The odds of winning the game is 3 to 5.
The probability of losing the game is:**
15. A number is drawn from $\{1, 2, 3, 6, 18\}$. The probability that the number drawn is not a prime number is:
16. The odds of drawing a red 7 from a standard 52-card deck is:
17. A number is randomly drawn from the set $\{1, 2, 3, 4, 5, 6, 7, 8, 9\}$. What are the odds that the number drawn is odd:
18. A number is drawn from the set $\{1, 2, 3, 4, 5\}$. What is the probability that the number drawn is a factor of 6:
19. The odds of randomly drawing a prime number from the set $\{1, 2, 3, 4, 5\}$ is:
20. When two dice are tossed, the probability that the sum of the faces will be 3 is:
21. A pair of dice is thrown. The probability that their sum is 7 is:
22. A pair of dice is thrown. The odds that their sum is 7 is:
23. A pair of dice is thrown. The odds that the sum is 6 or 8 is:
24. Two dice are tossed. What is the probability the sum is a multiple of 4:
25. **Two dice are tossed. What is the probability the sum is a multiple of 5:**
26. A die is rolled. What is the probability that a multiple of 2 is shown:
27. A die is rolled. What is the probability that a composite number is rolled:
28. A die is rolled. What is the probability that a factor of 12 is shown:
29. The probability of losing is 4-to-7.
What are the odds of winning:
30. A pair of dice are rolled. What are the odds that the same number is shown:
31. **The odds of drawing an ace followed by a king from a standard 52-card deck with replacement is:**

3.1.14 Sets

Questions concerning sets are by far the easiest problems on the number sense tests. The only topics that are actively questioned are the definitions of intersection, union, complement, and subsets. Let sets $A = \{M, E, N, T, A, L\}$ and $B = \{M, A, T, H\}$ then:

Intersection: The intersection between A and B (notated as $C = A \cap B$) is defined to be elements which are in *both* sets A and B. So in our case $C = A \cap B = \{M, A, T\}$ which consists of **3** elements.

Union: The union between A and B (notated as $D = A \cup B$) is defined to be a set which contains all elements in A and all elements in B. So $D = A \cup B = \{M, E, N, T, A, L, H\}$ which consists of **7** elements.

Complement Let's solely look at set A and define a new set $E = \{T, E, N\}$. Then the complement of E (notated a variety of ways, typically \bar{E} or E') with respect to Set A consists of simply all elements in A which aren't in E. So $\bar{E} = \{M, A, L\}$, which consists of **3** elements.

Subsets The number of possible subsets of a set is 2^n where n is the number of elements in the set. The number of *proper* subsets consists of all subsets which are strictly in the set. The result is that this disregards the subset of the set itself. So the number of proper subsets is $2^n - 1$. So in our example, the number of subsets of A is $2^7 = 128$ and the number of proper subsets is $2^7 - 1 = 127$. Another way to ask how many different subsets a particular set has is asking how many elements are in a set's *Power Set*. So the number of elements in the Power Set of B is simply $2^4 = 16$.

The following are questions concerning general set theory on the number sense test:

Problem Set 3.1.14

1. Set B has 15 proper subsets. How many elements are in B:
2. The number of subsets of $\{1, 3, 5, 7, 9\}$ is:
3. The number of elements in the power set of $\{M, A, T, H\}$ is:
4. If the power set for A contains 32 elements, then A contains how many elements:
5. The number of distinct elements of $[\{t, w, o\} \cup \{f, o, u, r\}] \cap \{e, i, g, h, t\}$ is:
6. The number of distinct elements of $\{m, a, t, h\} \cap \{e, m, a, t, i, c, s\}$ is:
7. The number of distinct elements of $[\{f, i, v, e\} \cap \{s, i, x\}] \cup \{t, e, n\}$ is:
8. If universal set $U = \{2, 3, 5, 7, 9, 11, 13, 17, 19\}$ and $A = \{3, 7, 13, 17\}$, then A' contains how many distinct elements:
9. If the universal set $U = \{n, u, m, b, e, r, s\}$ and set $A = \{s, u, m\}$ then the complement of set A contains how many distinct elements:
10. The universal set $U = \{n, u, m, b, e, r, s\}$, $A \subset U$ and $A = \{e, u\}$, then the complement of A contains how many elements:
11. The number of distinct elements in $[\{z, e, r, o\} \cap \{o, n, e\}] \cup \{t, w, o\}$ is:
12. The number of distinct elements in $[\{m, e, d, i, a, n\} \cap \{m, e, a, n\}] \cap \{m, o, d, e\}$ is:
13. The set $\{F, U, N\}$ has how many subsets:

14. The set $\{T, W, O\}$ has how many proper subsets:
15. Set A has 32 subsets. How many elements are in A :
16. The set P has 63 proper subsets. How many elements are in P :
17. Set A has 15 proper subsets. How many elements are in A :
18. **The set A has 8 distinct elements. How many proper subsets with at least one element does A have:**
19. Set $A = \{a, b, c, d\}$. How many proper subsets does set A have:
20. The number of proper subsets of $\{M, A, T, H\}$ is:
21. Set $A = \{o, p, q, r, s\}$ has how many improper subsets:

3.2 Changing Bases

3.2.1 Converting Integers

One of the topics I've found rather difficult teaching to students is the concept of changing bases. It seems that students have the concept of a base-10 system so ingrained in their mind (almost always unbeknownst to them) that it is difficult for them to consider other base systems. Hopefully this section will be a good introduction to the process of changing bases, and doing basic operations in other number systems. First, let's observe how we look at numbers in the usual base-10 fashion.

Everyone knows that 1294 means that you have one-thousand two-hundred and fifty-four of something, but expressing this in an unusual manner we can say:

$$1294 = 1 \cdot 1000 + 2 \cdot 100 + 5 \cdot 10 + 4 \cdot 1 = 1 \cdot 10^3 + 2 \cdot 10^2 + 9 \cdot 10^1 + 4 \cdot 10^0$$

From this we can see where this concept of "base-10" comes from, we are adding combinations of these powers of tens (depending on what 0 – 9 digit we multiply by). So, you can express any integer n in base-10 as:

$$n = a_m \cdot 10^m + a_{m-1}10^{m-1} + a_{m-2} \cdot 10^{m-2} + \cdots a_1 \cdot 10^1 + a_0 \cdot 10^0$$

Where all a_m 's are integers ranging from 0 – 9.

The fact that we are summing these various powers of 10 is completely an arbitrary one. We can easily change this to some other integer (like 6 for example) and develop a base-6 number system. Let's see what it would look like:

$$n = a_m \cdot 6^m + a_{m-1}6^{m-1} + a_{m-2} \cdot 6^{m-2} + \cdots a_1 \cdot 6^1 + a_0 \cdot 6^0$$

Where all a_m 's are integers ranging from 0 – 5.

So to use an example, let look at what the number 123_6 (where the subscript denotes we are in base-6) would look like in our usual base-10 system:

$$123_6 = 1 \cdot 6^2 + 2 \cdot 6^1 + 3 \cdot 6^0 = 1 \cdot 36 + 2 \cdot 6 + 3 \cdot 1 = 36 + 12 + 3 = \mathbf{51}_{10}$$

From this we have found the way to convert any base- n whole number to base-10!

Let's look at another example:

$$3321_4 = 3 \cdot 4^3 + 3 \cdot 4^2 + 2 \cdot 4^1 + 1 \cdot 4^0 = 3 \cdot 64 + 3 \cdot 16 + 2 \cdot 4 + 1 \cdot 1 = 192 + 48 + 8 + 1 = \mathbf{249}_{10}$$

So now that we know how to convert from base- n to base-10, let's look at the process on how to convert the opposite direction:

1. Find the highest power of n which divides the base-10 number (let's say it is the m^{th} power).
2. Figure out how many times it divides it and that will be your $(m + 1)^{th}$ digit in base- n .
3. Take the remainder and figure out how many times one less than the highest power divides it (so see how many times n^{m-1} divides it). That is your $(m)^{th}$ digit.
4. Take the remainder, and continue process.

I know that this might seem complicated, but let's look at an example we have already done in the "forward" direction to illustrate how to go "backwards." Convert 51_{10} to base-6:

1. Well we know $6^2 = 36$ and $6^3 = 216$, so the highest power which divides 51 is 6^2 .
2. 36 goes into 51 one time, so our 3^{rd} digit is 1.
3. The remainder when dividing 51 by 36 is 15.
4. Now we see how many times 6^1 goes into 15 (which is 2 times, so our 2^{nd} digit is 2).
5. The remainder when dividing 15 by 6 is 3.
6. $6^0 = 1$ divides 3 obviously 3 times, so our 1^{st} digit is 3
7. So after conversion, $51_{10} = \mathbf{123}_6$, which corresponds to what we expected.

As a warning, some digits might be zero when you do the base conversion. Let's look at an example of this: Convert 18_{10} to base-4:

$4^2 = 16$ and $4^3 = 64$, so $4^2 = 16$ goes into 18 once with a remainder of 2:	Third Digit is 1
Now $4^1 = 4$ doesn't go into 2:	Second Digit is 0
4^0 goes into 2 twice:	First Digit is 2
Answer:	102_4

This seems like a lot of steps in making a base conversion, but after substantial practice, it will become second nature. Here are some practice problems with just converting bases from base- n to base-10 and reverse.

Problem Set 3.2.1

- | | |
|---------------------------------|-------------------------------|
| 1. $212_5 = \text{-----}_{10}$ | 4. $3^4 + 3 = \text{-----}_9$ |
| 2. $108 = \text{-----}_4$ | 5. $2^4 + 2 = \text{-----}_4$ |
| 3. $2004_5 = \text{-----}_{10}$ | |

6. $82 = \text{-----}_5$
7. $4^3 + 4 = \text{-----}_4$
8. $24 = \text{-----}_2$
9. $3^3 + 3 = \text{-----}_3$
10. $48 = \text{-----}_3$
11. $4^3 + 2^3 = \text{-----}_8$
12. $2^4 + 1 = \text{-----}_8$
13. $200_{10} = \text{-----}_7$
14. $72 + 18 + 4 = \text{-----}_6$
15. $234_{10} = \text{-----}_5$
16. $123_4 = \text{-----}_{10}$
17. $2^5 + 2 = \text{-----}_4$
18. $430_{10} = \text{-----}_5$
19. $540_{10} = \text{-----}_6$
20. $243 + 27 + 3 = \text{-----}_9$
21. $200_5 = \text{-----}_{10}$
22. $200_6 = \text{-----}_{10}$
23. $4^4 + 4^2 + 4^0 = \text{-----}_4$
24. $3^3 + 3^2 + 3^0 = \text{-----}_3$
25. $216 + 108 + 30 + 5 = \text{-----}_6$
26. $44_b = 40$, then $b =$
27. $123_{10} = \text{-----}_5$
28. $123_4 = \text{-----}_5$
29. $8^2 + 2^4 + 4^0 = \text{-----}_4$
30. $234_5 = \text{-----}_4$
31. $686 + 98 + 14 = \text{-----}_7$
32. $77_{10} = \text{-----}_7$
33. $4^3 + 4 = \text{-----}_8$
34. $234_5 = \text{-----}_{10}$
35. $3^4 + 3^2 + 3^0 = \text{-----}_3$
36. $123_{10} = \text{-----}_4$
37. $125 + 75 + 15 + 1 = \text{-----}_5$
38. $234_{10} = \text{-----}_5$
39. $1728 + 288 + 36 + 4 = \text{-----}_{12}$
40. $128 + 48 + 12 + 2 = \text{-----}_4$
41. Find b when $4b_6 = 29$:
42. $45_6 = \text{-----}_9$
43. $210_4 = \text{-----}_6$
44. $43_8 = \text{-----}_9$
45. $34_5 = \text{-----}_7$

3.2.2 Converting Decimals

In the similar manner of how we analyzed an integer n in base-10, we can look at decimals in base-10 as well. For example, let's look at how we see .125 in base-10

$$.125 = 1 \cdot (.1) + 2 \cdot (.01) + 5 \cdot (.001) = 1 \cdot 10^{-1} + 2 \cdot 10^{-2} + 5 \cdot 10^{-3}$$

You can display this in terms of fractions as well:

$$= \frac{1}{10} + \frac{2}{100} + \frac{5}{1000} = \frac{1}{10} + \frac{1}{50} + \frac{1}{200} = \frac{20 + 4 + 1}{200} = \frac{1}{8}$$

Similar to the previous session, we can replace the powers of ten by the power of any fraction. Let's look at converting $.321_6$ to a base-10 fraction:

$$.321_6 = \frac{3}{6} + \frac{2}{36} + \frac{1}{216} = \frac{108 + 12 + 1}{216} = \frac{121}{216}$$

Going in the reverse direction is similar to what you do with integers. The following is a problem set to give you more practice:

Problem Set 3.2.2

1. Change $.32_5$ to a base-10 fraction:
2. Change $.34_5$ to a base-10 fraction:
3. **Change $.111_7$ to a base-10 fraction:**
4. Change $.33_4$ to a base-10 fraction:
5. **Change $.234_5$ to a base-10 fraction:**
6. Change $.44_8$ to a base-10 fraction:
7. Change $.33_6$ to a base-10 fraction:
8. Change $.66_{12}$ to a base-10 fraction:
9. Change $.202_5$ to a base-10 fraction:
10. Change $.55_6$ to a base-10 fraction:
11. **Change $.444_5$ to a base-10 fraction:**
12. Change $.44_5$ to a base-10 decimal:
13. Change $.14_5$ to a base-10 decimal:
14. Change $\frac{9}{16}$ to a base-4 decimal:
15. Change $\frac{35}{36}$ to a base-6 decimal:
16. Change $\frac{15}{16}$ to a base-4 decimal:

17. Change $\frac{15}{16}$ to a base-8 decimal:

decimal:

18. Change $\frac{11}{25}$ to a base-5 decimal:

19. Change $\frac{30}{49}$ to a base-7 decimal:

3.2.3 Performing Operations

For some basic operations in other bases, sometimes it is simpler to convert all numbers to base-10, perform the operations, then convert back to base- n . Let's look at an example where I would employ this technique:

$$23_4 \times 3_4 + 12_4 = 11 \times 3 + 6 = 39 = 213_4$$

However, when numbers are larger, this might not be the case, so let's look at the most popular operations on the number sense test which are addition (and subsequently subtraction) and multiplication (division is usually not tested, so I will exclude explaining this operation).

Addition:

For addition of two integers in base-10 we sum the digits one at a time writing down the answer digit (0–9) and carrying when necessary. Other base- n work in the same manner with the only difference being the answer digits range from 0 to $(n - 1)$. Let's look at an example:

$124_6 + 53_6 =$	First Digit:	$4_6 + 3_6$	11_6
	Second Digit:	$5_6 + 2_6 + 1_6$	12_6
	Third Digit:	$1_6 + 1_6$	2_6
	Answer:		221_6

Subtraction:

Subtraction works in a similar method, only the concept of “borrowing” when you can't subtract the digits is slightly altered. When you “borrow” in base-10 you lower the digit you are borrowing from and then “add” 10 to the adjacent digit to aid in the subtraction. In a different base- n , you will be borrowing in the same fashion but adding n to the adjacent digit. Let's look at an example:

$122_4 - 13_4 =$	First Digit:	Since you “can't” do $2 - 3$ you have to borrow	
		$(4_4 + 2_4) - 3_4$	3_4
	Second Digit:	$(2_4 - 1_4) - 1_4$	0_4
	Third Digit:	1_4	1_4
	Answer:		103_4

In the above expressions, everything in italics represents the borrowing process. When borrowing from

the second digit, you lower it by 1 (seen by the -1_4) and then add to the adjacent digit (the first one) 4_4 .

Multiplication: What I like to do for multiplication in a difference base is essentially perform the FOILING procedure in base-10 then convert your appropriate result to base- n and apply appropriate carry rules. Let's look at a couple of examples (one involving carries and the other one not):

	First Digit:	$1 \times 3 = 3_{10}$	3_9
	Second Digit:	$1 \times 1 + 2 \times 3 = 7_{10}$	7_9
$13_9 \times 21_9 =$	Third Digit:	$2 \times 1 = 2_{10}$	2_9
	Answer:		273_9

The above scenario was simple because no carries were involved and converting those particular single digits from base-10 to base-9 was rather simple. Let's look at one with carries involved:

	First Digit:	$3 \times 5 = 15_{10}$	16_9
	Second Digit:	$3 \times 4 + 2 \times 5 + 1 = 23_{10}$	25_9
$45_9 \times 23_9 =$	Third Digit:	$2 \times 4 + 2 = 10_{10}$	11_9
	Fourth Digit:	1	1
	Answer:		1156_9

The above example shows the procedure where you do the FOILING in base-10 then convert that to base-9, write down last digit, carry any remaining digits, repeat procedure. As one can see to perform multiplication in other bases it is important to have changing bases automatic so that the procedure is relatively painless.

To practice the above three operations here are some problems:

Problem Set 3.2.3

- | | |
|---------------------------------------|---------------------------------------|
| 1. $112_6 + 4_6 = \text{-----}_6$ | 6. $37_8 + 56_8 = \text{-----}_8$ |
| 2. $53_6 \times 4_6 = \text{-----}_6$ | 7. $88_9 + 82_9 = \text{-----}_9$ |
| 3. $101_2 - 11_2 = \text{-----}_2$ | 8. $100_6 - 44_6 = \text{-----}_6$ |
| 4. $44_5 \times 4_5 = \text{-----}_5$ | 9. $104_8 - 47_8 = \text{-----}_8$ |
| 5. $26_9 \div 6_9 = \text{-----}_9$ | 10. $143_5 \div 4_5 = \text{-----}_5$ |

11. $22_9 - 66_9 = \text{-----}_9$

25. $123_4 \div 3_4 = \text{-----}_4$

12. $135_7 \times 4_7 = \text{-----}_7$

26. $431_5 \div 4_5 = \text{-----}_5$

13. $132_4 - 33_4 = \text{-----}_4$

27. $222_3 \times 2_3 = \text{-----}_3$

14. $42_5 - 34_5 + 23_5 = \text{-----}_5$

28. $(21_5 - 12_5) \times 11_5 = \text{-----}_5$

15. $123_5 \times 4_5 = \text{-----}_5$

29. $(33_4 + 22_4) \times 11_4 = \text{-----}_4$

16. $33_4 \times 3_4 - 21_4 = \text{-----}_4$

30. $235_6 \div 5_6 = \text{-----}_6$

17. $22_7 \times 4_7 = \text{-----}_7$

31. $543_7 \div 6_7 = \text{-----}_7$

18. $33_6 \times 3_6 = \text{-----}_6$

32. $234_5 + 432_5 = \text{-----}_5$

19. $22_6 + 33_6 + 44_6 = \text{-----}_6$

33. $33_4 \times 2_4 - 11_4 = \text{-----}_4$

20. $44_8 \times 4_8 = \text{-----}_8$

34. $44_5 \times 2_5 + 33_5 = \text{-----}_5$

21. $32_6 \div 5_6 \times 4_6 = \text{-----}_6$

35. $(13_5 + 12_5) \times 11_5 = \text{-----}_5$

22. $24_7 \div 6_7 + 24_7 = \text{-----}_7$

36. $11_4 \times 21_4 - 3_4 = \text{-----}_4$

23. $23_6 + 45_6 - 50_6 = \text{-----}_6$

37. $12_5 + 23_5 + 34_5 = \text{-----}_5$

24. $23_5 \times 4_5 - 10_5 = \text{-----}_5$

38. $(22_4 + 33_4) \times 11_4 = \text{-----}_4$

3.2.4 Changing Between Bases: Special Case

When changing between two bases m and n , the standard procedure is to convert the number from base- m to base-10 then convert that into base- n . However, there are special cases when the middle conversion into base-10 is unnecessary: when n is an integral power of m (say $n = m^a$, a an integer) or vice versa. The procedure is relatively simple, take the digits of m in groups of a and convert each group into base- n . For example, if we are converting 1001001_2 into base-4, you would take 1001001 in groups of two (since $2^2 = 4$) and converting each group into base-4. Let's see how it would look:

Convert 1001001_2 to base-4	First Digit:	01_2	1_4
	Second Digit:	10_2	2_4
	Third Digit:	00_2	0_4
	Fourth Digit:	1_2	1_4
	Answer:		1021_4

Let's look at an example where the converting base is that of the original base cubed (so you would take it in groups of 3):

Convert 110001011_2 to base-8	First Digit:	011_2	3_8
	Second Digit:	001_2	1_8
	Third Digit:	110_2	6_8
	Answer:		613_8

Similarly, you can perform the method in reverse. So when converting from base-9 to base-3 you would take each digit in base-9 and convert it to two-digit base-3 representation. For example:

Convert 643_9 to base-3	First/Second Digits:	3_9	10_3
	Third/Fourth Digits:	4_9	11_3
	Fifth/Sixth Digits:	6_9	20_3
	Answer:		201110_3

Problem Set 3.2.4

- | | |
|---------------------------------|--------------------------------|
| 1. $46_9 = \text{-----}_3$ | 7. $123_4 = \text{-----}_2$ |
| 2. $48_9 = \text{-----}_3$ | 8. $101011_2 = \text{-----}_4$ |
| 3. $1011011_2 = \text{-----}_8$ | 9. $231_4 = \text{-----}_2$ |
| 4. $123_4 = \text{-----}_2$ | 10. $432_8 = \text{-----}_2$ |
| 5. $2122_3 = \text{-----}_9$ | 11. $312_4 = \text{-----}_2$ |
| 6. $345_8 = \text{-----}_2$ | 12. $1111_2 = \text{-----}_4$ |

13. $1011_2 = \text{-----}_4$

15. $11011_2 = \text{-----}_4$

14. $123_4 = \text{-----}_2$

3.2.5 Changing Bases: Sum of Powers

When asked the sum of a series of powers of two ($1 + 2 + 4 + 8 + \dots + 2^n$), it is best to represent the number in binary, then we can see the result. For example purposes let's look at the sum $1 + 2 + 4 + 8 + 16 + 32 + 64$. If we represented them as binary it would be:

$$1 + 2 + 4 + 8 + 16 + 32 + 64 = 1 \cdot 2^0 + 1 \cdot 2^1 + 1 \cdot 2^2 + 1 \cdot 2^3 + 1 \cdot 2^4 + 1 \cdot 2^5 + 1 \cdot 2^6 = 1111111_2$$

$$1111111_2 = 1000000_2 - 1_2 \Rightarrow 2^7 - 1 = 128 - 1 = \mathbf{127}$$

Although this method is easiest with binary, you can apply it to other powers as well, as long as you are carefully. For example:

$$2 + 2 \cdot 3 + 2 \cdot 9 + 2 \cdot 27 + 2 \cdot 81 + 2 \cdot 243 = 2 \cdot 3^0 + 2 \cdot 3^1 + 2 \cdot 3^2 + 2 \cdot 3^3 + 2 \cdot 3^4 + 2 \cdot 3^5 = 222222_3$$

$$222222_3 = 1000000_3 - 1 = 3^6 - 1 = \mathbf{728}$$

3.2.6 Changing Bases: Miscellaneous Topics

There are a handful of topics involving changing bases that rely on understanding other tricks previously discussed in this book. Take this problem for example:

Problem: Convert the decimal $.333\dots_7$ into a base-10 fraction.

Solution: The above problem relies on using the formula for the sum of an infinite geometric series:

$$.333\dots_7 = \frac{3}{7} + \frac{3}{49} + \frac{3}{343} + \dots = \frac{\frac{3}{7}}{1 - \frac{1}{7}} = \frac{3}{7} \times \frac{7}{6} = \frac{1}{2}$$

Another problem which relies on understanding of how the derivation of finding the remainder of a number when dividing by 9, only in a different base is:

Problem: The number $123456_7 \div 6$ has what remainder?

Solution: The origins of this is rooted in modular arithmetic (see that section) and noticing that: $7^n \cong 1 \pmod{6}$. So our integer can be represented as:

$$123456_7 = 1 \cdot 7^5 + 2 \cdot 7^4 + 3 \cdot 7^3 + 4 \cdot 7^2 + 5 \cdot 7^1 + 6 \cdot 7^0 \cong (1 + 2 + 3 + 4 + 5 + 6) = \frac{6 \cdot 7}{2} = 21 \cong \mathbf{3} \pmod{6}$$

So an important result is that when you have a base- n number and divide it by $n - 1$, all you need to do is sum the digits and see what the remainder *that is* when dividing by $n - 1$.

Problem Set 3.2.6

1. $.555\dots_7 = \text{-----}_{10}$

2. The remainder when 123456_7 is divided by 6 is:

4. $.777\dots_9 = \text{-----}_{10}$

3. $.666\dots_8 = \text{-----}_{10}$

5. $.111\dots_5 = \text{-----}_{10}$

3.3 Repeating Decimals

The following sections are concerned with expressing repeating decimals as fractions. All of the problems of this nature have their root in sum of infinite geometric series.

3.3.1 In the form: $.aaaa\dots$

Any decimal in the form $.aaaa\dots$ can be re written as:

$$.aaaa\dots = \frac{a}{10} + \frac{a}{100} + \frac{a}{1000} + \dots$$

Which we can sum appropriately using the sum of an infinite geometric sequence with the common difference being $\frac{1}{10}$ (See Sum of Series Section):

$$\frac{a}{10} + \frac{a}{100} + \frac{a}{1000} + \dots = \frac{\frac{a}{10}}{1 - \frac{1}{10}} = \frac{a}{10} \times \frac{10}{9} = \frac{\mathbf{a}}{\mathbf{9}}$$

Which is what we expected knowing what the fractions of $\frac{1}{9}$ are. For example:

$$.44444\dots = \frac{\mathbf{4}}{\mathbf{9}}$$

3.3.2 In the form: $.ababa\dots$

In a similar vein, fractions in the form $.ababab\dots$ can be treated as:

$$.ababab\dots = \frac{ab}{100} + \frac{ab}{10000} + \frac{ab}{1000000} + \dots = \frac{\frac{ab}{100}}{1 - \frac{1}{100}} = \frac{ab}{100} \times \frac{100}{99} = \frac{\mathbf{ab}}{\mathbf{99}}$$

Where ab represents the digits (not $a \times b$). Here is an example:

$$.242424\dots = \frac{24}{99} = \frac{\mathbf{8}}{\mathbf{33}}$$

You can extend the concept for any sort of continuously repeating fractions. For example, $.abcabcabc\dots = \frac{abc}{999}$, and so on.

Here are some practice problems to help you out:

Problem Set 3.3.2

1. $.272727\dots =$

7. $.727272\dots =$

2. $.414141\dots =$

8. $.151515\dots =$

3. $.212121\dots =$

9. $.308308\dots =$

4. $.818181\dots =$

10. $.231231\dots =$

5. $.363636\dots =$

11. $.303303\dots =$

6. $.020202\dots =$

12. $.099099099\dots =$

3.3.3 In the form: .abbbb...

Fractions in the form $.abbbb\dots$ are treated in a similar manner (sum of an infinite series) with the inclusion of one other term (the $.a$ term). Let's see how it would look:

$$.abbb\dots = \frac{a}{10} + \frac{b}{100} + \frac{b}{1000} + \dots = \frac{a}{10} + \frac{\frac{b}{100}}{1 - \frac{1}{10}} = \frac{a}{10} + \frac{b}{90}$$

However we can continue and rewrite the fraction as:

$$\frac{a}{10} + \frac{b}{90} = \frac{9 \cdot a + b}{90} = \frac{(10 \cdot a + b) - a}{90}$$

Lets take a step back to see what this means. The numerator is composed of the sum $(10 \cdot a + b)$ which represents the two-digit number ab . Then you subtract from that the non-repeating digit and place that result over 90. Here is an example to show the process:

$$.27777\dots = \frac{27 - 2}{90} = \frac{25}{90} = \frac{5}{18}$$

Here are some more problems to give you more practice:

Problem Set 3.3.3

1. $.23333\dots =$

4. $.32222\dots =$

2. $.32222\dots =$

5. $.01222\dots =$

3. $.21111\dots =$

3.3.4 In the form: .abcabc...

Again, you can repeat the process above for variances. In this example we can represent .abcabc... can be represented in fraction form:

$$.abcabc... = \frac{abc - a}{990}$$

Where the abc represents the three-digit number abc (not the product $a \cdot b \cdot c$). Here is an example:

$$.437373737... = \frac{437 - 4}{990} = \frac{433}{990}$$

It is important for the number sense test to reduce all fractions. This can sometimes be the tricky part. The easiest way to check for reducibility is to see if you can divide the numerator by 2, 3, or 5. In the above example, 433 is not divisible by 2, 3, 5 so the fraction is in its lowest form.

Here is an example where you can reduce the fraction:

$$.2474747... = \frac{247 - 2}{990} = \frac{245}{990} = \frac{49}{198}$$

Problem Set 3.3.4

1. $.2131313... =$
2. $.1232323... =$
3. $.2313131... =$
4. $.3050505... =$
5. $.2050505... =$
6. $.3141414... =$
7. $.2717171... =$
8. $.2353535... =$
9. $.0474747... =$
10. $.2141414... =$
11. $.1232323... =$

3.4 Modular Arithmetic

A lot has been made about the uses of modular arithmetic (for example all of the sections dealing with finding remainders when dividing by 3, 9, 11, etc...). Here is a basic understanding of what is going on with modular arithmetic.

When dividing two numbers a and b results in a quotient q and a remainder of r we say that $a \div b = q + \frac{r}{b}$. With modular arithmetic, we are only concerned with the remainder so the expression of $a \div b = q + \frac{r}{b} \Rightarrow a \cong r(\text{mod } b)$.

So you know $37 \div 4$ has a remainder of 1, so we say $37 \cong 1(\text{mod } 4)$. As noted before, what's great about modular arithmetic is you can do the algebra of remainders (See: Remainders of Expressions Section). From this rule alone is where all of our divisibility rules come from. For example, let's see where we get our divisibility by 9 rule:

Recall we can express any base-10 number n by: $n = a_m 10^m + a_{m-1} 10^{m-1} + \cdots + a_1 10^1 + a_0 10^0$

So when we are trying to see the remainder when dividing by 9, we want to find what x is in the expression

$$n \cong x \pmod{9}.$$

However we do know that $10 \cong 1 \pmod{9}$, meaning $10^a \cong 1 \pmod{9}$ for all $a \geq 0$. So:

$$n = a_m 10^m + a_{m-1} 10^{m-1} + \cdots + a_1 10^1 + a_0 10^0 \cong (a_m + a_{m-1} + \cdots + a_1 + a_0) \pmod{9}$$

. Well $a_m + a_{m-1} + \cdots + a_1 + a_0$ is just the sum of the digits, so we just proved that in order for a number n to be divisible by 9 then the sum of it's digits have to be divisible by 9!

Learning the basics in modular arithmetic is not only crucial for recognizing and forming divisibility rules but also they pop up as questions on the number sense test. For example:

$$\text{Find } x, 0 \leq x \leq 4, \text{ if } x + 3 \cong 9 \pmod{5}$$

Here we know that $9 \cong 4 \pmod{5}$, so the problem reduces to finding x restricted to $0 \leq x \leq 4$ such that $x + 3 \cong 4 \pmod{5}$, which simply makes $x = 1$.

The following are some more problems to get you some practice on modular arithmetic:

Problem Set 3.4

1. $x + 6 \cong 9 \pmod{7}$, $0 \leq x \leq 6$
 $x =$

8. $x + 4 \cong 1 \pmod{8}$, $0 \leq x \leq 7$
 $x =$

2. $4^7 \div 7$ has a remainder of:

9. $3^8 \div 7$ has a remainder of:

3. $2^5 \times 3^5 \div 5$ has a remainder of:

10. $3x \cong 17 \pmod{5}$, $0 \leq x \leq 5$
 $x =$

4. $2^6 \times 3^4 \div 5$ has a remainder of:

11. $3x - 2 \cong 4 \pmod{7}$, $0 \leq x \leq 7$
 $x =$

5. $8^7 \div 6$ has a remainder of:

6. **If N is a positive integer and $4N \div 5$ has a remainder of 2 then $N \div 5$ has a remainder of:**

12. $6^8 \div 7$ has a remainder of:

13. $3^7 \div 7$ has a remainder of:

7. $x + 3 \cong 9 \pmod{5}$, $0 \leq x \leq 4$
 $x =$

14. $5^4 \div 11$ has a remainder of:

3.5 Fun with Factorials!

All of these problems incorporate common factorial problems.

3.5.1 $1 \cdot 1! + 2 \cdot 2! + \cdots + n \cdot n!$

The sum of $1 \cdot 1! + 2 \cdot 2! + \cdots + n \cdot n!$ is a fairly simple problem if you know the formula (its derivation is left to the reader).

$$1 \cdot 1! + 2 \cdot 2! + \cdots + n \cdot n! = (n + 1)! - 1$$

The simplest case would be to compute sums like:

$$1 \cdot 1! + 2 \cdot 2! + 3 \cdot 3! + 4 \cdot 4! = (4 + 1)! - 1 = 120 - 1 = \mathbf{119}$$

There are slight variations which could be asked (the easiest of which would be leaving out some terms).

$$1 \cdot 1! + 3 \cdot 3! + 5 \cdot 5! = (5 + 1)! - 1 - 2 \cdot 2! - 4 \cdot 4! = 720 - 1 - 4 - 96 = \mathbf{619}$$

The following are some practice problems:

Problem Set 3.5.1

1. $1 \cdot 1! + 2 \cdot 2! + 3 \cdot 3! + 4 \cdot 4! + 5 \cdot 5! =$

4. $1 \cdot 1! - 2 \cdot 2! - 3 \cdot 3! - 4 \cdot 4! =$

2. $1 \cdot 1! + 2 \cdot 2! + \cdots + 6 \cdot 6! =$

5. $2 \cdot 1! + 3 \cdot 2! + 4 \cdot 3! + 5 \cdot 4! =$

3. $1 \cdot 1! + 2 \cdot 2! + \cdots + 7 \cdot 7! =$

3.5.2 $\frac{a! \pm b!}{c!}$

This problem has pretty much nothing to do with factorials and mostly with basic fraction simplification. Take the following example:

$$\frac{8! + 6!}{7!} = \frac{8!}{7!} + \frac{6!}{7!} = 8\frac{1}{7}$$

Sometimes it is easier to just factor out the common factorial, for example:

$$\frac{3! + 4! - 5!}{3!} = \frac{3! \cdot (1 + 4 - 5 \cdot 4)}{3!} = 1 + 4 - 20 = \mathbf{-15}$$

Problem Set 3.5.2

1. $\frac{8! + 6!}{7!} =$

4. $\frac{11! - 9!}{10!} =$

2. $\frac{10! + 8!}{9!} =$

5. $\frac{10! - 11!}{9!} =$

3. $\frac{7! - 5!}{6!} =$

6. $6 \cdot 5 \cdot 4! - 5! =$

$$7. (2! + 3!) \div 5! =$$

$$8. (2! \times 3!) - 4! =$$

$$9. 7! \div 6! - 5! =$$

$$10. 7 \times 5! - 6! =$$

$$11. 2! - 3! \times 5! =$$

$$12. 8! \div 6! - 4! =$$

$$13. \frac{5! \cdot 4!}{6!} =$$

$$14. \frac{4 \times 5! - 5 \times 4!}{4!} =$$

$$15. \frac{4 \times 5! + 5 \times 4!}{4!} =$$

$$16. \frac{6 \times 7! - 7 \times 6!}{6!} =$$

$$17. \frac{10 \times 9! - 10! \times 9}{9!} =$$

$$18. \frac{8! \times 7 - 8 \times 7!}{7!} =$$

$$19. \frac{11 \times 10! - 11! \times 10}{11!} =$$

$$20. 6! \div (3! \times 2!) =$$

3.5.3 Wilson's Theorem

I've seen a couple of questions in the latter stages of the number sense question which asks something along the lines of:

$$6! \cong x \pmod{7}, 0 \leq x \leq 6, x = ?$$

Questions like this use the result from Wilson's Theorem which states:

$$\text{For prime } p, (p-1)! \cong (p-1) \pmod{p}$$

So using the above Theorem, we know that $6! \cong x \pmod{7}$, $0 \leq x \leq 6$, $x = \mathbf{6}$.

It is essentially for p to be prime Wilson's Theorem to be applicable. Usually, with non-prime factorial problems, you can lump common factors and then can check divisibility. For example:

$$4! \cong x \pmod{6}, 0 \leq x \leq 5, x = ?$$

Well we know that $4! = 4 \cdot 3 \cdot 2 \cdot 1 = 4 \cdot 6 \cong 0 \pmod{6} \Rightarrow x = \mathbf{0}$.

The following are some more problems to give you some practice:

Problem Set 3.5.3

$$1. (4!)(3!)(2!) \cong x \pmod{8}, 0 \leq x \leq 7.$$

$$x =$$

$$x =$$

$$2. (4+2)! \cong x \pmod{7}, 0 \leq x \leq 6.$$

$$x =$$

$$4. \frac{5! \cdot 3!}{4!} \cong k \pmod{8}, 0 \leq k \leq 7.$$

$$k =$$

$$3. (5-2)! \cong x \pmod{5}, 0 \leq x \leq 5.$$

$$5. \frac{5! \cdot 4!}{3!} \cong k \pmod{9}, 0 \leq k \leq 8.$$

$k =$

$$6. \mathbf{5! \cdot 3! \cong k \pmod{8}, 0 \leq k \leq 7.}$$

$k =$

3.6 Basic Calculus

If you are one of the fortunate people to reach the end of the fourth column, you will experience usually two or three calculus related problems which are relatively simple if you know the basics of calculus. If you haven't had basic calculus preparation, the following is a rough introduction on the computations of limits, derivatives, and integrals associated with the number sense test.

3.6.1 Limits

Usually the limits are the simplest kind where simple substitution can be used to get an appropriate answer. For example:

$$\lim_{x \rightarrow 3} 3x^2 - 4 = 3 \cdot 3^2 - 4 = \mathbf{23}$$

However certain problems, which when passing the limit, might lead to a $\frac{0}{0}$ violation. In this case, you want to see if there are any common factors that you can cancel so that passing the limit *doesn't* give you an indeterminate form. Let's look at an example:

$$\lim_{x \rightarrow 2} \frac{(x-2)(x+3)}{(x+5)(x-2)} = \lim_{x \rightarrow 2} \frac{(x+3)}{(x+5)} = \frac{\mathbf{5}}{\mathbf{7}}$$

If we had plugged in $x = 2$ into the original problem, we would have gotten a $\frac{0}{0}$ form, however after canceling the factors, we were able to pass the limit.

The final testable question concerning limits involve l'hôpital's rule (this requires the understanding of derivatives in order to compute it, see the next section for instructions on how to compute that). L'hôpital's rule states that if you come across a limit that gives an indeterminate form (either $\frac{0}{0}$ or $\frac{\infty}{\infty}$) you can compute the limit by taking the derivative of both the numerator and the denominator then passing the limit. So:

$$\lim_{x \rightarrow n} \frac{f}{g} = \frac{0}{0} \text{ or } \frac{\infty}{\infty} \Rightarrow \lim_{x \rightarrow n} \frac{f}{g} = \lim_{x \rightarrow n} \frac{f'}{g'}$$

Let's look at an example of l'hôpital's rule with computing the limit $\lim_{x \rightarrow 0} \frac{\sin x}{x}$:

$$\lim_{x \rightarrow 0} \frac{\sin x}{x} = \frac{0}{0} \Rightarrow \lim_{x \rightarrow 0} \frac{\sin x}{x} = \lim_{x \rightarrow 0} \frac{(\sin x)'}{x'} = \lim_{x \rightarrow 0} \frac{\cos x}{1} = \mathbf{1}$$

The following are some more practice problems with Limits:

Problem Set 3.6.1

1. $\lim_{x \rightarrow \infty} \frac{3x+8}{7x-4} =$

2. $\lim_{x \rightarrow 4} 2x - 6 =$

$$3. \lim_{x \rightarrow 2} \frac{x^2 - 4}{x - 2} =$$

$$6. \lim_{x \rightarrow 3} \frac{x^3 - 27}{x - 3} =$$

$$4. \lim_{x \rightarrow 2} \frac{x^3 - 8}{x^2 - 4} =$$

$$7. \lim_{x \rightarrow 0} \frac{x^2 - 3x}{x} =$$

$$5. \lim_{x \rightarrow \infty} \frac{3x - 1}{x} =$$

$$8. \lim_{x \rightarrow 3} \frac{x^3 - 27}{x^2 - 9} =$$

3.6.2 Derivatives

Usually on the number sense test, there is guaranteed to be a derivative (or double derivative) of a polynomial. Almost every single time, the use of the power rule is all that is required, so let's see how we can take the derivative of a polynomial:

$$\text{Define } f(x) = a_n x^n + a_{n-1} x^{n-1} + \dots + a_1 x^1 + a_0 x^0$$

Then

$$f'(x) = a_n(n)x^{n-1} + a_{n-1}(n-1)x^{n-2} + \dots + a_1(1)x^0$$

So the procedure is you multiply the coefficient by the power and then lower the power (notice that a constant after differentiating disappears). Let's look at an example:

$$\text{Let } f(x) = x^3 - 3x^2 + x - 3, \text{ solve for } f'(2).$$

$$f'(x) = 1 \cdot 3x^2 - 3 \cdot 2x + 1 \Rightarrow f'(2) = 1 \cdot 3 \cdot 2^2 - 3 \cdot 2 \cdot 2 + 1 = \mathbf{1}$$

When approached with taking double derivatives ($f''(x)$), then just follow the procedure twice:

$$\text{Let } f(x) = 5x^3 + 3x^2 - 7, \text{ solve for } f''(1).$$

$$f'(x) = 5 \cdot 3x^2 + 3 \cdot 2x = 15x^2 + 6x$$

$$f''(x) = 15 \cdot 2x + 6 \Rightarrow f''(1) = 30 \cdot 1 + 6 = \mathbf{36}$$

In the off chance that the derivative of sine/cosine or $e^x/\ln x$ is needed (like for using l'hôpitals rule), here is a chart showing these functions and their derivatives:

Function	Derivative
$\sin x$	$\cos x$
$\cos x$	$-\sin x$
e^x	e^x
$\ln x$	$\frac{1}{x}$

For more derivative rules, consult a calculus textbook (it would be good to be familiar with more derivative rules for the math test, but unlikely those rules will be applied to the number sense test).

Here are some problems to practice taking derivatives:

Problem Set 3.6.2

$$1. f(x) = 3x^2 + x - 5, f'(-2) =$$

$$2. f(x) = x^2 - 2x + 22, f'(2) =$$

$$3. f(x) = x^3 - 3x + 3, f''(2) =$$

$$4. g(x) = 2x^2 - 3x + 1, g'(2) =$$

$$5. f(x) = 3x^3 - 3x + 3, f'(-3) =$$

$$6. f(x) = 4x^3 + 2x^2, f''(-5) =$$

$$7. f(x) = x^3 - 3x + 3, f'(3) =$$

$$8. f(x) = x^4 - 4x + 4, f'(4) =$$

$$9. f(x) = 3x^2 + 4x - 5, f'(-6) =$$

$$10. f(x) = 2x^3 - 3x^4, f''(-1) =$$

$$11. f(x) = 4x^3 - 3x^2 + 1, f'(-1) =$$

$$12. f(x) = x^2 - 3x + 4, f''(-1) =$$

$$13. f(x) = 3x + 5x^2 - 7x^4, f'(1) =$$

$$14. f(x) = 3x^3 - 2x^2 + x, f''(1) =$$

$$15. f(x) = 2x^3 - 4x^2 + 6x, f'(1) =$$

$$16. f(x) = x^5 + x^3 - x, f''(2) =$$

$$17. f(x) = 4x^3 - 3x^2 + x, f'(-1) =$$

$$18. f(x) = x^3 - 3x^2 + 5x, f''(2) =$$

$$19. \mathbf{f(x) = 4x^3 - 3x^2 + 2x, \mathbf{f''(1) =}}$$

$$20. f(x) = 2x^2 - 3x + 4, f'(-1) =$$

$$21. f(x) = 4 - 3x - 2x^2, f'(-1) =$$

$$22. g(x) = x^3 - 3x - 3, g'(-3) =$$

$$23. g(x) = 2x^3 + 3x^2 + 5, g''(4) =$$

$$24. h(x) = 1 + 2x^2 - 3x^3, h''(4) =$$

$$25. f(x) = 4 - 3x^2 + 2x^3, f''(5) =$$

$$26. f(x) = x^3 - 3x + 3, f'(-3) =$$

$$27. f(x) = x^4 - 4x^2 + 4, f'(-4) =$$

$$28. f(x) = 3x^3 + 3x - 3, f'(-3) =$$

$$29. f(x) = 3x^2 - 4x + 2, f'\left(\frac{1}{3}\right) =$$

3.6.3 Integration

Again, only basic integration is required for the number sense test. The technique for integrating is essentially taking the derivative backwards (or anti-derivative) and then plugging in the limits of integration. The following shows a generic polynomial being integrated:

$$\int_a^b a_n x^n + a_{n-1} x^{n-1} + \dots + a_1 x^1 + a_0 x^0 dx = F(x) = \left(\frac{a_n}{n+1} x^{n+1} + \frac{a_{n-1}}{n} x^n + \dots + \frac{a_1}{2} x^2 + \frac{a_0}{1} x^1 \right)_a^b = F(b) - F(a)$$

Let's look at an example:

Evaluate $\int_0^2 3x^2 - x \, dx$.

$$\int_0^2 3x^2 - x \, dx = \left(x^3 - \frac{1}{2}x^2\right)_0^2 = (2^3 - \frac{1}{2}2^2) - (0^3 - \frac{1}{2} \cdot 0) = \mathbf{6}$$

Again, you can apply the table in the previous section for computing integrals of functions (just go in reverse).

To end this section on Integration, there is one special case when integrating which makes the integral trivial, and that is:

$$\int_{-a}^a \text{Odd Function} \, dx = 0$$

So when you are integrating an odd function whose limits are negatives of each other, the result is zero. Let's look at an example of where to apply this:

$$\int_{-\frac{\pi}{4}}^{\frac{\pi}{4}} \sin(x) \, dx = 0$$

Since sine is an odd function, the integral (with the appropriate negative limits) is simply zero!

The following are some more practice problems concerning integration:

Problem Set 3.6.3

1. $\int_0^2 x^2 + 3 \, dx =$

10. $\int_0^\pi \cos x \, dx =$

2. $\int_2^4 2x - 3 \, dx =$

11. $\int_0^3 \frac{x}{3} \, dx =$

3. $\int_1^4 2x \, dx =$

12. $\int_1^3 x^2 \, dx =$

4. $\int_{-3}^3 x^2 \, dx =$

13. $\int_1^3 \frac{3x}{2} \, dx =$

5. $\int_0^4 \frac{x}{2} \, dx =$

14. $\int_1^3 x^{-2} \, dx =$

6. $\int_0^1 x^{\frac{3}{4}} \, dx =$

15. $\int_1^{\frac{3}{2}} x^{-2} \, dx =$

7. $\int_1^3 (x^2 - 2) \, dx =$

16. $\int_0^1 1 - x^2 \, dx =$

8. $\int_{-2}^4 x + 1 \, dx =$

17. $\int_0^4 \sqrt{x} \, dx =$

9. $\int_0^\pi \sin x \, dx =$

18. $\int_{-1}^2 4x \, dx =$

$$19. \int_0^3 x^2 dx =$$

$$20. \int_1^e \frac{2}{x} dx =$$

$$21. \int_0^4 x - 1 dx =$$

$$22. \int_0^2 x^3 dx =$$

$$23. \int_1^e \frac{-3}{x} dx =$$

$$24. \int_0^3 2\mathbf{x} + 1 d\mathbf{x} =$$

$$25. \int_0^1 x^{\frac{2}{3}} dx =$$

$$26. \int_0^{14} 13 - \mathbf{x} d\mathbf{x} =$$

$$27. \int_{-1}^1 x + 1 dx =$$

$$28. \int_0^1 \sqrt{x} dx =$$

$$29. \int_0^1 \sqrt[3]{x} dx =$$

$$30. \int_{-1}^2 3x^2 dx =$$

$$31. \int_2^4 \frac{3}{5}x dx =$$

$$32. \int_1^2 x^3 dx =$$

$$33. \int_0^2 x^3 dx =$$

$$34. \int_0^2 \mathbf{x}^3 + 1 d\mathbf{x} =$$

$$35. \int_0^2 x dx =$$

$$36. \int_{-1}^2 2x dx =$$

$$37. \int_0^4 3 - x dx =$$

$$38. \int_0^2 \frac{3x}{4} dx =$$

$$39. \int_0^3 \frac{4x}{3} dx =$$

4 Additional Problems

The following are assortment of problems which don't occur frequently enough to warrant a section in this book:

5 Solutions

The following are solutions to the practice problems proposed in the previous sections.

Problem Set 1.1:

2850	7020	561	4233
3392	5265	6992	1150
8648	1728	918	847
3828	5418	4968	5644
4680	7776	5073	2450
1088	6688	242	2484
1820	1320	2976	2240
4836	3819	1232	5680
3111	1215	896	1739
7663	8613	2646	1943
3724	3525	2618	4606
2059	5610	845	704
930	2795	5328	2009
1610	5400	3127	7462
238	1809	2125	2475
4384	31672	46812	29430
17277	36950	21489	48312
42658	3564	5994	69030
22270	27664	11022	13545
294849	128472	211554	124890
13431	397946	185364	283251
190005	66789	293007	176712

Problem Set 1.2.1:

- | | | | |
|-----------|------------|------------|-----------------------|
| 1. 594 | 16. 4884 | 31. 36663 | 46. 13794 |
| 2. 792 | 17. 34 | 32. 704 | 47. 12100 |
| 3. 418 | 18. 9657 | 33. 333 | 48. 1815 |
| 4. 5082 | 19. 26 | 34. 22.077 | 49. (*) 648 – 717 |
| 5. 814 | 20. 5883 | 35. 2.42% | 50. 182 |
| 6. 726 | 21. 27 | 36. 1573 | 51. 6776 |
| 7. 2.53 | 22. 203 | 37. 252 | 52. (*) 31181 – 34465 |
| 8. 572 | 23. 2178 | 38. 22066 | 53. 14641 |
| 9. 2706 | 24. 27 | 39. 14641 | 54. 23 |
| 10. 50616 | 25. 4551 | 40. 1452 | 55. 6006 |
| 11. 18 | 26. 3885 | 41. 858 | 56. 136653 |
| 12. 3927 | 27. 38295 | 42. 2662 | 57. 2310 |
| 13. 25 | 28. 222333 | 43. 2420 | 58. 15004 |
| 14. 35631 | 29. 1155 | 44. 2310 | 59. (*) 75897 – 83853 |
| 15. .275 | 30. 14641 | 45. 23 | |

Problem Set 1.2.2:

- | | | | |
|-----------|--------|------------|----------------------|
| 1. 124634 | | 6. 24846 | |
| 2. 2363.4 | 4. 345 | 7. \$15.15 | 9. (*) 14488 – 16014 |
| 3. 37269 | 5. 222 | 8. 448844 | 10. (*) 2398 – 2652 |

Problem Set 1.2.3:

- | | | | |
|----------|--------------------|---------------------|--------------------------------|
| 1. 6000 | 9. 24.64 | 17. (*) 1265 – 1400 | 25. 1280 |
| 2. 10800 | 10. 101 | 18. 85.6% | 26. (*) 185 – 205 |
| 3. 6.5 | 11. .44 | 19. 16.16 | 27. 50075 |
| 4. 3700 | 12. 80800 | 20. 7575 | 28. 4125 |
| 5. 825 | 13. (*) 376 – 417 | 21. (*) 376 – 417 | 29. 6600 |
| 6. 2.56 | 14. $\frac{6}{25}$ | 22. $\frac{7}{25}$ | 30. 4950 |
| 7. 3675 | 15. 5225 | 23. 80.24 | 31. (*) 14842800 –
16405200 |
| 8. 10450 | 16. 850 | 24. 7675 | |

Problem Set 1.2.4:

- | | | | |
|---------|-------------------------------|-------------------|-----------|
| 1. 3600 | 5. (*) 560 – 620 | 12. 19800 | |
| 2. 4800 | 6. 2100 | 9. .64 | 13. 64 |
| 3. .88 | 7. 1800 | 10. (*) 719 – 796 | 14. 54000 |
| 4. 6300 | 8. (*) 10504127 –
11609825 | 11. 1.28 | 15. 72.6 |

Problem Set 1.2.5:

- | | | | |
|---------------------------|--------------------|---------------------------|----------------------------|
| 1. 40000 | | | 12. (*) 4620 – 5107 |
| 2. (*) 189992 –
209992 | 5. (*) 2212 – 2446 | 9. (*) 192850 –
213150 | 13. (*) 3917 – 4330 |
| 3. 1.104 | 6. (*) 1054 – 1166 | 10. 6000 | 14. (*) 384750 –
425250 |
| 4. (*) 425 – 471 | 7. 200000 | 11. (*) 139 – 154 | 15. 153000 |
| | 8. (*) 6628 – 7326 | | |

- | | | | |
|----------------------------------|----------------------------|----------------------------------|----------------------------|
| 16. (*) 307 – 341 | 34. (*) 321 – 356 | 51. (*) 84142 –
93000 | 67. (*) 5277 – 5834 |
| 17. 121 | 35. (*) 474999 –
525000 | 52. (*) 583 – 646 | 68. (*) 118 – 132 |
| 18. (*) 597668 –
660582 | 36. (*) 1030 – 1140 | 53. (*) 58163 –
64286 | 69. (*) 8130 – 8986 |
| 19. (*) 8957133 –
9899991 | 37. (*) 326 – 362 | 54. (*) 7546054 –
8340376 | 70. (*) 6332 – 7000 |
| 20. (*) 114 – 126 | 38. (*) 1576 – 1743 | 55. (*) 664694 –
734662 | 71. (*) 54204 – 59910 |
| 21. 183000 | 39. (*) 461428 –
510000 | 56. (*) 1644 – 1818 | 72. (*) 237 – 263 |
| 22. (*) 7440353 –
8223549 | 40. (*) 38240 – 42267 | 57. 40625 | 73. (*) 50805 – 56154 |
| 23. (*) 1261 – 1395 | 41. (*) 182076 –
201242 | 58. (*) 99071 –
109500 | 74. (*) 14776 – 16332 |
| 24. (*) 646 – 714 | 42. 60.25 | 59. (*) 232071 –
256500 | 75. (*) 12324 – 13622 |
| 25. (*) 22757 – 25153 | 43. (*) 593749 –
656250 | 60. (*) 113491195 –
125437637 | 76. (*) 200163 –
221233 |
| 26. 210000 | 44. (*) 652 – 721 | 61. (*) 18457124 –
20399980 | 77. (*) 577 – 639 |
| 27. (*) 3360 – 3715 | 45. (*) 775848 –
857518 | 62. (*) 484306 –
535286 | 78. (*) 21855 –
24157 |
| 28. 9300 | 46. (*) 1056 – 1168 | 63. (*) 6641817 –
7340957 | 79. (*) 632 – 700 |
| 29. (*) 347699652 –
384299616 | 47. (*) 2253 – 2492 | 64. (*) 24 – 28 | 80. (*) 605 – 670 |
| 30. .02 | 48. (*) 93755 –
103625 | 65. (*) 35624 – 39375 | 81. (*) 1159 – 1283 |
| 31. (*) 5842616 –
6457630 | 49. (*) 4303 – 4757 | 66. (*) 47362 – 52348 | 82. (*) 3167 – 3502 |
| 32. (*) 2020 – 2233 | 50. (*) 450570 –
498000 | | 83. (*) 139 – 155 |
| 33. (*) 3528 – 3900 | | | 84. (*) 117040 –
129362 |

Problem Set 1.2.6:

- | | | | |
|---------|---------|----------|-----------|
| 1. 7.8 | | 10. 2016 | |
| | 6. 4368 | | |
| 2. 72 | | 11. 378 | 15. 1.5 |
| | 7. 840 | | |
| 3. 96 | | 12. 4410 | |
| | 8. 4368 | | 16. 10.56 |
| 4. 720 | | 13. 22.5 | |
| | 9. 3.6 | | 17. 700 |
| 5. 2842 | | 14. 4140 | |

Problem Set 1.2.7:

- | | | | |
|----------|-----------|-------------|--------------------------|
| 1. 8633 | 9. 8544 | 17. 8554 | 25. 9672 |
| 2. 9312 | 10. 8924 | 18. 8918 | 26. 9888 |
| 3. 11227 | 11. 10712 | 19. 987042 | 27. 982081 |
| 4. 9021 | 12. 10506 | 20. 9888 | 28. 10088 |
| 5. 11021 | 13. 8556 | 21. 9579 | 29. 1011024 |
| 6. 8277 | 14. 11342 | 22. 980099 | |
| 7. 11016 | 15. 8633 | 23. 1013036 | 30. (*) 18062 –
19964 |
| 8. 11663 | 16. 8212 | 24. 10379 | 31. 12996 |

Problem Set 1.2.8:

- | | | | |
|----------|----------|----------------------|-------|
| 1. 6.25 | | 6. -24.75 | |
| | 4. 13225 | | 9. 14 |
| 2. 1.225 | | 7. (*) 19699 – 21773 | |
| | 5. 3025 | | |
| 3. 625 | | 8. 255025 | |

Problem Set 1.2.9:

- | | | | |
|-----------|---------|---------------------------|---------|
| 1. 3364 | 3. 2209 | 5. (*) 111720 –
123480 | 7. 3481 |
| 2. 260100 | 4. 2809 | 6. 3136 | 8. 1681 |

Problem Set 1.2.10:

- | | | | |
|----------|----------|-----------|------------------------------|
| 1. 7224 | 13. 1225 | 25. 4842 | 37. 9856 |
| 2. 3021 | 14. 5625 | 26. 900 | 38. (*) 4745 – 5245 |
| 3. 2496 | 15. 4225 | 27. 5625 | 39. (*) 4016 – 4440 |
| 4. 3596 | 16. 1225 | 28. 5625 | 40. (*) 9305 – 10285 |
| 5. 48.96 | 17. 441 | 29. 3025 | 41. (*) 3035 – 3355 |
| 6. 7216 | 18. 4225 | 30. 9975 | 42. (*) 26270 – 29036 |
| 7. 864 | 19. 2025 | 31. 4200 | 43. (*) 101076
111716 |
| 8. 63.84 | 20. 7225 | 32. 936 | 44. 14400 |
| 9. 24.91 | 21. 3025 | 33. 7200 | 45. (*) 62132 – 68673 |
| 10. 3025 | 22. 1064 | 34. 625 | 46. (*) 1423267 –
1573085 |
| 11. 9984 | 23. 2000 | 35. 1073 | |
| 12. 7225 | 24. 5625 | 36. –7200 | |

Problem Set 1.2.11:

- | | | | |
|---------|---------|---------|----------|
| 1. 1462 | 4. 252 | 7. 765 | 10. 2268 |
| 2. 736 | 5. 1944 | 8. 574 | 11. 1008 |
| 3. 403 | 6. 976 | 9. 1458 | 12. 1612 |

Problem Set 1.3.1:

- | | | | |
|------------|------------|------------|---------------|
| 1. 40804 | 12. 91809 | 23. 509796 | 34. 38688 |
| 2. 164836 | 13. 826281 | 24. 49374 | 35. 37942 |
| 3. 253009 | 14. 161604 | 25. 23632 | 36. 274576 |
| 4. 368449 | 15. 499849 | 26. 67196 | 37. 41363 |
| 5. 43264 | 16. 34013 | 27. 24969 | 38. 19881 |
| 6. 93636 | 17. 644809 | 28. 49731 | 39. 108332 |
| 7. 259081 | 18. 163216 | 29. 46144 | 40. 25864 |
| 8. 646416 | 19. 262144 | 30. 204020 | 41. 144288144 |
| 9. 495616 | 20. 37942 | 31. 35143 | 42. 444889 |
| 10. 166464 | 21. 374544 | 32. 15004 | |
| 11. 362404 | 22. 96942 | 33. 842724 | |

Problem Set 1.3.2:

- | | | | |
|---------|-----------|------------|----------|
| 1. 640 | 8. 10020 | 15. 5100 | 22. 1024 |
| 2. 810 | 9. 1280 | 16. 660 | 23. 330 |
| 3. 450 | 10. 380 | 17. 490 | 24. 484 |
| 4. 0 | 11. 12030 | 18. -10030 | 25. 2450 |
| 5. 1210 | 12. 0 | 19. 384 | 26. 870 |
| 6. -660 | 13. 441 | 20. 14.4 | 27. 256 |
| 7. 16.9 | 14. 960 | 21. 196 | 28. 3540 |

- | | | | |
|----------|-----------|------------------------|----------------------|
| 29. -196 | 40. 14280 | 51. 910 | 61. (*) 2050 - 2266 |
| 30. 289 | 41. 1560 | 52. (*) 1825 - 2019 | 62. 1584 |
| 31. -289 | 42. -324 | 53. 3300 | 63. (*) 4698 - 5194 |
| 32. 1080 | 43. 3300 | 54. 720 | 64. 2250 |
| 33. 4830 | 44. 9900 | 55. (*) 12108
13384 | 65. 4662 |
| 34. 2002 | 45. 0 | 56. (*) 9076 - 10032 | 66. -588 |
| 35. 1210 | 46. -1210 | 57. 1056 | 67. (*) 9516 - 10518 |
| 36. 2160 | 47. 2775 | 58. 11990 | 68. 2100 |
| 37. 6320 | 48. 540 | 59. (*) 8015 - 8859 | 69. 3774 |
| 38. 1188 | 49. 576 | 60. 672 | 70. (*) 3659 - 4045 |
| 39. 363 | 50. 16770 | | |

Problem Set 1.3.3:

- | | | | |
|---------|--------|---------|----------|
| 1. 2521 | | 4. 1301 | |
| | 3. 481 | | 6. 12961 |
| 2. 313 | | 5. 3281 | |

Problem Set 1.3.4:

- | | | | |
|---------|----------|----------|----------|
| 1. 462 | 7. 1920 | 13. 2280 | 19. 550 |
| 2. 1920 | 8. 12960 | 14. 3360 | |
| | | | 20. 128 |
| 3. 380 | 9. 550 | 15. 128 | |
| 4. 128 | 10. 3760 | 16. 1680 | 21. 5300 |
| 5. 160 | 11. 6380 | 17. 880 | |
| | | | 22. 2300 |
| 6. 124 | 12. 120 | 18. 450 | |

Problem Set 1.3.5:

- | | | | |
|---------|---------|---------|---------|
| 1. 9090 | 3. 5353 | 5. 4141 | 7. 6161 |
| 2. 505 | 4. 6868 | 6. 4545 | 8. 5858 |

Problem Set 1.3.6:

- | | | | |
|---------|-----------|-----------|-----------|
| 1. 145 | 13. -172 | 25. 170 | 37. 78 |
| 2. 140 | 14. 11.2 | 26. 438 | 38. -238 |
| 3. -115 | 15. -6.72 | 27. -363 | 39. -900 |
| 4. 133 | 16. 254 | 28. 302 | 40. 168 |
| 5. 272 | 17. 540 | 29. 720 | 41. 1014 |
| 6. -109 | 18. 218 | 30. 218 | 42. -1540 |
| 7. 264 | 19. 300 | 31. -1560 | 43. -616 |
| 8. 175 | 20. -30 | 32. -70 | 44. 715 |
| 9. -97 | 21. -94 | 33. -170 | 45. -272 |
| 10. 193 | 22. 525 | 34. 288 | 46. 672 |
| 11. 153 | 23. 326 | 35. 105 | 47. 894 |
| 12. 107 | 24. 321 | 36. 18 | |

Problem Set 1.3.7:

- | | | | |
|---------|---------|---------|---------|
| 1. 1575 | 3. 2275 | 5. 2925 | 7. 6175 |
| 2. 4275 | 4. 4675 | 6. 2975 | 8. 5225 |

Problem Set 1.3.8:

- | | | | |
|-----------------------|-----------------------|-----------------------|-----------------------|
| 1. $35\frac{1}{16}$ | 9. $137\frac{4}{25}$ | 17. $138\frac{2}{3}$ | 25. 41 |
| 2. $72\frac{2}{9}$ | 10. 53.04 | 18. $21\frac{7}{12}$ | 26. $44\frac{4}{9}$ |
| 3. $12\frac{4}{25}$ | 11. $40\frac{4}{9}$ | 19. 131 | 27. 79.04 |
| 4. $29\frac{1}{6}$ | 12. $101\frac{1}{49}$ | 20. $64\frac{4}{9}$ | 28. $21\frac{7}{15}$ |
| 5. $101\frac{1}{16}$ | 13. $53\frac{1}{25}$ | 21. $160\frac{4}{9}$ | 29. 5 |
| 6. $139\frac{1}{36}$ | 14. $131\frac{1}{25}$ | 22. $351\frac{1}{49}$ | 30. 5.7 |
| 7. $75\frac{1}{36}$ | 15. $29\frac{4}{25}$ | 23. 9 | 31. $12\frac{24}{25}$ |
| 8. $245\frac{1}{121}$ | 16. $131\frac{1}{64}$ | 24. 9.03 | |

Problem Set 1.3.9:

- | | | | |
|----------------------|-----------------------|-----------------------|-----------------------|
| 1. $8\frac{9}{14}$ | 7. $28\frac{9}{34}$ | 13. $-\frac{17}{18}$ | 19. $-3\frac{11}{15}$ |
| 2. $19\frac{9}{25}$ | 8. $8\frac{9}{17}$ | 14. $-2\frac{16}{25}$ | 20. $-2\frac{8}{17}$ |
| 3. $15\frac{16}{23}$ | 9. $11\frac{9}{14}$ | 15. $-2\frac{8}{17}$ | 21. $-1\frac{13}{17}$ |
| 4. $22\frac{25}{32}$ | 10. $23\frac{9}{16}$ | 16. $30\frac{16}{21}$ | 22. $37\frac{9}{38}$ |
| 5. $13\frac{9}{19}$ | 11. $13\frac{16}{17}$ | 17. $-2\frac{7}{16}$ | 23. $-1\frac{11}{15}$ |
| 6. $24\frac{25}{34}$ | 12. $-\frac{13}{14}$ | 18. $-\frac{11}{12}$ | |

Problem Set 1.3.10:

- | | | | |
|----------------------|--------------------|-----------------|----------------------|
| 1. (*) 2553 – 2823 | 4. (*) 5958 – 6586 | 118041 | |
| 2. (*) 25356 – 28026 | 3. (*) 3149 – 3481 | 5. (*) 106799 – | 6. (*) 34108 – 37700 |

- | | | | |
|-------------------------|-------------------------|-------------------------|-----------------------------|
| 7. (*) 39398 – 43545 | 15. (*) 80548 – 89028 | 24. (*) 523488 – 578592 | 33. (*) 53437500 – 59062500 |
| 8. (*) 126445 – 139755 | 16. (*) 65555 – 72457 | 25. (*) 25536 – 28224 | 34. (*) 25650 – 28350 |
| 9. (*) 14630 – 16170 | 17. (*) 60693 – 67083 | 26. (*) 298452 – 329868 | 35. (*) 95000 – 105000 |
| 10. (*) 624255 – 689967 | 18. (*) 60762 – 67158 | 27. (*) 260646 – 288084 | 36. (*) 475089 – 525099 |
| 11. (*) 97917 – 108225 | 19. (*) 2048 – 2265 | 28. (*) 1740 – 1924 | 37. (*) 3910 – 4322 |
| 12. (*) 760958 – 841060 | 20. (*) 86184 – 95256 | 29. (*) 8257 – 9127 | 38. (*) 150292 – 166114 |
| 13. (*) 31005 – 34269 | 21. (*) 157586 – 174174 | 30. (*) 5728 – 6332 | 39. 2592 |
| 14. (*) 9771 – 10801 | 22. (*) 7524 – 8316 | 31. (*) 28260 – 31236 | 40. (*) 3406 – 3766 |
| | 23. (*) 34108 – 37700 | 32. (*) 3513 – 3883 | |

Problem Set 1.4.1:

- | | | | |
|------|------|------|------|
| 1. 0 | 3. 3 | 5. 0 | 7. 4 |
| 2. 2 | 4. 3 | 6. 5 | |

Problem Set 1.4.2:

- | | | | |
|------|------|------|------|
| 1. 2 | | 4. 7 | |
| 2. 5 | 3. 0 | 5. 2 | 6. 8 |

Problem Set 1.4.3:

- | | | | |
|------|------|------|------|
| 1. 8 | 3. 9 | 5. 0 | 7. 5 |
| 2. 5 | 4. 9 | 6. 8 | 8. 7 |

9. 4

10. 4

Problem Set 1.4.4:

1. 4

5. 0

9. 7

13. 0

2. 2

6. 3

10. 0

14. 2

3. 2

7. 0

11. 8

4. 6

8. 6

12. 7

15. 6

Problem Set 1.4.5:

1. 1

8. 0

15. 3

22. 6

2. 3

9. 4

16. 3

23. 2

3. 0

10. 1

17. 0

24. 3

4. 2

11. 2

18. 5

25. 2

5. 2

12. 5

19. 2

6. 2

13. 2

20. 4

26. 2

7. 4

14. 2

21. 3

27. 2

Problem Set 1.4.6:

1. $39\frac{1}{3}$

3. $222\frac{5}{9}$

5. $50\frac{2}{3}$

7. $1371\frac{2}{3}$

2. $55\frac{8}{9}$

4. $35\frac{2}{3}$

6. $137\frac{1}{9}$

8. 55

Problem Set 1.4.7:

- | | | | |
|----------|---------------------|-----------|---------------------|
| 1. 2.5% | 6. .075 | 11. 27.5% | 16. 6.25% |
| 2. 7.5% | 7. $1\frac{1}{4}\%$ | 12. .045 | 17. $\frac{11}{40}$ |
| 3. 17.5% | 8. 20% | 13. 18 | 18. .32 |
| 4. 52.5% | 9. 17.5% | 14. .025 | 19. 8% |
| 5. 1.075 | 10. 40 | 15. 435% | 20. .0081 |

Problem Set 1.5.1:

- | | | | |
|---------|--------|----------|-----------|
| 1. 198 | | 6. 2997 | |
| 2. -396 | 4. 495 | 7. -3996 | 9. -198 |
| 3. 1998 | 5. 99 | 8. -999 | 10. -4995 |

Problem Set 1.5.2:

- | | | | |
|----------------------|---------------------|----------------------|----------------------|
| 1. $-1\frac{1}{6}$ | 5. $-1\frac{4}{7}$ | 9. $-1\frac{8}{9}$ | 13. $-8\frac{1}{12}$ |
| 2. $-1\frac{14}{15}$ | 6. $-\frac{7}{8}$ | 10. $-7\frac{1}{14}$ | 14. $-6\frac{1}{12}$ |
| 3. -2 | 7. $-4\frac{1}{8}$ | 11. $-3\frac{1}{6}$ | 15. $-4\frac{1}{2}$ |
| 4. $-1\frac{17}{20}$ | 8. $-5\frac{1}{10}$ | 12. $-1\frac{5}{6}$ | 16. $-1\frac{3}{5}$ |

Problem Set 1.5.3:

- | | | | |
|-------------------|-------------------|-------------------|-------------------|
| 1. $\frac{4}{21}$ | 2. $\frac{1}{24}$ | 3. $\frac{3}{40}$ | 4. $1\frac{1}{6}$ |
|-------------------|-------------------|-------------------|-------------------|

Problem Set 1.5.4:

- | | | | |
|----------------------|---------------------|-----------------------|------------------------|
| 1. $2\frac{1}{156}$ | 6. $1\frac{4}{143}$ | 12. $-\frac{31}{35}$ | 17. $3\frac{1}{156}$ |
| 2. $2\frac{1}{30}$ | 7. $1\frac{36}{91}$ | 13. $1\frac{4}{255}$ | 18. $1\frac{2}{35}$ |
| 3. $2\frac{16}{285}$ | 8. $\frac{1}{30}$ | 14. $1\frac{16}{165}$ | 19. $1\frac{1}{132}$ |
| 4. $\frac{4}{15}$ | 9. $1\frac{4}{195}$ | 15. $1\frac{4}{143}$ | 20. $\frac{49}{330}$ |
| 5. $1\frac{4}{35}$ | 10. 2 | 16. $1\frac{1}{210}$ | 21. $-\frac{145}{154}$ |
| | 11. 3 | | |

Problem Set 1.5.5:

- | | | | |
|----------------------|------------------------|-----------------------|-----------------------|
| 1. $\frac{13}{252}$ | 7. $\frac{11}{584}$ | 13. $-\frac{22}{435}$ | 19. $\frac{11}{448}$ |
| 2. $\frac{9}{203}$ | 8. $\frac{9}{430}$ | 14. $\frac{13}{328}$ | |
| 3. $\frac{17}{520}$ | 9. $-\frac{11}{42}$ | 15. $\frac{17}{333}$ | 20. $-\frac{11}{414}$ |
| 4. $\frac{22}{915}$ | 10. $\frac{17}{900}$ | 16. $\frac{7}{165}$ | 21. $\frac{11}{328}$ |
| 5. $\frac{19}{495}$ | 11. $-\frac{37}{1620}$ | 17. $\frac{27}{784}$ | |
| 6. $\frac{19}{1342}$ | 12. $\frac{11}{328}$ | 18. $\frac{19}{1342}$ | 22. $\frac{18}{979}$ |

Problem Set 2.1.1:

- | | | | |
|----------|---------|----------|----------|
| 1. 784 | 6. 4.84 | 11. 324 | 16. 196 |
| 2. 10.24 | 7. 1156 | 12. 5.76 | 17. 441 |
| 3. 841 | 8. 289 | 13. 529 | 18. 576 |
| 4. 256 | 9. 529 | 14. 1024 | 19. 9.61 |
| 5. 961 | 10. 361 | 15. 484 | 20. 7.29 |

- | | | | |
|----------|-----------------------|-----------------------|----------------------------|
| 21. 784 | 27. (*) 972 – 1075 | 33. (*) 36495 – 40337 | 39. (*) 79344 – 87698 |
| 22. 1156 | 28. (*) 372 – 412 | 34. (*) 379 – 420 | 40. (*) 241 – 267 |
| 23. 676 | 29. –224 | 35. (*) 28227 – 31200 | 41. (*) 496 – 549 |
| 24. 289 | 30. .324 | 36. (*) 27132 – 29990 | 42. (*) 975 – 1078 |
| 25. 1089 | 31. (*) 14546 – 16078 | 37. (*) 9098 – 10057 | |
| 26. –27 | 32. (*) 7553 – 8349 | 38. (*) 13166 – 14553 | 43. (*) 184756 –
204206 |

Problem Set 2.1.2:

- | | | | |
|----------|---------------------|------------------------------|------------------------------|
| 1. 12 | 12. 512 | 22. $\frac{1}{2}$ | 32. –216 |
| 2. 1331 | 13. 3375 | 23. 370 | 33. $\frac{1}{2}$ |
| 3. 3744 | 14. 1728 | 24. –2 | 34. 225 |
| 4. –7 | 15. $\frac{5}{4}$ | 25. 1.2 | 35. –217 |
| 5. 1728 | 16. 2197 | 26. 64000 | 36. (*) 169059 –
186855 |
| 6. 4096 | 17. 343 | 27. 1331 | 37. 343000 |
| 7. 2 | 18. –11 | 28. 1.1 | 38. (*) 1641486 –
1814374 |
| 8. 1331 | 19. 216 | 29. (*) 692464 –
765356 | 39. (*) 2669363 –
2950349 |
| 9. –1728 | 20. 3375 | 30. .9 | 40. 4096 |
| 10. 13 | 21. (*) 1653 – 1828 | 31. (*) 1682982 –
1860140 | |
| 11. –9 | | | |

Problem Set 2.1.3:

- | | | | |
|--------|---------------------|-----------------|-------------|
| 1. 160 | 10. -26 | 19. 7 | 28. 648000 |
| 2. -83 | 11. 81 | 20. -144 | 29. 98000 |
| 3. 32 | 12. 3200000 | 21. .081 | 30. 14400 |
| 4. 243 | 13. 729 | 22. 288 | 31. 21600 |
| 5. -61 | 14. .04 | 23. 29 | 32. 2025000 |
| 6. 3 | 15. 4000 | 24. (*) 61 - 69 | 33. 2500 |
| 7. 160 | 16. 2560000 | 25. 2.5 | 34. 64800 |
| 8. 98 | 17. (*) 3242 - 3584 | 26. 64000 | 35. 8100000 |
| 9. 40 | 18. 40000 | 27. 512000 | 36. 144000 |

Problem Set 2.1.4:

- | | | | |
|------------------|----------------------|-----------------------|----------------------|
| 1. $\frac{1}{8}$ | 11. -.875 | 20. $43\frac{6}{7}\%$ | 29. -.375 |
| 2. 220% | 12. 275% | 21. $77\frac{7}{9}\%$ | 30. $-\frac{4}{3}$ |
| 3. .56 | 13. $\frac{5}{9}$ | 22. $\frac{1}{2}$ | 31. $7\frac{1}{7}\%$ |
| 4. $\frac{5}{8}$ | 14. $\frac{2}{3}$ | 23. .56 | 32. $\frac{7}{16}$ |
| 5. 2.125 | 15. $-\frac{16}{17}$ | 24. $\frac{9}{11}$ | 33. 15 |
| 6. 1 | 16. .46 | 25. $\frac{4}{3}$ | 34. 3 |
| 7. 60% | 17. $-\frac{8}{9}$ | 26. 43.75% | 35. -.27 |
| 8. .125 | 18. $\frac{3}{8}$ | 27. 176 | 36. 121 |
| 9. .81 | 19. .8 | 28. $28\frac{4}{7}\%$ | 37. $-\frac{7}{18}$ |
| 10. 6.25% | | | 38. 31.25% |

- | | | | |
|-----------------------|-----------------------|------------------------|------------------------|
| 39. 1331 | 47. 10021 | 55. 13.31 | 63. $\frac{1}{14}$ |
| 40. $\frac{3}{14}$ | 48. $8\frac{1}{3}\%$ | 56. 18.75% | 64. 2400 |
| 41. 8 | 49. $78\frac{4}{7}\%$ | 57. $\frac{1}{16}$ | 65. $\frac{11}{160}$ |
| 42. $10\frac{4}{5}$ | 50. 800 | 58. $121\frac{3}{7}\%$ | 66. $92\frac{6}{7}\%$ |
| 43. $21\frac{3}{7}\%$ | 51. 1331 | 59. $\frac{3}{7}$ | 67. $\frac{2}{65}$ |
| 44. $\frac{5}{14}$ | 52. $80\frac{1}{3}$ | 60. $\frac{3}{80}$ | 68. $107\frac{1}{7}\%$ |
| 45. 6 | 53. 135 | 61. $\frac{11}{1000}$ | 69. $\frac{3}{14}$ |
| 46. $\frac{11}{14}$ | 54. $\frac{9}{14}$ | 62. $\frac{13}{14}$ | |

Problem Set 2.1.5:

- | | | | |
|-----------|------------|---------------------|-------------------|
| 1. 12012 | 12. 36036 | 23. 7070.7 | 34. 36 |
| 2. 54 | 13. 7 | 24. 121.121 | 35. (*) 712 – 788 |
| 3. 505.05 | 14. 1073 | 25. 35035 | 36. $\frac{1}{3}$ |
| 4. 25025 | 15. 30030 | 26. 909.09 | 37. 6006 |
| 5. 70707 | 16. 70070 | 27. 505505 | 38. 49 |
| 6. 37 | 17. 999 | 28. 9009 | 39. 13 |
| 7. 20020 | 18. 55055 | 29. 1111.11 | 40. 48 |
| 8. 15015 | 19. 75075 | 30. 303303 | 41. 10010 |
| 9. 27027 | 20. 153153 | 31. 5005 | 42. 96 |
| 10. 29 | 21. 10010 | 32. $28\frac{7}{9}$ | 43. 256 |
| 11. 60 | 22. 18018 | 33. 7007 | 44. 11011 |

- | | | | |
|--------------------|---------------------|----------------------|-----------|
| 45. 9009 | 51. 9009 | 57. 147 | 63. 13013 |
| 46. 384 | 52. 90 | 58. $32\frac{8}{9}$ | 64. 324 |
| 47. 36036 | 53. $16\frac{4}{9}$ | 59. 81 | 65. 185 |
| 48. 7007 | 54. 11011 | 60. 74 | 66. 175 |
| 49. 60 | 55. 96 | 61. $789\frac{1}{3}$ | 67. 9009 |
| 50. $8\frac{2}{9}$ | 56. $24\frac{2}{3}$ | 62. 37 | 68. 15015 |

Problem Set 2.1.6:

- | | | | |
|---------|----------|---------------------|-----------|
| 1. 2042 | 10. 2222 | 19. 1364 | 27. 20 |
| 2. 44 | 11. -89 | 20. 2006 | 28. 401 |
| 3. 2003 | 12. 2100 | 21. 556 | 29. 2997 |
| 4. 199 | 13. 999 | 22. 505 | 30. 11011 |
| 5. 1666 | 14. 534 | 23. 1530 | 31. 50175 |
| 6. 1544 | 15. 2017 | 24. $66\frac{8}{9}$ | 32. 84 |
| 7. 277 | 16. 2007 | 25. 34 | 33. 22066 |
| 8. 1459 | 17. 1664 | 26. 2005 | 34. 10.1 |
| 9. 999 | 18. 1666 | | |

Problem Set 2.1.7:

- | | | | |
|-------|-------|-------|--------|
| 1. 20 | 4. 10 | 7. 6 | 10. 12 |
| 2. 20 | 5. 4 | 8. 20 | 11. 20 |
| 3. 16 | 6. 12 | 9. 6 | |

Problem Set 2.1.8:

- | | | | |
|------------------|--------------------|-----------------------|-----------------------|
| 1. (*) 185 – 205 | 5. (*) 5052 – 5585 | 9. (*) 995 – 1100 | 13. (*) 5052 – 5585 |
| 2. (*) 683 – 756 | 6. (*) 342 – 379 | 10. (*) 664 – 734 | 14. (*) 493 – 546 |
| 3. (*) 51 – 58 | 7. (*) 1608 – 1778 | 11. (*) 15384 – 17005 | 15. (*) 46339 – 51218 |
| 4. (*) 290 – 322 | 8. (*) 7495 – 8285 | 12. (*) 1221 – 1350 | 16. (*) 524 – 581 |

Problem Set 2.1.9:

- | | | | |
|--------|----------|----------|----------|
| 1. 22 | 6. 220 | 11. 30 | 16. 8 |
| 2. 126 | 7. 440 | 12. 10 | 17. 66 |
| 3. 4.5 | 8. 1760 | 13. 44 | 18. 22.5 |
| 4. 240 | 9. 81 | 14. .5 | 19. 11 |
| 5. 132 | 10. 3520 | 15. 2160 | |

Problem Set 2.1.10:

- | | | | |
|---------|---------|--------------------|----------|
| 1. 81 | 5. 27 | 9. 10000 | 13. 36 |
| 2. 1728 | 6. 3 | 10. $1\frac{1}{3}$ | 14. 3456 |
| 3. 81 | 7. 5184 | 11. 1.5 | |
| 4. 3 | 8. 2.5 | 12. 1 | 15. 500 |

Problem Set 2.1.11:

- | | | | |
|-------|-------|--------|--------|
| 1. 4 | | 4. 693 | |
| 2. 32 | 3. 48 | 5. 154 | 6. 308 |

- | | | | |
|------------------|-----------|-------------------|---------|
| 7. 96 | 14. 12.5% | 21. 2 | 28. 231 |
| 8. 16 | 15. 225% | 22. 1.75 | 29. 63 |
| 9. $\frac{3}{8}$ | 16. 400% | 23. 5 | 30. 12 |
| 10. 6 | 17. 11 | 24. 3 | 31. 147 |
| 11. 2 | 18. 600% | 25. 2.5 | 32. 128 |
| 12. 50% | 19. 37.5% | 26. 112 | 33. 320 |
| 13. 400% | 20. 1155 | 27. $\frac{1}{8}$ | |

Problem Set 2.1.12:

- | | | |
|-------|--------|-------|
| 1. 77 | 2. -40 | 3. 37 |
|-------|--------|-------|

Problem Set 2.2.1.:

- | | | | |
|------------------|---------------------|--------------------|--------------------|
| 1. 132 | 10. 132 | 19. 143 | 28. 528 |
| 2. 231 | 11. 81 | 20. $\frac{4}{5}$ | 29. $4\frac{1}{6}$ |
| 3. 169 | 12. 506 | 21. 117 | 30. 98 |
| 4. 123 | 13. 1.5 | 22. $5\frac{1}{3}$ | 31. 126 |
| 5. 18 | 14. 441 | 23. 2.5 | 32. 207 |
| 6. 240 | 15. 396 | 24. 108 | 33. 91 |
| 7. 100 | 16. 255 | 25. 462 | 34. $6\frac{1}{4}$ |
| 8. $\frac{2}{5}$ | 17. 4 | 26. -3 | 35. $2\frac{2}{3}$ |
| 9. 96 | 18. $-1\frac{1}{8}$ | 27. 264 | 36. 255 |

37. 147	45. 81	53. 37	61. 80
38. 98	46. 6	54. 3.2	62. 4.8
39. 1150	47. 294	55. (*) 179763 – 198687	63. 77
40. 16	48. 726	56. (*) 4138 – 4574	64. (*) 7866 – 8696
41. 264	49. 161	57. 11	65. 3
42. (*) 418 – 464	50. 273	58. 141	66. 16
43. 396	51. 168	59. 1.5	67. (*) 25863 – 28587
44. 242	52. 528	60. $9\frac{1}{3}$	68. (*) 1231 – 1361

Problem Set 2.2.2:

1. 750	6. 610	11. 372	16. 143
2. 372	7. 893	12. 114	17. 319
3. 514	8. 534	13. 88	18. 693
4. 660	9. 284	14. 6	19. 748
5. 804	10. 304	15. 196	

Problem Set 2.2.3:

1. 3	6. 10	11. 36	16. 15
2. 9	7. 8	12. 56	17. 192
3. 96	8. 12	13. 20	18. 78
4. 4	9. 9	14. 42	19. 42
5. 10	10. 124	15. 5	20. 8

21. 56		30. 35	
	26. 24		35. 7
22. 70		31. 240	
	27. 24		36. 35
23. 7		32. 7	
	28. 39		37. 7
24. 55		33. 10	
	29. 54		38. 124
25. 385		34. 24	

Problem Set 2.2.4:

1. 5	3. 5	5. 20	7. 2
2. 9	4. 27	6. 35	8. 14

Problem Set 2.2.5:

1. 140	3. 45	5. 120	7. 133
2. 108	4. 1080	6. 1440	8. 540

Problem Set 2.2.6:

1. 70		8. 51	
	5. 276		12. 66
2. 40		9. 45	
	6. 112		13. 36
3. 35		10. 66	
	7. 35		14. 18
4. 176		11. 78	

Problem Set 2.2.7:

1. 9	2. 40	3. 26	4. 15
------	-------	-------	-------

- | | | | |
|-------|-------|-------|--------|
| 5. 4 | 8. 5 | 11. 7 | 14. 8 |
| 6. 8 | 9. 15 | 12. 6 | 15. 84 |
| 7. 33 | 10. 6 | 13. 9 | 16. 84 |

Problem Set 2.2.8:

- | | | | |
|------|----------------|-------|------|
| 1. 3 | 3. $2\sqrt{3}$ | 5. 4 | 7. 3 |
| 2. 6 | 4. 12 | 6. 18 | 8. 9 |

Problem Set 2.2.9:

- | | | | |
|-------------|-------|---------|--------|
| 1. 726 | | 6. 216 | |
| 2. 144π | 4. 96 | 7. 512 | 9. 224 |
| 3. 27 | 5. 64 | 8. 1728 | |

Problem Set 2.2.10:

- | | | | |
|--------|--------|---------------------|-------------------|
| 1. 60 | 7. 28 | 13. 6 | 18. 12 |
| 2. 10 | 8. 10 | 14. 24 | 19. 720 |
| 3. 20 | 9. 336 | | |
| 4. 35 | 10. 56 | 15. $\frac{1}{120}$ | 20. $\frac{1}{6}$ |
| 5. 840 | 11. 36 | 16. 6 | 21. 10 |
| 6. 30 | 12. 2 | 17. 4 | 22. 200 |

Problem Set 2.2.11:

- | | | | |
|-------------------|--------------------|--------------------------|--------------------|
| 1. $-\frac{1}{2}$ | 16. $\frac{1}{2}$ | 31. -4 | 46. $-\frac{1}{2}$ |
| 2. $\frac{8}{3}$ | 17. 1 | 32. 3 | 47. 45 |
| 3. 0 | 18. -2 | 33. 225 | 48. $\frac{1}{2}$ |
| 4. 1 | 19. -2 | 34. $\frac{1}{3}$ | 49. $-\frac{3}{4}$ |
| 5. $\frac{1}{2}$ | 20. -1 | 35. $\frac{\sqrt{2}}{2}$ | 50. $\frac{1}{4}$ |
| 6. -10 | 21. 108 | 36. -1 | 51. $\frac{6}{5}$ |
| 7. $-\frac{1}{2}$ | 22. -1 | 37. -1 | 52. $\frac{1}{4}$ |
| 8. -1 | 23. -1 | 38. $\frac{3}{2}$ | 53. $\frac{1}{3}$ |
| 9. 10 | 24. 45 | 39. -1 | 54. $\frac{1}{4}$ |
| 10. $\frac{1}{3}$ | 25. $\frac{14}{9}$ | 40. $\frac{1}{2}$ | 55. 4 |
| 11. 112.5 | 26. 1 | 41. 0 | 56. $\frac{1}{2}$ |
| 12. 36 | 27. 12 | 42. $-\frac{1}{3}$ | 57. $-\frac{3}{4}$ |
| 13. 0 | 28. -1 | 43. 1 | 58. $-\frac{1}{4}$ |
| 14. -1 | 29. 2 | 44. $-\frac{1}{3}$ | 59. $3\frac{1}{2}$ |
| 15. 0 | 30. 15 | 45. $-\frac{3}{4}$ | |

Problem Set 2.2.12:

- | | | | |
|------------------|-------------------|-------------------|--------------------|
| 1. 1 | 4. $-\frac{1}{2}$ | 7. $\frac{3}{4}$ | 10. $\frac{1}{2}$ |
| 2. $\frac{1}{2}$ | 5. $\frac{3}{4}$ | 8. $-\frac{1}{2}$ | 11. $-\frac{1}{4}$ |
| 3. $\frac{1}{2}$ | 6. 68 | 9. 3 | 12. 308 |

13. $-\frac{1}{2}$

14. $\frac{1}{2}$

15. $\frac{1}{4}$

16. $\frac{1}{2}$

17. $\frac{1}{2}$

18. $\frac{7}{25}$

19. $\frac{1}{2}$

20. $\frac{1}{4}$

21. $-\frac{1}{2}$

22. 1

23. $-\frac{1}{2}$

24. 1

25. $\frac{1}{2}$

26. $-\frac{1}{2}$

27. -2

Problem Set 2.2.13:

1. 4

2. 5

3. 2

4. 3

5. 8

6. -3

7. 5

8. -2

9. $\frac{\pi}{6}$

10. $\frac{1}{2}$

11. 10π

12. 2

Problem Set 2.2.14:

1. -9

2. -5

3. $1\frac{1}{4}$

Problem Set 2.2.15:

1. -2

2. $-\frac{1}{24}$

3. $\frac{2}{3}$

4. $\sqrt{17}$

5. $\frac{4}{3}$

6. $-\frac{1}{3}$

Problem Set 3.1.1:

1. 7

2. 320

3. 108

4. 24

8. 285

5. 9

6. 315

7. 12

9. 13

10. 364

11. 6

12. 18

13. 108

14. 324

- | | | | |
|---------|----------|---------|---------|
| 15. 4 | 22. 144 | 29. 288 | 36. 63 |
| 16. 102 | 23. 17 | 30. 108 | 37. 144 |
| 17. 216 | 24. -260 | 31. 22 | 38. -76 |
| 18. 84 | 25. 160 | 32. 693 | 39. 14 |
| 19. 432 | 26. 420 | 33. 96 | 40. 720 |
| 20. 420 | 27. 72 | 34. 201 | 41. 360 |
| 21. 11 | 28. 42 | 35. 336 | 42. 168 |

Problem Set 3.1.3:

- | | | | |
|---------|--------|----------|-----------|
| 1. -64 | 6. 4 | 11. 128 | 16. -2 |
| 2. 1728 | 7. 32 | 12. 9 | 17. -3456 |
| 3. 0 | 8. 0 | 13. 128 | 18. 16 |
| 4. 8 | 9. 16 | 14. 1458 | 19. 16000 |
| 5. 64 | 10. 96 | 15. 2500 | |

Problem Set 3.1.4:

- | | | | |
|------------------|--------------------|-------------------|--------------------|
| 1. $\frac{3}{2}$ | 6. $-\frac{3}{4}$ | 11. 0 | 16. -2 |
| 2. 9 | 7. 3 | 12. $\frac{5}{2}$ | 17. $-\frac{1}{4}$ |
| 3. $\frac{2}{3}$ | 8. $-\frac{3}{5}$ | 13. $\frac{3}{5}$ | 18. 2 |
| 4. -7 | 9. $\frac{1}{4}$ | 14. $\frac{1}{6}$ | 19. 7 |
| 5. 0 | 10. $-\frac{2}{3}$ | 15. -3 | 20. -36 |

21. 3

22. $-\frac{3}{4}$

23. $-\frac{1}{4}$

25. -4

24. 0

Problem Set 3.1.5:

1. 9

6. 7

4. 3

9. 6

2. 9

7. -1

5. 3

3. 6

8. -1

Problem Set 3.2.6:

1. 1224

7. 289

13. $\frac{4}{3}$

19. 144

2. 630.9

8. 29.2

14. 0

20. 2

3. $\frac{1}{8}$

9. 2.5

15. -6

21. 10

4. 2

10. 324

16. 343

5. $\frac{2}{7}$

11. 216

17. 13

22. 0

6. $\frac{1}{64}$

12. $2\frac{2}{3}$

18. 4

23. 25

Problem Set 3.1.7:

1. $\frac{1}{9}$

6. $\frac{4}{3}$

11. 3

16. 1

2. 2

7. 9

12. 1

17. 6

3. 2

8. 7

13. 3

18. 2

4. 6

9. -3

14. 1

19. 5

5. -3

10. 1

15. $\frac{8}{3}$

20. 3

- | | | | |
|--------------------|-------------------|--------------------|-------------------|
| 21. 3 | 29. 1 | 37. 0 | 45. 9 |
| 22. $1\frac{1}{2}$ | 30. 4 | 38. 8 | 46. 12 |
| 23. 2 | 31. 2 | 39. 5 | 47. 8 |
| 24. 1 | 32. 0 | 40. (*) 791 – 876 | 48. 1 |
| 25. –.5 | 33. $\frac{3}{2}$ | 41. .5 | 49. 16 |
| 26. 0 | 34. 0 | 42. 2 | 50. $\frac{1}{3}$ |
| 27. $\frac{3}{4}$ | 35. –1.5 | 43. 8 | 51. $\frac{1}{2}$ |
| 28. 243 | 36. 22 | 44. $\frac{1}{16}$ | |

Problem Set 3.1.8:

- | | | | |
|-------|-------|-------|-------------------|
| 1. 45 | 4. 78 | 7. 22 | 10. $\frac{3}{4}$ |
| 2. 60 | 5. 36 | 8. 48 | 11. $\frac{2}{3}$ |
| 3. 66 | 6. 28 | 9. 36 | 12. 40 |

Problem Set 3.1.9:

- | | | | |
|--------------------|--------------------|-------------------|-------------------------|
| 1. (*) 117 – 131 | 7. (*) 2368 – 2618 | 13. (*) 170 – 189 | 19. (*) 831 – 919 |
| 2. 94 | 8. (*) 395 – 438 | 14. (*) 128 – 142 | 20. (*) 270 – 299 |
| 3. (*) 145 – 161 | 9. (*) 2407 – 2661 | 15. (*) 150 – 167 | 21. (*) 296 – 328 |
| 4. (*) 172 – 191 | 10. (*) 887 – 981 | 16. (*) 489 – 541 | 22. (*) 279 – 309 |
| 5. (*) 2430 – 2686 | 11. (*) 496 – 549 | 17. (*) 271 – 301 | 23. (*) 7276 – 8043 |
| 6. 87 | 12. (*) 186 – 207 | 18. (*) 486 – 539 | 24. (*) 200220 – 221297 |

25. (*) $26596 - 29397$ 217917 29. (*) $1258 - 1392$
 26. (*) $197162 -$ 27. (*) $217 - 241$ 28. (*) $62366 - 68932$

Problem Set 3.1.10:

- | | | | |
|------------------|-----------|------------|---------------------|
| 1. 15 | 9. 1600 | 18. 50 | 27. $16 + 16i$ |
| 2. 61 | 10. 1 | 19. -41 | 28. 1 |
| 3. -6 | 11. -44 | 20. 7 | 29. $\frac{12}{13}$ |
| 4. 24 | 12. 31 | 21. 0 | 30. -64 |
| 5. 54 | 13. 9 | 22. 41 | 31. $\frac{1}{5}$ |
| 6. 48 | 14. 15 | 23. 3721 | 32. 625 |
| 7. $\frac{3}{2}$ | 15. 53 | 24. -243 | 33. $\frac{12}{13}$ |
| 8. 25 | 16. -7 | 25. 4 | 34. 169 |
| | 17. -7 | 26. 0 | |

Problem Set 3.1.11:

- | | | | |
|-------------------|-------------------|--------------------|----------|
| 1. $-\frac{4}{3}$ | 7. 1 | 13. 1 | 19. 0 |
| 2. 2.5 | 8. $3\frac{1}{2}$ | 14. -5 | 20. 2 |
| 3. $\frac{1}{3}$ | 9. -3 | 15. $-\frac{2}{3}$ | 21. -1 |
| 4. 3 | 10. -4 | 16. 1 | 22. 7 |
| 5. $-\frac{7}{3}$ | 11. 1 | 17. $\frac{7}{3}$ | 23. 1 |
| 6. $\frac{2}{3}$ | 12. -1 | 18. -2 | 24. 7 |

Problem Set 3.1.12:

- | | | | |
|--------|--------|---------|---------|
| 1. 17 | 5. 217 | 9. 110 | 13. 513 |
| 2. 110 | 6. 720 | 10. 398 | 14. 511 |
| 3. 65 | 7. 26 | 11. 101 | 15. 45 |
| 4. 91 | 8. 256 | 12. 46 | 16. .25 |

Problem Set 3.1.13:

- | | | | |
|--------------------|--------------------|--------------------|---------------------|
| 1. $\frac{1}{12}$ | 9. $\frac{3}{8}$ | 17. $\frac{5}{4}$ | 25. $\frac{7}{36}$ |
| 2. $\frac{1}{18}$ | 10. $\frac{9}{8}$ | 18. $\frac{3}{5}$ | 26. $\frac{1}{2}$ |
| 3. $\frac{3}{18}$ | 11. $\frac{1}{3}$ | 19. $\frac{3}{2}$ | 27. $\frac{1}{3}$ |
| 4. $\frac{1}{7}$ | 12. $\frac{7}{29}$ | 20. $\frac{1}{18}$ | 28. $\frac{5}{6}$ |
| 5. $\frac{9}{13}$ | 13. $\frac{5}{4}$ | 21. $\frac{1}{6}$ | 29. $\frac{3}{4}$ |
| 6. $\frac{4}{5}$ | 14. $\frac{5}{8}$ | 22. $\frac{1}{5}$ | 30. $\frac{1}{5}$ |
| 7. $\frac{13}{20}$ | 15. $\frac{3}{5}$ | 23. $\frac{5}{13}$ | 31. $\frac{1}{169}$ |
| 8. $\frac{1}{216}$ | 16. $\frac{1}{25}$ | 24. $\frac{1}{4}$ | |

Problem Set 3.1.14:

- | | | | |
|-------|------|-------|-------|
| 1. 4 | | 8. 5 | |
| 2. 32 | 5. 1 | 9. 4 | 12. 2 |
| 3. 16 | 6. 3 | 10. 5 | 13. 8 |
| 4. 5 | 7. 4 | 11. 4 | 14. 7 |

15. 5 17. 4 19. 15 21. 1

16. 6 18. 254 20. 15

Problem Set 3.2.1:

1. 57 13. 404 25. 1355 37. 1331
2. 1230 14. 234 26. 9 38. 1414
3. 254 15. 1414 27. 443
4. 103 16. 27 28. 102 39. 1234
5. 102 17. 202 29. 1101 40. 2332
6. 312 18. 3210 30. 1011
7. 1010 19. 2300 31. 2220 41. 5
8. 11000 20. 333 32. 140 42. 32
9. 1010 21. 250 33. 104 43. 100
10. 1210 22. 72 34. 69
11. 110 23. 10101 35. 10101 44. 38
12. 21 24. 1101 36. 1323 45. 25

Problem Set 3.2.2:

1. $\frac{17}{25}$ 8. $\frac{13}{24}$
2. $\frac{19}{25}$ 5. $\frac{69}{125}$ 12. $\frac{24}{25}$
3. $\frac{57}{343}$ 6. $\frac{9}{16}$ 9. $\frac{52}{125}$ 13. $\frac{9}{25}$
4. $\frac{15}{16}$ 7. $\frac{7}{12}$ 10. $\frac{35}{36}$ 14. .21
11. $\frac{124}{125}$

- | | | |
|---------|---------|---------|
| 15. .55 | | 18. .21 |
| | 17. .74 | |
| 16. .33 | | 19. .42 |

Problem Set 3.2.3:

- | | | | |
|--------|----------|----------|----------|
| 1. 120 | 11. -44 | 21. 24 | 31. 64 |
| 2. 340 | 12. 606 | 22. 30 | 32. 1221 |
| 3. 10 | 13. 33 | 23. 32 | |
| 4. 341 | 14. 31 | 24. 142 | 33. 121 |
| 5. 4 | 15. 1102 | 25. 21 | 34. 231 |
| 6. 115 | 16. 210 | 26. 34 | 35. 330 |
| 7. 181 | 17. 121 | 27. 1221 | |
| 8. 12 | 18. 201 | 28. 44 | 36. 222 |
| 9. 35 | 19. 143 | 29. 1331 | 37. 124 |
| 10. 22 | 20. 220 | 30. 31 | 38. 1331 |

Problem Set 3.2.4:

- | | | | |
|----------|-------------|---------------|-----------|
| 1. 1120 | 5. 78 | 9. 101101 | 13. 23 |
| 2. 1122 | 6. 11100101 | 10. 100011010 | |
| 3. 133 | 7. 11011 | 11. 110110 | 14. 11011 |
| 4. 11011 | 8. 223 | 12. 33 | 15. 123 |

Problem Set 3.2.6:

1. $\frac{5}{6}$

4. $\frac{7}{8}$

5. $\frac{1}{4}$

2. 3

3. $\frac{6}{7}$

Problem Set 3.3.2:

1. $\frac{3}{11}$

4. $\frac{9}{11}$

7. $\frac{8}{11}$

10. $\frac{77}{333}$

2. $\frac{41}{99}$

5. $\frac{4}{11}$

8. $\frac{5}{33}$

11. $\frac{101}{333}$

3. $\frac{7}{33}$

6. $\frac{2}{99}$

9. $\frac{308}{999}$

12. $\frac{11}{111}$

Problem Set 3.3.3:

1. $\frac{7}{30}$

4. $\frac{29}{90}$

2. $\frac{29}{90}$

3. $\frac{19}{90}$

5. $\frac{11}{900}$

Problem Set 3.3.4:

1. $\frac{211}{990}$

4. $\frac{151}{494}$

7. $\frac{269}{990}$

10. $\frac{106}{495}$

2. $\frac{61}{495}$

5. $\frac{203}{990}$

8. $\frac{233}{990}$

11. $\frac{61}{495}$

3. $\frac{229}{990}$

6. $\frac{311}{990}$

9. $\frac{47}{990}$

Problem Set 3.4:

1. 3

8. 5

2. 4

5. 2

9. 2

12. 1

3. 1

6. 3

10. 4

13. 3

4. 4

7. 1

11. 2

14. 9

Problem Set 3.5.1:

- | | | |
|---------|----------|--------|
| 1. 719 | | 4. -20 |
| 2. 5039 | 3. 40319 | 5. 152 |

Problem Set 3.5.2:

- | | | | |
|---------------------|-------------------|----------|---------|
| 1. $8\frac{1}{7}$ | 6. 600 | 11. -718 | 16. 35 |
| 2. $10\frac{1}{9}$ | 7. $\frac{1}{15}$ | 12. 32 | 17. -80 |
| 3. $6\frac{5}{6}$ | 8. -12 | 13. 4 | 18. 48 |
| 4. $10\frac{9}{10}$ | 9. -113 | 14. 15 | 19. -9 |
| 5. -100 | 10. 120 | 15. 5 | 20. 60 |

Problem Set 3.5.3:

- | | | | |
|------|------|------|------|
| 1. 0 | | 4. 6 | |
| 2. 6 | 3. 1 | 5. 3 | 6. 0 |

Problem Set 3.6.1:

- | | | | |
|------------------|------|-------|------------------|
| 1. $\frac{3}{7}$ | 3. 4 | 5. 4 | 7. 0 |
| 2. 2 | 4. 3 | 6. 27 | 8. $\frac{9}{2}$ |

Problem Set 3.6.2:

- | | | | |
|--------|-------|-------|-------|
| 1. -11 | 3. 12 | 5. 78 | 7. 24 |
| 2. 2 | 4. 5 | 6. -8 | 8. 60 |

- | | | | |
|---------|---------|---------|----------|
| 9. -32 | 15. 4 | 21. 1 | 27. -224 |
| 10. -52 | 16. 172 | 22. 24 | 28. 84 |
| 11. 18 | 17. 19 | 23. 54 | 29. -2 |
| 12. 2 | 18. 6 | 24. -68 | |
| 13. -15 | 19. 18 | 25. 54 | |
| 14. 14 | 20. -7 | 26. 24 | |

Problem Set 3.6.3:

- | | | | |
|-------------------|--------------------|-------------------|--------------------|
| 1. $8\frac{2}{3}$ | 11. $\frac{3}{2}$ | 21. 4 | 31. $\frac{18}{5}$ |
| 2. 6 | 12. $8\frac{2}{3}$ | 22. 4 | 32. $3\frac{3}{4}$ |
| 3. 15 | 13. 6 | 23. -3 | 33. 4 |
| 4. 18 | 14. $\frac{2}{3}$ | 24. 12 | 34. 6 |
| 5. 4 | 15. $\frac{1}{3}$ | 25. $\frac{3}{5}$ | 35. 2 |
| 6. $\frac{4}{7}$ | 16. $\frac{2}{3}$ | 26. 84 | 36. 3 |
| 7. $4\frac{2}{3}$ | 17. $5\frac{1}{3}$ | 27. 2 | 37. 4 |
| 8. 12 | 18. 6 | 28. $\frac{2}{3}$ | 38. $1\frac{1}{2}$ |
| 9. 2 | 19. 9 | 29. $\frac{3}{4}$ | 39. 6 |
| 10. 0 | 20. 2 | 30. 9 | |