



OKLAHOMA
State Department
of Health

April 8, 2024

Dear Parent/Guardian,

The Bryan County Health Department will be at your child's school on Thursday, April 25, 2024 providing Tetanus, Diphtheria and Pertussis (Tdap) vaccines to 6th grade students. A Tdap vaccine is required for Oklahoma children going into 7th grade. Included in this packet, you will find a vaccine information sheet for additional information regarding Tdap vaccine.

In order for your child to be able to receive a vaccine the following forms must be completed and returned to school: Immunization Data Entry Form (front page only) and Screening Checklist for Contraindications to Vaccines for Children and Teens. Unfortunately, your child will not be immunized if the required paperwork is not completed and returned with parent/guardian signatures.

For students who have private insurance, a claim will be sent to the insurance provider. If you receive a bill, please contact the Bryan County Health Department at (580) 924-4285.

If students are not cooperative, the vaccine will not be administered. Parents/guardians are welcome to be in attendance.

Sincerely,

Amy Potts RN
Coordinating Nurse

Immunizations Data Entry Form

Please complete each field below with the information that applies to the client receiving services today. **If client is a minor, please sign the Consent in both places if parent/legal guardian is not present with minor at visit.**

Last Name		First Name		Middle Initial	Suffix (eg., Jr, III)	Age	Date of Birth
Street Address		City		State	Zip	Phone Number ()) ☐ Cell ☐ Home	
Hispanic Origin ☐ Yes ☐ No	Race ☐ American Indian/Alaskan Native ☐ Asian ☐ Black/African American ☐ Native Hawaiian/Other Pacific Islander ☐ White ☐ Other		Marital Status ☐ Single ☐ Married ☐ Divorced ☐ Widowed ☐ Legally Separated ☐ Unknown		Primary Language ☐ American Sign Language ☐ English ☐ Spanish ☐ Other		
Gender ☐ Female ☐ Male ☐ Other	State of Birth	Country of Birth		May we contact you? ☐ Yes, at address provided ☐ Yes, at phone provided ☐ No			
If the client is under 18 years of age, please complete guardian information.* Guardian Relationship to Client: ☐ Father ☐ Mother ☐ Legal Guardian ☐ Other Guardian Name (Last, First) _____							
Medical Insurance and Household Income Information							
Does client have health insurance? ☐ Yes ☐ No		Annual Household Income \$ _____		Number of people supported by income _____			
☐ SoonerCare/Medicaid	SoonerCare/Medicaid Number		Member First and Last name as it appears on policy or card		Mother's Maiden Name		
☐ Private Insurance	Primary Insurance Co & EDI/Payer ID		Policyholder		Client Relation to Policyholder		Member ID
	Secondary Insurance Co & EDI/Payer ID		Policyholder		Client Relation to Policyholder		Member ID
☐ Medicare	Do you have Medicare Part B? ☐ Yes ☐ No		Is Medicare Primary? ☐ Yes ☐ No		Medicare Number		
Consent for Service							
I, the undersigned, give my consent for the services that I am requesting from the Oklahoma State Department of Health (OSDH) and its entities/contractors. I understand that the risks and benefits for these services will be explained to me and that I will have the opportunity to ask questions. I also understand that: • The information regarding myself and the services I receive will be entered into OSDH management information systems and may be used for program evaluation, management, and billing purposes. • I will not be denied service because of my inability to pay. • I may refuse service at any time.							
AUTHORIZATION FOR RELEASE OF INFORMATION AND ASSIGNMENT OF THIRD-PARTY PAYMENTS: It is ultimately the client's responsibility to know your coverage and benefits. You may be responsible for any amount not covered by your insurance. If your insurance carrier denies any part of your claim, or if you elect to continue services past your coverage/policy period, you may be responsible for your balance in full. • I authorize the OSDH to furnish information to my insurance carrier(s) concerning my care. • I authorize my insurer(s) to pay any benefits directly to OSDH. I understand that any amount remaining after such payment has been made by my insurance carrier becomes my responsibility. • I have read the above policy regarding my financial responsibility to OSDH for providing medical services to me or the above-named patient. I certify that the information is, to the best of my knowledge, true and accurate. • I acknowledge that I have received a copy of the Oklahoma State Department of Health Privacy Statement as required by the Health Information Portability and Accountability Act (HIPAA). I can also find a copy on the agency website.							
Client/Guardian Signature: _____				Date: _____			
*For clients less than 18 years old without a parent/guardian present: I consent for my child/ward to receive the provider's recommended vaccines today without my presence, and understand if my child is not cooperative, vaccines will not be given. In the event of an emergency situation, emergency medications and/or oxygen may be administered to my child.							
Parent/Guardian Signature: _____				Date: _____			

Date of Service: _____

OFFICE USE ONLY – DO NOT WRITE BELOW									
VACCINE NAME: Lot #: Exp. Date:		Site:		VIS given? Reaction?		Y N Y N		Provider Signature:	
VACCINE NAME: Lot #: Exp. Date:		Site:		VIS given? Reaction?		Y N Y N		Provider Signature:	
VACCINE NAME: Lot #: Exp. Date:		Site:		VIS given? Reaction?		Y N Y N		Provider Signature:	
VACCINE NAME: Lot #: Exp. Date:		Site:		VIS given? Reaction?		Y N Y N		Provider Signature:	
VACCINE NAME: Lot #: Exp. Date:		Site:		VIS given? Reaction?		Y N Y N		Provider Signature:	
VACCINE NAME: Lot #: Exp. Date:		Site:		VIS given? Reaction?		Y N Y N		Provider Signature:	
VACCINE NAME: Lot #: Exp. Date:		Site:		VIS given? Reaction?		Y N Y N		Provider Signature:	
VACCINE NAME: Lot #: Exp. Date:		Site:		VIS given? Reaction?		Y N Y N		Provider Signature:	
(Circle one) JSN / MOD / MOD 6mo / MOD 6-11yr / PFR / PFR 6mo / PFR 5-11yr / NVX Lot #: Exp. Date:		Site:		EUA/VIS given? Reaction?		Y N Y N		Provider Signature:	
0 – P.O.		1 – RT VAST LAT IM		2 – LT VAST LAT IM		3 – LT DELTOID IM		4 – RT DELTOID IM	
8 – LT GLUTEUS IM		9 – OTHER		10 – RT LEG		11 – LT LEG		12 – RT DELT REG ID	
16 – RT LOW FOREARM ID		17 – LT FOREARM ID		18 – LT LOW FOREARM ID		13 – LT DELT REG ID		5 – RT UPPER ARM SQ	
								6 – LT UPPER ARM SQ	
								7 – RT GLUTEUS IM	
								14 – NASAL	
								15 – RT FOREARM ID	

Immunizations BCD for **Monkeypox Vaccine only** --- Jynneos SDV/MDV _____ (Service Code ID 3289)
(Provider initials)

Screening Checklist for Contraindications to Vaccines for Children and Teens

PATIENT NAME _____

DATE OF BIRTH ____/____/____
month day year

For parents/guardians: The following questions will help us determine which vaccines your child may be given today. If you answer "yes" to any question, it does not necessarily mean your child should not be vaccinated. It just means additional questions must be asked. If a question is not clear, please ask your healthcare provider to explain it.

	yes	no	don't know
1. Is the child sick today?	☐	☐	☐
2. Does the child have allergies to medicine, food, a vaccine component, or latex?	☐	☐	☐
3. Has the child had a serious reaction to a vaccine in the past?	☐	☐	☐
4. Does the child have a long-term health problem with heart, lung (including asthma), kidney, liver, nervous system, or metabolic disease (e.g., diabetes), a blood disorder, no spleen, a cochlear implant, or a spinal fluid leak? Are they taking regular aspirin or salicylate medication?	☐	☐	☐
5. For children age 2 through 4 years: Has a healthcare provider told you that the child had wheezing or asthma in the past 12 months?	☐	☐	☐
6. For babies: Have you ever been told the child had intussusception?	☐	☐	☐
7. Has the child, a sibling, or a parent had a seizure; has the child had a brain or other nervous system problem?	☐	☐	☐
8. Has the child ever been diagnosed with a heart condition (myocarditis or pericarditis) or have they had Multisystem Inflammatory Syndrome (MIS-C) after an infection with the virus that causes COVID-19?	☐	☐	☐
9. Does the child have an immune-system problem such as cancer, leukemia, HIV/AIDS?	☐	☐	☐
10. In the past 6 months, has the child taken medications that affect the immune system such as prednisone, other steroids, or anticancer drugs; drugs to treat rheumatoid arthritis, Crohn's disease, or psoriasis; or had radiation treatments?	☐	☐	☐
11. Does the child's parent or sibling have an immune system problem?	☐	☐	☐
12. In the past year, has the child received immune (gamma) globulin, blood/blood products, or an antiviral drug?	☐	☐	☐
13. Is the child/teen pregnant?	☐	☐	☐
14. Has the child received vaccinations in the past 4 weeks?	☐	☐	☐
15. Has the child ever felt dizzy or faint before, during, or after a shot?	☐	☐	☐
16. Is the child anxious about getting a shot today?	☐	☐	☐

Did you bring your immunization record card with you? yes ☐ no ☐

It is important to have a personal record of your child's vaccinations. If you don't have one, ask the child's healthcare provider to give you one with all your child's vaccinations on it. Keep it in a safe place and bring it with you every time you seek medical care for your child. Your child will need this document to enter day care or school, for employment, or for international travel.

FORM COMPLETED BY _____ DATE _____

OFFICE USE ONLY: Circle vaccine(s) given.

DTaP HepA HepB Hib HPV IPV MenACWY MenB MMR Pneumococcal Rotavirus Td Tdap Varicella Zostavax Other _____

FORM REVIEWED BY/ PROVIDER SIGNATURE _____ DATE _____



Information for Healthcare Professionals about the Screening Checklist for Contraindications to Vaccines (Children and Teens)

Read the information below for help interpreting answers to the screening checklist.
To learn even more, consult the references in Note below.

NOTE: For additional details, see CDC's "Child and Adolescent Immunization Schedule" (www.cdc.gov/vaccines/schedules/hcp/imz/child-adolescent.html) and General Best Practice Guidelines for Immunization sections on "Contraindications and Precautions" (www.cdc.gov/vaccines/hcp/acip-recs/general-recs/contraindications.html) and "Altered Immunocompetence" (www.cdc.gov/vaccines/hcp/acip-recs/general-recs/immunocompetence.html). For more details on COVID-19 vaccines, see "Use of COVID-19 Vaccines in the United States: Interim Clinical Considerations" at www.cdc.gov/vaccines/covid-19/clinical-considerations/covid-19-vaccines-us.html.

1. Is the child sick today? [all vaccines]

There is no evidence that acute illness reduces vaccine effectiveness or safety. However, as a precaution, all vaccines should be delayed until moderate or severe acute illness has improved. Mild illnesses with or without fever (e.g., otitis media, "colds," and diarrhea) and antibiotic use are not contraindications to routine vaccination.

2. Does the child have allergies to medicine, food, a vaccine component, or latex? [all vaccines]

Gelatin: If a person has anaphylaxis after eating gelatin, do not give vaccines containing gelatin. **Latex:** An anaphylactic reaction to latex is a contraindication to vaccines with latex as part of the vaccine's packaging (e.g., vial stoppers, prefilled syringe plungers, prefilled syringe caps). For details on latex in vaccine packaging, refer to the package insert (listed at www.fda.gov/vaccines-blood-biologics/vaccines/vaccines-licensed-use-united-states). **COVID-19 vaccine:** History of a severe allergic reaction (e.g., anaphylaxis) after a previous dose or to a COVID-19 vaccine component is a contraindication to use of the same vaccine type. People may receive the alternative COVID-19 vaccine type (either mRNA or protein subunit) if they have a contraindication or an allergy-related precaution to one COVID-19 vaccine type. Allergy-related precautions include history of 1) diagnosed non-severe allergy to a COVID-19 vaccine component; 2) non-severe, immediate (onset less than 4 hours) allergic reaction after a dose of one COVID-19 vaccine type (see Note).

Not contraindications: Eggs: ACIP and CDC do not consider egg allergy of any severity to be a contraindication or precaution to any egg-based influenza vaccine. **Injection site reaction** (e.g., soreness, redness, delayed-type local reaction) to a prior dose or vaccine component is not a contraindication to a subsequent dose or vaccine containing that component.

3. Has the child had a serious reaction to a vaccine in the past? [all vaccines]

- Anaphylaxis to a previous vaccine dose or vaccine component is a contraindication for subsequent doses of corresponding vaccines (see question 2).
- Usually, one defers vaccination when a precaution is present, unless the benefit outweighs the risk (e.g., during an outbreak).
- A history of encephalopathy within 7 days of DTP/DTaP is a contraindication for further doses of any pertussis-containing vaccine.
- Other "serious reactions" that this child experienced following vaccination might constitute contraindications or precautions to future doses. See the appendix on vaccine contraindications and precautions in the Note section above.

4. Does the child have a long-term health problem with heart, lung (including asthma), kidney, liver, nervous system, or metabolic disease (e.g., diabetes), a blood disorder, no spleen, a cochlear implant, or a spinal fluid leak? Are they taking regular aspirin or salicylate medication? [MMR, MMRV, LAIV, VAR]

LAIV is not recommended for children with cerebrospinal fluid leak, anatomic or functional asplenia, cochlear implant, a child age 2 through 4 years with a history of asthma or wheezing, or current aspirin or salicylate-containing medication use. Precautions to LAIV include any underlying health condition that increases the risk of influenza complications (see package insert or CDC schedule for details). **MMR & MMRV:** A history of thrombocytopenia or thrombocytopenic purpura is a precaution to MMR and MMRV. **VAR:** Aspirin use is a precaution to VAR due to the association of aspirin use, chickenpox, and Reye syndrome in children and adolescents.

5. For children age 2 through 4 years: Has a healthcare provider told you that the child had wheezing or asthma in the past 12 months? [LAIV]

Children ages 2 through 4 years who had a wheezing episode within the past 12 months should not get LAIV. Give IIV or RIV instead.

6. For babies: Have you ever been told the child had intussusception? [Rotavirus]

Infants who have a history of intussusception (i.e., the telescoping of one portion of the intestine into another) should **not** be given rotavirus vaccine.

7. Has the child, a sibling, or a parent had a seizure; has the child had a brain or other nervous system problem? [DTaP, Td, Tdap, IPV, LAIV, MMRV, RIV]

For patients with stable neurologic disorders (including seizures) unrelated to vaccination, or with a family history of seizures, vaccinate as usual (exception: children with a first degree relative [e.g., parent or sibling] or personal history of seizures generally should receive separate MMR and VAR, not MMRV). **Pertussis-containing vaccines:** DTaP and Tdap are contraindicated in children who have a history of encephalopathy within 7 days

following DTP/DTaP. An unstable progressive neurologic problem is a precaution to using DTaP and Tdap. A history of Guillain-Barré syndrome (GBS): a) Td/Tdap: GBS within 6 weeks of a tetanus-toxoid vaccine is a precaution; if the decision is made to vaccinate, give Tdap instead of Td; b) all influenza vaccines: GBS within 6 weeks of an influenza vaccine is a precaution; influenza vaccination should generally be avoided unless the benefits outweigh the risks (e.g., for those at higher risk for influenza complications).

8. Has the child ever been diagnosed with a heart condition (myocarditis or pericarditis) or have they had Multisystem Inflammatory Syndrome (MIS-C) after an infection with the virus that causes COVID-19?

Precautions to COVID-19 vaccination include a history of myocarditis or pericarditis within 3 weeks after a dose of any COVID-19 vaccine or a history of Multisystem Inflammatory Syndrome (MIS-C). Myocarditis or pericarditis within 3 weeks after a dose of any COVID-19 vaccine is a precaution: the person should generally not receive additional COVID-19 vaccine. A child with a history of myocarditis or pericarditis unrelated to vaccination may receive a COVID-19 vaccine once the condition has completely resolved. A child with a history of MIS-C may be vaccinated if the condition has fully resolved and it has been at least 90 days since diagnosis. Refer to CDC COVID-19 vaccine guidance for additional considerations for myocarditis, pericarditis, and MIS (see Note).

9. Does the child have an immune-system problem, such as cancer, leukemia, HIV/AIDS? [LAIV, MMR, MMRV, Rotavirus, VAR]

Live virus vaccines are usually contraindicated in immunocompromised people with exceptions. For example, MMR is recommended for asymptomatic HIV-infected patients who are not severely immunosuppressed. VAR should be administered (if indicated) to people with isolated humoral immunodeficiency. LAIV is contraindicated in immunosuppressed people; give IIV or RIV instead. Infants with severe combined immunodeficiency (SCID) should not be given a live virus vaccine, including rotavirus vaccine, but other forms of immunosuppression are a precaution, not a contraindication, to rotavirus vaccine. See "General Best Practice Guidelines: Altered Immunocompetence" at www.cdc.gov/vaccines/hcp/acip-recs/general-recs/immunocompetence.html.

10. In the past 6 months, has the child taken medications that affect the immune system such as prednisone, other steroids, or anticancer drugs; drugs to treat rheumatoid arthritis, Crohn's disease, or psoriasis; or had radiation treatments? [LAIV, MMR, MMRV, VAR]

Live virus vaccines should be postponed until after chemotherapy or long-term high-dose steroid therapy has ended. See Note above. Some immune mediator and modulator drugs (especially anti-necrosis factor [TNF] agents) may be immunosuppressive. Avoid live virus vaccines in people taking immunosuppressive drugs. A list of these is in CDC's *Yellow Book* at www.wnc.cdc.gov/travel/yellowbook/2024/additional-considerations/immunocompromised-travelers.

11. Does the child's parent or sibling have an immune system problem? [MMR, MMRV, VAR]

MMR, MMRV, and VAR vaccines should not be given to a patient with a family history of congenital or hereditary immunodeficiency in first-degree relatives (e.g., parents, siblings) unless the patient's immune competence has been verified clinically or by a laboratory.

12. In the past year, has the child received immune (gamma) globulin, blood/blood products, or an antiviral drug? [MMR, MMRV, LAIV, VAR]

See Note (schedule) for antiviral drug information (VAR, LAIV). See "Timing and Spacing of Immunobiologics" (www.cdc.gov/vaccines/hcp/acip-recs/general-recs/timing.html#antibody) for intervals between MMR, VAR, and certain blood/blood products, immune globulin.

13. Is the child/teen pregnant? [HPV, IPV, LAIV, MenB, MMR, MMRV, VAR]

Live virus vaccines (e.g., LAIV, MMR, MMRV, VAR) are contraindicated in pregnancy due to the theoretical risk of virus transmission to the fetus. People who could become pregnant and receive a live virus vaccine should be instructed to avoid pregnancy for 1 month after vaccination. IPV and MenB should not be given except to those with an elevated risk of exposure during pregnancy. HepB: HepBisav-B and PreHevbrio are not recommended during pregnancy, use Engerix-B or Recombivax-HB. HPV is not recommended during pregnancy.

14. Has the child received vaccinations in the past 4 weeks? [LAIV, MMR, MMRV, VAR, yellow fever]

Children given live virus vaccines, such as those listed above, should wait 28 days before receiving another live virus vaccine (wait 30 days for yellow fever vaccine). Inactivated vaccines may be given at the same time or at any spacing interval.

15. Has the child ever felt dizzy or faint before, during or after a shot?

Fainting (syncope) or dizziness is not a contraindication or precaution to vaccination; it may be an anxiety-related response to any injection. CDC recommends vaccine providers consider observing all patients for 15 minutes after vaccination. See Immunize.org's resource on vaccination and syncope at www.immunize.org/catg.d/p4260.pdf.

16. Is the child anxious about getting a shot today?

Anxiety can lead to vaccine avoidance. Simple steps can ease a patient's anxiety about vaccination. Visit Immunize.org's "Addressing Vaccination Anxiety" clinical resources at www.immunize.org/clinical/topic/addressing-anxiety/

VACCINE ABBREVIATIONS

DTaP = Diphtheria, tetanus, & acellular pertussis vaccine
HPV = Human papillomavirus vaccine
IIV = Inactivated influenza vaccine
cclIV - cell culture inactivated influenza vaccine

IPV = Inactivated poliovirus vaccine
LAIV = Live attenuated influenza vaccine
MenB = Meningococcal B vaccine
MMR = Measles, mumps, and rubella vaccine

MMRV = MMR+VAR vaccine
RIV = Recombinant influenza vaccine
Td, Tdap = Tetanus, diphtheria, (acellular pertussis) vaccine
VAR = Varicella vaccine

Tdap (Tetanus, Diphtheria, Pertussis) Vaccine: *What You Need to Know*

Many vaccine information statements are available in Spanish and other languages. See www.immunize.org/vis

Hojas de información sobre vacunas están disponibles en español y en muchos otros idiomas. Visite www.immunize.org/vis

1. Why get vaccinated?

Tdap vaccine can prevent **tetanus, diphtheria, and pertussis**.

Diphtheria and pertussis spread from person to person. Tetanus enters the body through cuts or wounds.

- **TETANUS (T)** causes painful stiffening of the muscles. Tetanus can lead to serious health problems, including being unable to open the mouth, having trouble swallowing and breathing, or death.
- **DIPHTHERIA (D)** can lead to difficulty breathing, heart failure, paralysis, or death.
- **PERTUSSIS (aP)**, also known as “whooping cough,” can cause uncontrollable, violent coughing that makes it hard to breathe, eat, or drink. Pertussis can be extremely serious especially in babies and young children, causing pneumonia, convulsions, brain damage, or death. In teens and adults, it can cause weight loss, loss of bladder control, passing out, and rib fractures from severe coughing.

2. Tdap vaccine

Tdap is only for children 7 years and older, adolescents, and adults.

Adolescents should receive a single dose of Tdap, preferably at age 11 or 12 years.

Pregnant people should get a dose of Tdap during every pregnancy, preferably during the early part of the third trimester, to help protect the newborn from pertussis. Infants are most at risk for severe, life-threatening complications from pertussis.

Adults who have never received Tdap should get a dose of Tdap.

Also, **adults should receive a booster dose of either Tdap or Td** (a different vaccine that protects against tetanus and diphtheria but not pertussis) **every 10 years**, or after 5 years in the case of a severe or dirty wound or burn.

Tdap may be given at the same time as other vaccines.

3. Talk with your health care provider

Tell your vaccination provider if the person getting the vaccine:

- Has had an **allergic reaction after a previous dose of any vaccine that protects against tetanus, diphtheria, or pertussis**, or has any severe, life-threatening allergies
- Has had a **coma, decreased level of consciousness, or prolonged seizures within 7 days after a previous dose of any pertussis vaccine (DTP, DTaP, or Tdap)**
- Has **seizures or another nervous system problem**
- Has ever had **Guillain-Barré Syndrome** (also called “GBS”)
- Has had **severe pain or swelling after a previous dose of any vaccine that protects against tetanus or diphtheria**

In some cases, your health care provider may decide to postpone Tdap vaccination until a future visit.

People with minor illnesses, such as a cold, may be vaccinated. People who are moderately or severely ill should usually wait until they recover before getting Tdap vaccine.

Your health care provider can give you more information.



U.S. Department of
Health and Human Services
Centers for Disease
Control and Prevention

4. Risks of a vaccine reaction

- Pain, redness, or swelling where the shot was given, mild fever, headache, feeling tired, and nausea, vomiting, diarrhea, or stomachache sometimes happen after Tdap vaccination.

People sometimes faint after medical procedures, including vaccination. Tell your provider if you feel dizzy or have vision changes or ringing in the ears.

As with any medicine, there is a very remote chance of a vaccine causing a severe allergic reaction, other serious injury, or death.

5. What if there is a serious problem?

An allergic reaction could occur after the vaccinated person leaves the clinic. If you see signs of a severe allergic reaction (hives, swelling of the face and throat, difficulty breathing, a fast heartbeat, dizziness, or weakness), call 9-1-1 and get the person to the nearest hospital.

For other signs that concern you, call your health care provider.

Adverse reactions should be reported to the Vaccine Adverse Event Reporting System (VAERS). Your health care provider will usually file this report, or you can do it yourself. Visit the VAERS website at www.vaers.hhs.gov or call 1-800-822-7967. *VAERS is only for reporting reactions, and VAERS staff members do not give medical advice.*

6. The National Vaccine Injury Compensation Program

The National Vaccine Injury Compensation Program (VICP) is a federal program that was created to compensate people who may have been injured by certain vaccines. Claims regarding alleged injury or death due to vaccination have a time limit for filing, which may be as short as two years. Visit the VICP website at www.hrsa.gov/vaccinecompensation or call 1-800-338-2382 to learn about the program and about filing a claim.

7. How can I learn more?

- Ask your health care provider.
- Call your local or state health department.
- Visit the website of the Food and Drug Administration (FDA) for vaccine package inserts and additional information at www.fda.gov/vaccines-blood-biologics/vaccines.
- Contact the Centers for Disease Control and Prevention (CDC):
 - Call 1-800-232-4636 (1-800-CDC-INFO) or
 - Visit CDC's website at www.cdc.gov/vaccines.

