Algebra II Summer Assignment Mini Lessons Table of contents

- A. Given the Slope and the Y-Intercept, Write Equation of the Line page 2
 - B. Writing an Equation of a Line Given a point and the Slope page 3
 - C. Given Two Points, Write the Equation of the Line page 4
 - D. Solving Literal Equations page 5
 - E. Changing Equations into Slope Intercept Form page 6
 - F. Graphing from Slope-Intercept Form page 7
 - G. Graphing Systems of Equations page 8
 - H. Solving Systems Using Substitution page 9
 - I. Solving Systems of Equations by Linear Combination page 10
 - J. Word Problem Applications of Linear Systems page 11
 - K. Product of Powers Property of Exponents page 12
 - L. Quotient of Powers Property of Exponents pages 12-13
 - M. Power of a Power Property of Exponents pages 13-14
 - N. Zero and Negative Exponents pages 14-15
 - O. Scientific Notation page 16
 - P. Direct Variation page 17
 - Q. Solving Inequalities page 18
 - R. Two-Variable Inequalities page 19
 - S. Adding & Subtracting Polynomials page 20
 - T. Multiplying Binomials page 21
 - U. Function Notation page 22
 - V. Translations of Graphs Review page 23
 - W. Absolute Value Equations and Inequalities page 24
 - X. Changing Forms of Linear Equations page 25
 - Y. Practice with non-calculator computations page 26

A. Given the Slope and the Y-Intercept,

Write the Equation of the Line

GOAL

Use the slope-intercept form to write an equation of a line

VOCABULARY

In the **slope-intercept form** of the equation of a line, y = mx + b, m is the slope and b is the y-intercept.

Example 1 Writing an Equation of a Line

Write an equation of the line whose slope is 4 and whose y-intercept is -3.

SOLUTION

y = mx + b	Write in slope - intercept form.
y = 4x + (-3)	Substitute 4 for m, and - 3 for b.
y = 4x - 3	Simplify.

Web Enhancement

Refer to the following sites for help with this concept.

This site will explain the concept in greater detail.

http://www.sparknotes.com/math/algebra1/writingequations/section1.html

This site will give examples to practice with answers to check.

http://www.algebralab.net/practice/practice.aspx?file=Algebra1_5-1.xml

B. Writing an Equation of a Line Given a point and the Slope

GOAL

Use slope and any point on a line to write an equation of the line

Example 1 Writing an Equation of a Line

Write an equation of the line that passes through the point (--2, 5) and has a slope of 3.

SOLUTION

Find the y-intercept.

y = mx + b	Write in slope - intercept form.
5 = 3(-2) + b	Substitute 3 for m, - 2 for x, and 5 for y.
5 = -6 + b	Simplify.
11 = b	Solve for b.

The y-intercept is b = 11.

Now write an equation of the line, using slope-intercept form.

y = mx + b	Write the slope - intercept form.
y = 3x + 11	Substitute 3 for m and 11 for b.

Web Enhancement

This site gives directions and an example of writing the equation given a slope and a point.

http://www.purplemath.com/modules/strtlneq.htm

This site will give examples to practice with answers to check.

http://www.algebralab.net/practice/practice.aspx?file=Algebra1_5-1.xml

C. Given Two Points, Write the Equation of the Line

GOAL

Write an equation of a line given two points on the line

Example 1 Writing an Equation Given Two Points

Write an equation of the line that passes through the points (1, 5) and (2, 3).

SOLUTION

Find the slope of the line. Let $(x_1, y_1) = (1, 5)$ and $(x_2, y_2) = (2, 3)$.

$m = \frac{y_2 - y_1}{x_2 - x_1}$	Write formula for slope.
$m = \frac{3-5}{2-1}$	Substitute.
$m = \frac{-2}{1} = -2$	Simplify.

Find the y-intercept. Let m = -2, x = 1, and y = 5 and solve for b. (You could use either point.)

y = mx + b	Write the slope - intercept form.
5 = (-2)(1) + b	Substitute -2 for m, 1 for x, and 5 for y.
5 = -2 + b	Simplify.
7 = b	Solve for b.

Write an equation of the line.

y = mx + b Write slope - intercept form. y = -2x + 7 Substitute - 2 for m and 7 for b.

Web Enhancement This site will take you step by step through examples.

http://webmath.com/equline1.html

D. Solving Literal Equations

GOAL To solve an equation for one variable in terms of other variables.

Equations with several variables (letters) are called literal equations.

Solving a literal equation means that you solve the equation for one of the variables.

Example 1 Solve d = rt for r.

$$\frac{d}{t} = rt$$
$$\frac{d}{t} = \frac{rt}{t}$$
$$\frac{d}{t} = r$$

Example 2 Solve P = 2l + 2w for w.

$$P = 2l + 2w$$

$$P - 2l = 2w$$

$$\frac{P - 2l}{2} = \frac{2w}{2}$$

$$\frac{P - 2l}{2} = w$$

Web Enhancement This site will give further examples of solving literal equations.

http://www.purplemath.com/modules/solvelit.htm

E. Changing Equations into Slope Intercept Form

GOAL

Graph a linear equation in slope-intercept form and interpret equations in slope-intercept form.

VOCABULARY

The linear equation y = mx + b is written in **slope-intercept form**. The slope of the line is m. The y-intercept is b.

Example 1 Writing Equations in Slope-Intercept Form

Example	Equation	Slope-Intercept Form	Slope	y-Intercept
А	y = 3x	y = 3x + 0	m = 3	b = 0
В	$y = \frac{2x - 3}{5}$	$y = \frac{2}{5}x - \frac{3}{5}$	$m = \frac{2}{5}$	$b = -\frac{3}{5}$
C	4x + 8y = 24 $8y = 24 - 4x$ $y = 35x$	y = -0.5x + 3	m = -0.5	<i>b</i> = 3

Web Enhancement This site is shows you step-by-step how to convert an equation into slope-intercept form.

http://www.mathops.com/free/a1le015.php

F. Graphing from Slope-Intercept Form

Example 2 Graphing Using Slope and y-Intercept

Graph the equation 5x - y = 3.

SOLUTION

Write the equation in slope-intercept form: y = 5x - 3

Find the slope and the y-intercept:

m = 5 and b = -3

Plot the point (0, b). Draw a slope triangle to locate a second point on the line.

 $m = \frac{5}{1} = \frac{rise}{run}$

Draw a line through the two points.

Web Enhancement

This video will explain how to graph from slope intercept form.

https://www.khanacademy.org/math/algebra/linear-equations-and-inequalitie/graphing-slopeintercept/v/graphing-a-line-in-slope-intercept-form

This site will let you practice graphing

http://www.ltcconline.net/greenl/java/BasicAlgebra/LineGraph/LineGraph.htm



G. Graphing Systems of Equations

OBJECTIVE Solving a system by graphing

VOCABULARY

As you solve a system of equations, remember the following ideas.

- Lines that have the same slopes but different y-intercepts are parallel and will never intersect. These
 systems are inconsistent.
- Lines that have both the same slope and the same y-intercept are the same line and will intersect at every point. These systems are **dependent.**
- Lines that have different slopes will intersect, and the system will have one solution. These systems are independent.



http://www.purplemath.com/modules/systlin2.htm

H. Solving Systems Using Substitution

OBJECTIVE	Solving systems of linear ed	quations by substitution
Example #1	Solve using substitution.	-4x + y = -13
		x-1=y
y = 4x - 13 $y = x - 1$	Rewrite each equat	ion in the form $y = mx + b$
4x - 13 = x - 1	Since both parts equal y, th equal	ey are equal to each other. Set them to each other.
3x = 12 $x = 4$ $y = 4x - 13$	S	olve for x.
y = 4(4) - 13 y = 3	Substitute 4 for <i>x</i> in	either equation. Solve for <i>y</i> .
y = 5 Example #2	Solve using substitution.	-4x + y = -13 $x - 1 = y$
$\begin{aligned} x - 1 &= y \\ x &= y + 1 \end{aligned}$	Solve one equation fo	r one of the variables. In this example solve e second equation for x.
-4x + y = -13 -4(y + 1) + y =	Substitute what x e^{-13}	equals in the other equation in place of x.
-4y - 4 + y = $-3y = -9$ $y = 3$	-13 D	istribute and solve for y.
x = y + 1 x = (3) + 1 x = 4	Substitute 3 f	for <i>y</i> in either equation. Solve for <i>x</i> .

Web Enhancement This site shows several examples of using substitution to solve systems of equations.

http://www.purplemath.com/modules/systlin4.htm

I. Solving Systems of Equations by Linear Combination

OBJECTIVE Using elimination (linear combination) to solve a system of equations

Instructions Follow these steps when using elimination to solve steps.

- **Step 1:** Arrange the equations with like terms in columns.
- Step 2: Circle the like terms for which you want to obtain coefficients that are opposites.
- **Step 3:** Multiply each term of one or both equations by an appropriate number.
- **Step 4:** Add the equations.
- **Step 5:** Solve for the remaining variable.
- **Step 6:** Substitute the value obtained in step 5 into either of the original equations, and solve for the other variable.
- **Step 7:** Check the solution in the other original equation.

Example Solve the system using the elimination method. $\begin{cases} 2x + 5y = 11 \\ 3x - 2y = -12 \end{cases}$ Highlight the terms that you want to make opposite.

6x + 15y = 33	Multiply each term of the first equation by 3.
-6x + 4y = 24	Multiply each term of the second equation by -2 .
19y = 57 $y = 3$	Add the equations. Solve for the remaining variable.
3x - 2(3) = -12 x = -2	Substitute 3 for <i>y</i> to solve for <i>x</i> .
2(-2) + 5(3) = 11 - 4 + 15 = 11	Check using the other equation.
11 = 11	

The solution is (-2,3). You can also check the solution by using a graphing calculator.

Web Enhancement This site has examples of solving using linear combination. http://www.algebra-class.com/systems-of-equations.html

J. Word Problem Applications of Linear Systems

OBJECTIVE Writing and solving systems of linear equations

Instructions As you solve multi-step systems of linear equations, remember these strategies:

- Determine which form each equation is in: Ax + By = C or y = mx + b
- If the equations are in the form Ax + By = C and a variable can easily be eliminated, use elimination.
- If the equations are in y = mx + b form, use graphing or substitution.

Example Last year, Zach received \$469.75 in interest from two investments. The interest rates were 7.5% on one account and 8% on the other. If the total amount invested was \$6,000, how much was invested at each rate?

Define the variables; x = investment in first account; y = investment in second account

Relate The total amount invested was \$6,000.

Write out the mathematical steps.

x + y = 6,000		
0.075x + 0.08y = 469.75	Determine the form of each equation: $Ax + By = C$.	
y = -x + 6,000	Since a variable cannot be easily eliminated, rewrite 1 equation in	
	the form $y = mx + b$.	
0.075x + 0.08(-x + 6,000) = 469.75	Substitute $x + 6,000$ for y.	
-0.005x = -10.25	Solve for a	
x = 2050	Solve for x.	
2050 + y = 6,000	Substitute 2050 for x in the first equation and solve for y .	
<i>y</i> = 3950		
The encount increased in the first eccount -	x_{22} \$2,050 and in the second account was \$2,050	

The amount invested in the first account was \$2,050 and in the second account was \$3,950.

Web Enhancement

http://www.purplemath.com/modules/systprob.htm

K. Product of Powers Property of Exponents

OBJECTIVE Multiplying exponential expressions with the same base

VOCABULARY

An exponential expression is an expression in the form a^x .

To multiply exponential expressions with the same base, add the exponents: $a^m \star a^n = a^{m+n}$.

Example

Simplify $4^6 \star 4^3$.

 $4^{6} * 4^{3}$ 4^{6+3}

4⁹

Rewrite as one base with the exponents added. Add the exponents.

So, $4^6 * 4^3 = 4^9$.

Web Enhancement This website reviews some of the exponent rules including multiplication.

http://www.wtamu.edu/academic/anns/mps/math/mathlab/beg_algebra/beg_alg_tut26_exp.htm

L. Quotient of Powers Property of Exponents

OBJECTIVE Applying division properties of exponents

VOCABULARY

To divide powers with the same base, subtract the exponents.

$$\frac{4^5}{4^3} = 4^2$$

Example

Simplify $\frac{5^9}{5^{-6}}$

 $\frac{5^9}{5^{-6}} = 5^{9-(-6)} = 5^{9+6} = 5^{15}$ Subtract the exponents from the original equation.

Web <u>Enhancement</u>

More practice with exponent rules.

http://www.algebralab.org/lessons/lesson.aspx?file=Algebra_ExponentsRules.xml

M. Power of a Power Property of Exponents

OBJECTIVE Using more multiplication properties of exponents

VOCABULARY
To raise a power to a power, multiply the exponents. $(x^3)^2 = x^6$
Every number and variable inside parenthesis is being raised to the power to the right of the parenthesis. $(4x^3)^2 = (4^2x^{3(2)}) = 16x^6$

Simplify $(4x^3)^2$. Example #1

Rewrite each number and variable with an exponent.
Multiply the exponents.
Simplify.

Example #2	Simplify $4(x^3)^2$.
$4\left(x^{2^{\star}3}\right)$ $4x^{6}$	Rewrite, showing the exponents to be multiplied for the bases that are affected by the exponents. Multiply the exponents.

Web Enhancement This website goes through power to a power.

http://www.purplemath.com/modules/exponent.htm

N. Zero and Negative Exponents

OBJECTIVE Evaluating and simplifying expressions in which zero and negative numbers are used in exponents

VOCABULARY: When a nonzero number *a* has a zero exponent, then $a^0 = 1$.

For any nonzero number *a* and any integer, *n*, $a^{-n} = \frac{1}{a^n}$

For any nonzero numbers *a* and *b* and any integer, *n*, $\left(\frac{a}{b}\right)^{-n} = \frac{a^{-n}}{b^{-n}} = \frac{b^n}{a^n}$

Example Write each expression as an integer or a simple fraction.

Example A $2.7^0 \rightarrow 1$ Rewrite, using the property of zero as an exponent.Example B $5^{-2} \rightarrow \frac{1}{5^2} \rightarrow \frac{1}{25}$ Rewrite as a fraction, using the property of negative exponents.
Then simplify.Example C $\left(\frac{4}{7}\right)^{-2} \rightarrow \frac{7^2}{4^2} \rightarrow \frac{49}{16}$ Distribute the exponent, switch the placement in the fraction, and
simplify.Example D $\frac{1}{5^{-2}} \rightarrow 1(5^2) \rightarrow 25$ Move the amount with the negative exponent to the other side of
the fraction, make the exponent positive and simplify.

Example E	$\frac{(-4)^2}{(-4)^7} = (-4)^{2-7}$	Quotient of Powers Property
	$= (-4)^{-5}$	Simplify.
	$=\frac{1}{(-4)^5}=-\frac{1}{1024}$	Definition of negative exponent
Example F	$(z^4)^{-3} = z^4 \cdot {}^{(-3)}$	Power of a Power Property
	$= z^{-12}$	Simplify.
	$=\frac{1}{z^{12}}$	Definition of negative exponent
Example G	$\left(\frac{3d}{2}\right)^4 = \frac{(3d)^4}{2^4}$	Power of a Quotient Property
	$=\frac{3^4d^4}{2^4}$	Power of a Product Property
	$=\frac{81d^4}{16}$	Simplify.
Example H	$\left(\frac{2x}{3}\right)^{-5} = \frac{(2x)^{-5}}{3^{-5}}$	Power of a Quotient Property
	$=\frac{3^5}{(2x)^5}$	Definition of negative exponent
	$=\frac{3^5}{2^5x^5}$	Power of a Product Property
	$=\frac{243}{32x^5}$	Simplify.

Example Simplify each expression. Use only positive exponents in your final answer.

Web Enhancement This site has a great video showing why the exponents are as they are: http://www.homeschoolmath.net/teaching/negative_zero_exponents.php

O. Scientific Notation

OBJECTIVE Writing numbers in scientific notation

VOCABULARY

To write a number in **scientific notation**, follow these steps:

- Move the decimal to the right of the first integer.
- If the original number is greater than 1, multiply by 10ⁿ, where *n* represents the number of places the decimal was moved to the left.
- If the original number is less than 1, multiply by 10⁻ⁿ, where *n* represents the number of places the decimal was moved to the right.

Example

Write each number in scientific notation.

А	9,040,000,000	Standard Form
	9.040000000	Move the decimal to the left nine places.
	$9.04*10^{9}$	Drop all significant 0's. Multiply by the appropriate power of
		10.
В	0.0000008	Standard Form
	000008.	Move the decimal to the right seven places.
	$8.0*10^{-7}$	Multiply by the appropriate power of 10.

Web Enhancement

Practice converting from standard to scientific notation.

http://janus.astro.umd.edu/astro/scinote/

http://www.purplemath.com/modules/exponent3.htm

P. Direct Variation

OBJECTIVE Writing and interpreting direct variation

VOCABULARY

A linear function defined by an equation of the form y = kx, where $k \neq 0$, represents **direct variation.** The constant, k, the slope of the line, is called the **constant of variation.** The **y intercept is (0,0).**

Given the value of *y* corresponding to a specific value of *x*, you can find the constant of variation, *k*, by substituting the given values of *x* and *y* into the equation $k = \frac{y}{x} \text{ or } \frac{y-0}{x-0}$.

The equation y = kx can be used to find the values of y that correspond to other values of x or vice versa.

Examples Find the value for each direct variation.

A. If y = 5 when x = 2, find y when x = 7.

$$k = \frac{y}{x} = \frac{5}{2}$$
Use y = 5, x = 2, and k = $\frac{7}{x}$ to find the value of k.

$$y = \frac{5}{2}x$$
Now use the form y = kx and $k = \frac{5}{2}$ to write the equation of the direct variation.

 $y = \frac{5}{2}x = \frac{5}{2}(7) = \frac{35}{2} = 17\frac{1}{2}$ To find the value of y when x = 7, replace x with 7 in the direct variation equation and simplify to find y.

B. If
$$y = 6$$
 when $x = -3$, find x when $y = -4$.

$k = \frac{y}{x} = \frac{6}{-3} = -2$	Use y = 6, x = -3 , and k = $\frac{y}{x}$ to find the value of k.
y = -2x	Now use the form $y = kx$ and $k = -2$ to write the equation of the direct
	variation.
-4 = -2x	
2 = x	

Web Enhancement This site explains direct variation and shows an example.

http://www.freemathhelp.com/direct-variation.html

Q. Solving Inequalities

OBJECTIVE Solving one variable inequalities and graphing inequalities on a number line

To solve an inequality, use the techniques used to solve an equation with one difference; when multiplying or dividing each side by a negative number, reverse the inequality. One variable inequalities can be graphed on a number line or a coordinate plane. When graphing on a number line the equal sign in combination with the inequality means a closed circle.

Examples Solve each inequality. Graph the solutions.

a. $2x-5 \ge 13$ b. 4+3(1-2x) > 37

Use the properties of real numbers and the properties of inequalities to rewrite each inequality in an equivalent form.

When dividing each side by a positive number, do not reverse the inequality.



When dividing (or multiplying) each side by a negative number, reverse the inequality sign.

4 + 3(1 - 2x) > 37 4 + 3 - 6x > 37	Use the distributive property.
7 - 6x > 37	Simplify.
-6x > 30	Subtract 7 from each side.
<i>x</i> < -5	Divide each side by -6 and reverse the inequality.
-10 -9 -8 -7 -	6 -5 -4 -3 -2 -1 0 +1 +2 +3 +4 +5 +6 +7 +8 · -• + + + + + + + + + + + + + + + + +

Web Enhancement

http://www.algebralab.org/lessons/lesson.aspx?file=Algebra_OneVariableSolvingInequalities.x ml

R. Two-Variable Inequalities

OBJECTIVE Graphing inequalities with two variables on a coordinate plane

Examples Graph the inequality $6x - 2y \le 12$.



To graph the boundary line, write the inequality in slope-intercept form as if it were an equation.

The boundary line is solid if the inequality contains \leq or \geq . The boundary line is dashed if the inequality contains < or >. Graph the boundary line y = 3x - 6 as a solid line.





Since the boundary line does not contain the origin, substitute the point (0,0) into the inequality. Simplify. The resulting inequality is true.

Use your highlighting marker to shade the region that contains the origin. If the resulting inequality were false, then you would shade the region that does not contain the origin.

Web Enhancement Link for more step by step instructions or watch the video from the second site listed.

http://www.purplemath.com/modules/ineqgrph.htm

http://www.brightstorm.com/math/algebra/solving-and-graphing-inequalities/graphing-2-variable-inequalities/

S. Adding & Subtracting Polynomials

OBJECTIVE Adding and subtracting polynomials

Vocabulary Term- each part of the polynomial that is being added

Like terms- terms that contain the same variables raised to the same power; only the numerical coefficients are or may be different.

When adding and subtracting polynomials only like terms can be added or subtracted. In an expression, only like terms can be combined. We combine like terms to shorten and simplify algebraic expressions, so we can work with them more easily. To combine like terms, we add the coefficients and keep the variables the same.

Example Add these polynomials: $(2a^2 + 4a - 6) + (a^2 - 2a + 4)$

$(2a^{2} + 4a - 6) + (a^{2} - 2a + 4)$ $(2a^{2} + a^{2}) + (4a - 2a) + (-6 + 4)$ $3a^{2} + 2a - 2$	Put like terms together and add. Be sure and bring the sign with the term.
Subtract these polynomials: $(4x^2 + 2x - 7) - (4x^2 + 2x - 7) = (4x^2 + 2x - 7) - (4x^2 + 2x - 7) = $	$-3x^2-6x+2\Big)$
$(4x^{2} + 2x - 7) - (-3x^{2} - 6x + 2)$ $(4x^{2} - (-3x^{2})) + (2x - (-6x)) + (-7 - 2)$ $7x^{2} + 8x - 9$	Put like terms together and subtract each pair. Be sure and bring the sign with the term.

Web Enhancement The first site describes like terms. http://www.math.com/school/subject2/lessons/S2U2L4DP.html

This website describes adding and subtracting polynomials.

http://www.purplemath.com/modules/polyadd.htm

T. Multiplying Binomials

OBJECTIVE

Vocabulary :

Multiplying binomials

Multiply each term in one binomial by each term of the other binomial. Combine like terms. (FOIL or "sneaky squares")

To multiply two binomials, follow these steps:

Example Find the product (x + 7)(x + 2).

(x+7)(x+2)	Identify the first term of the first binomial
$x^2 + 2x$	Multiply each term of the second binomial by that first term of the first binomial.
(x+7)(x+2)	Identify the second term of the first binomial
7x + 14	Multiply each term of the second binomial by that second term of the first binomial.
$x^{2} + 2x + 7x + 14$	Add the two expressions. Combine like terms.
$x^{2} + 9x + 14$	Solution
	OR you may use "sneaky squares".

	Х	7	$x^{2} + 2x + 7x + 14$
Х	x ²	7x	2 0 11
2	2x	14	$x^2 + 9x + 14$

Web Enhancement These sites show binomial multiplication by FOIL.

http://www.freemathhelp.com/using-foil.html

https://www.khanacademy.org/math/algebra/multiplying-factoring-expression/multiplying-binomials/v/multiplying-binomials

U. Function Notation

OBJECTIVE

Apply function notation to evaluate expressions

Function notation replaces y with f(x). We can identify which equation was used to find the y-value and which input was used as well. Equations are commonly named as f(x), but other letters can be used.

Example Evaluate f(5) for f(x) = 3x + 4.

This means that we are finding the y value when x equals 5. We start by replacing x with 5. Then we simplify.

$$f(5) = 3(5) + 4 = 15 + 4 = 19$$

Example Evaluate g(-3) for $g(x) = x^2 - 8$.

It is important to use parentheses when replacing x with -3. Also note that $(-3)^2$ is positive 9.

$$g(-3) = (-3)^2 - 8 = 9 - 8 = 1$$

Web Enhancement These sites show more function notation examples.

http://www.purplemath.com/modules/fcnnot.htm

http://www.virtualnerd.com/algebra-1/relations-functions/functions/function-notation/f-of-x-definition

V. Translations of Graphs Review

OBJECTIVE

To graph equations by translating parent graphs

Examples



Here is the graph of $y = x^2$ and $y = (x-(-2))^2$. The parent graph is $y = x^2$ has been translated 2 units to the left by subtracting a negative number from x in the function. This can be written as $y = (x+2)^2$.

Here is the graph of y = |x| and y = |x|-3

The parent graph is y = |x| has been translate 3 units down by adding a negative number to the entire function.





Here is the graph of $y = x^2$ and $y = (x+5)^2+2$

The parent graph is $y = x^2$ has been translate 5 units to the left by subtracting a negative number from x in the function and up 2 units by adding 2 to the entire function.

Here is the graph of y = |x| and y = -|x|-2

The parent graph is y = |x| has been reflected over the x axis by changing the sign of the x value and translated 2 units down by adding a negative number to the entire function.





Here is the graph of $y = \sqrt{x}$ and $y = -\sqrt{(x-1)} + 3$. The parent graph is $y = \sqrt{x}$ has been reflected over the x axis by changing the sign of the x value and translated 1 unit to the right by adding a negative number to the x value and translated 3 units up by adding a positive number to the entire function. You can create an infinite number of functions by translating a parent graph.

http://www.mathsisfun.com/sets/function-transformations.html

W. Absolute Value Equations and Inequalities

OBJECTIVE Solving absolute value equations

Vocabulary For every positive real number, *a*, both *a* and –a satisfy the equation |x| = a. To solve an absolute value equation, first rewrite the equation as an equivalent equation with an absolute value expression on the left side by itself. Then rewrite this equation as a compound equality using the rule that if |x| = a then x = a or x = -a.

Example Solve the equation 2|x-3|+1 = 6x+7. Check for extraneous solutions.

Use the proporties of equality to rewrite the equation as an equivalent equation with the absolute value expression on one side by itself. Then write the equation as a compound equality and solve each resulting equation.

2 x-3 +1 = 6x + 7 2 x-3 = 6x + 6	Subtract 1 from each side.
x-3 = 3x+3	Divide each side by 2.
x-3 = 3x + 3 or $x-3 = -(3x+3)$	Rewrite as a compound equality.
-2x = 6 or x - 3 = -3x - 3	
x = -3 or 4x = 0	Solve each equation.
x = -3 or $x = 0$	

To check for extraneous solutions, substitute each value for x in the original absolute value equation. Any value that does not satisfy the original equation must be discarded.

Check	2 -3-3 +1=6(-3)+7	2 0-3 +1 = 6(0) + 7
	2 -6 +1 = -18+7	2 -3 +1=0+7
	2(6) + 1 = -11	2(3) + 1 = 7
	13≠11	7 = 7
The only solution	n is 0; -3 is an extraneous solution.	

Web Enhancement

http://www.purplemath.com/modules/solveabs.htm

X. Changing Forms of Linear Equations

VocabularyForms of the Linear Function Equationy = b + ax or y = mx + bSlope intercept Form: a or m is the slope and b is the y intercept $y = m(x - x_1) + y_1$ Point slope Form: (x_1, y_1) is a point on the graph and m is the slopeAx + By = CStandard or General Form: A, B, C are constants

OBJECTIVE Transform an equation into slope intercept form from standard or pointslope form.

Example Given $y = -\frac{3}{2}(x+1) + 5$ Transform into slope intercept form.

To transform to slope intercept form from point slope form:

$y = -\frac{3}{2}(x+1) + 5$	Distribute the slope.
$y = -\frac{3}{2}x - \frac{3}{2} + 5$	Combine like terms.
$y = -\frac{3}{2}x + 3.5$	This is now slope intercept form with a slope of $-\frac{3}{2}$ and a y intercept of 3.5 or $\frac{7}{2}$

Example Given 3x - 4y = 8 Transform into slope intercept form.

To transform to slope intercept form from standard form:

3x - 4y = 8 $-3x - 3x$	Subtract $3x$ from both sides.
$\frac{-4y = -3x + 8}{-4}$	Divide both sides by -4 .
$y = \frac{3}{4}x - 2$	This is now slope intercept form with a slope of $\frac{3}{4}$ and a y intercept of -2.

Web Enhancement

Refer to the following videos for help with this concept.

http://www.khanacademy.org/video/converting-to-slope-interceptform?playlist=Algebra%20I%20Worked%20Examples

http://www.khanacademy.org/video/point-slope-and-standardform?playlist=Algebra%20I%20Worked%20Examples

Y. Practice with non-calculator computations

Roughly half of the assessments in Algebra 2 are non-calculator. It is important to maintain your computational skills and identify areas where you may need extra practice. These examples let you know the level of computation you should be able to handle without a calculator. If you find that you are struggling after this practice, make sure you get extra help early!

OBJECTIVE Practice non-calculator computational skills

Example Subtract the fractions without using your calculator.

$\frac{7}{8} - \frac{5}{12}$	Identify 24 as the least common denomination (LCD).
$\frac{7}{8} \cdot \frac{3}{3} - \frac{5}{12} \cdot \frac{2}{2}$	Multiply the top and bottom of each term by the number that creates the LCD.
$\frac{21}{24} - \frac{10}{24}$ 21 - 10	Combine the numerators to make one fraction using the LCD.
24	Simplify the numerator.
$\frac{11}{24}$	

Examples Multiply these numbers without a calculator.

