

DIABETES - HYPOGLYCEMIA

Student:	Grade:	_ School Contact:	DOB:	
Mother:	MHome #:	MWork #:	MCell #:	
Father:	FHome #:	FWork #:	FCell #:	
Emergency Contact:	Relation	onship:	Phone:	
SYMPTOMS OF A HYPOGLYCEMIC EF Shaking, fast heartbeat, sweating, anxie Complaints of hunger, impaired vision Onset may be sudden and can prog SEVERE SYMPTOMS INCLUDE: Appears very pale, feels faint, loss of c Seizure activity	ety, irritability , weakness or fatigu ress to Insulin Sh	ie	Student Photo	
STAFF MEMBERS INSTRUCTED: ☐ Administration	☐ Classroom T☐ Support Staf		pecial Area Teacher(s) ransportation Staff	
TREATMENT: Stop any activity immediately. Accompany the student to the Health If off school grounds, provide a source 1/2 - 3/4 cup juice Glucose tabs Hard candy Regular soda (not diet!) Glucose gel Notify parents/guardian (do not delay	e of glucose:	, and the second		
STEPS TO FOLLOW FOR A HYPOGLYO	CEMIC EMERGI	ENCY:		
Glucagon ordered: Yes No If Glucagon is ordered, it should be given by a unconscious, unresponsive or having a seizure. After Glucagon is given, call 911. Notify parer Students receiving glucagon without their parer staff member should accompany the student to present and adequate supervision for other students.	willing volunteer water Preferred Hosport or guardian present the emergency root	ho has been trained by to ital if transported:ent should be transported	d to the hospital by ambulance. A	
Healthcare Provider:	Phone:			
Written by:		Date:		