


Emergency Care Plan



LATEX AND FOOD ALLERGY

Student: _____ Grade: _____ School Contact: _____ DOB: _____
Mother: _____ MHome #: _____ MWork #: _____ MCell #: _____
Father: _____ FHome #: _____ FWork #: _____ FCell #: _____
Emergency Contact: _____ Relationship: _____ Phone: _____

SYMPTOMS OF AN ALLERGIC REACTION MAY INCLUDE ANY/ALL OF THESE:

- **MOUTH** Itching & swelling of lips, tongue or mouth
- **THROAT** Itching, tightness in throat, tightness in chest
- **SKIN** Hives, warmth, itchy rash, generalized swelling
- **STOMACH** Nausea, abdominal cramps, vomiting and/or diarrhea
- **LUNG** Shortness of breath, repetitive cough, wheezing
- **HEART** "Thready pulse", "passing out"

**The severity of symptoms can change quickly –
it is important that treatment is give immediately.**

Student
Photo

STAFF MEMBERS INSTRUCTED:

- ☐ Administration ☐ Classroom Teacher(s) ☐ Special Area Teacher(s)
☐ Support Staff ☐ Transportation Staff

TREATMENT: Rinse contact area with water.

Benadryl ordered: ☐ Yes ☐ No Give _____ Benadryl per provider's orders

Call school nurse at _____. Call parent/guardian if off school grounds.

Epinephrine ordered: ☐ Yes ☐ No Special instructions: _____

IF ANY SYMPTOMS BEYOND REDNESS OR SWELLING ARE SEEN AT THE SITE AND EPINEPHRINE IS ORDERED, GIVE EPINEPHRINE IMMEDIATELY AND CALL 911.

Preferred Hospital if transported: _____

Epinephrine provides a 20 minute response window. After epinephrine, a student may feel dizzy or have an increased heart rate. This is a normal response. Students receiving epinephrine should be transported to the hospital by ambulance. A staff member should accompany the student to the emergency room if the parent, guardian or emergency contact is not present and adequate supervision for other students is present.

Transportation Plan: ☐ Medication available on bus ☐ Medication NOT available on bus ☐ Does not ride bus

Special instructions: _____

Healthcare Provider: _____ Phone: _____

Written by: _____ Date: _____