

ANAPHYLAXIS Individual Health Plan

Name: _____	Birth Date: _____
School: _____	Grade: _____

Severe allergy to: _____

EMERGENCY CARE PLAN

ALLERGY SYMPTOMS



LUNG

Short of breath, wheezing, repetitive cough



HEART

Pale, blue, faint, weak pulse, dizzy



THROAT

Tight, hoarse, trouble breathing/ swallowing



MOUTH

Significant swelling of the tongue and/or lips



SKIN

Many hives over body, widespread redness



GUT

Repetitive vomiting, severe diarrhea



OTHER

Feeling something bad is about to happen, anxiety, confusion

OR A COMBINATION of symptoms from different body areas.

IF YOU SUSPECT AN ALLERGIC REACTION IMMEDIATELY ADMINISTER EPINEPHRINE AS ORDERED BELOW AND CALL 911.

- If student has ingested food item with unknow ingredients – high risk of anaphylaxis – Don't hesitate
- Alert Main Office - Notify administrator of emergency.
- CALL Parent/Guardian. Contact numbers attached.
- May lay student flat, raise legs & keep warm. If breathing is difficult or student is vomiting, lie on side or sit up.
- Consider giving additional ordered medications following Epinephrine (if ordered)

MEDICATION ORDERS: *Can be filled out by licensed health care provider; MD, DO, ND, DMD, DC, PA, ARNP, CNM.*

IHP can be filled out by School Nurse based on BSD "Medication Authorization Form – 3416P Exhibit A" or other completed provider orders.

**In accordance with RCW 18.79.040, if a registered nurse is on site, they may utilize the nursing process, which includes assessment, specialized knowledge, judgment, and skill in determining plan of care or treatment.*

Epinephrine IM autoinjector: 0.3 mg. 0.15 mg.

If symptoms persist, may repeat dose in _____ minutes (may cause increased heart rate, jitteriness, nausea)

EPINEPHRINE is located:

Health Room Classroom Self-Carry*

*The health care provider must provide specific authorization for a student to self-carry.

Is an inhaler required for allergic reaction?**

**If student has asthma which is unrelated to anaphylaxis, please complete/see separate Asthma IHP.

No

Yes

See below for dosing/administration orders.

Inhaler is located:

Health Room Classroom Self-Carry*

*The health care provider must provide specific authorization for a student to self-carry.

Bronchodilator: _____ (Medication name) _____ puffs **after administration of epinephrine.**
(May cause shakiness, jitteriness, increased heart rate)

Other medication authorized to administer *AFTER* epinephrine:

_____ (name) _____ (dose) _____ (route) _____ (frequency)

SELF CARRY, Health Care Provider please check one *(BSD staff has been trained to assist with medication administration):*

YES** student may SELF CARRY. **YES**** student may SELF ADMINISTER.

****Checking "Yes"** indicates that student has been thoroughly instructed in the purpose and appropriate method/frequency of use and/or safe carrying and/or administration of medication. Student/parent/guardian/provider understand the responsibilities of self-carrying and self-administering at school. *Middle and high school students are encouraged to self-carry and self-administer medication as possible.*

NO, student MAY NOT self carry. **NO**, student MAY NOT SELF ADMINISTER.

Licensed Health Care Provider's signature: <small>Signature authorizes medication for length of school year</small>	Date: _____	Phone: _____
	School year: _____	Fax: _____

PARENT/GUARDIAN/STUDENT (age 18 and older) RESPONSIBILITIES:

- I will notify the school nurse of any changes in health status or healthcare provider instructions.
- I will keep track of expiration dates for the medication(s) and replace if expired or used.
- I will provide prescription medication(s) in original pharmacy container labeled with student name, medication name, strength of medication, dose, and time of administration OR I will provide non-prescription medication in the original container with the student name.
- I understand for "self-carrying" medication, back-up medication in the health room is recommended.
- Students who "self-carry" medication are responsible for carrying medication during school and for all school clubs, activities, sports, field trips.
- If my student attends extended day/childcare, clubs before- and after-school, evening, and summer activities, I will notify the program director of my student's medication and health care needs.

Parent/Guardian/Student (age 18 and older) - Please check only one box & sign below:

- I request that I/my student be allowed to **self-carry and/or self-administer this medication.**
- If I do not/my student **DOES NOT** self-carry, I request that the authorized/trained person(s) at school assist me/my student in taking the medicine(s) described above.

If I/my student has permission to SELF-CARRY and/or SELF-ADMINISTER THIS MEDICATION I/my student understands the responsibility of self-carrying medication at school and recognizes the school will not track compliance, expiration, or amount.

I agree to hold harmless and indemnify the school and Bellevue School District's officers, employees, and agents against all claims, judgments, or liabilities arising out of any injury resulting from reasonable and prudent administration of medication, the reasonable performance of health care procedures, or the liabilities arising out of self-administration and self-carrying of medication by my student/me. BSD staff have been trained to assist with medication administration.

I accept this Individual Health Plan. My signature gives permission for exchange of information between the School Nurse and the Health Care Provider regarding this medication order.

Parent/Guardian/Student (age 18) signature: _____	Date: _____	Phone: _____
		Fax: _____

INDIVIDUAL CONSIDERATIONS:

Cafeteria:

- **Elementary School (P-5th grade):** Parent may request the option for their child to eat in an allergen-aware area in the lunchroom.
- **Middle/High School (6th-12th grade):** Student will be making his/her own decision regarding food and seating choices.

Classroom:

- Teacher to inform substitute teachers of the student's Individual Health Plan
- Teachers will notify parent/guardian of classroom food celebrations
- **Elementary School:** Student should not be given food at school unless it is first approved by the parent/guardian
- Store bought, packaged food with clear ingredient list is strongly encouraged for use in food celebrations
- Teachers will notify parent/guardian of instructional materials or classroom projects which contain possible allergens





Field Trip Procedures: If your student does NOT self-carry, medication and IHP will accompany student during any off-campus activity. Student who self-carry are responsible for their own medication.

School Nurse: Will notify teachers, other school staff, transportation, and nutrition services of student's IHP



















Please return to:

School Nurse: _____	Email: _____@bsd405.org	Phone: 425-456-_____
		Fax: 425-456-_____

EPINEPHRINE DIRECTIONS

<p>AUVI-Q (Epinephrine)</p> <ol style="list-style-type: none"> 1. Remove outer case. 2. Pull off red safety guard. 3. Place black end against mid-outer thigh. 4. Press firmly and hold for 2 seconds. 5. Remove from thigh. 	<p>EPIPEN (Epinephrine)</p> <ol style="list-style-type: none"> 1. Remove from carrying case. 2. Pull off blue safety release (with orange tip facing down). 3. Place orange tip against mid-outer thigh. 4. Swing and push firmly until it "clicks" and hold for 3 seconds. 5. Remove from thigh and massage area for 10 seconds. 
<p>ADRENALICK (Epinephrine)</p> <ol style="list-style-type: none"> 1. Remove from carrying case. 2. Pull off grey end caps (with red tip facing down). 3. Place red tip against mid-outer thigh. 4. Press down hard for 10 seconds. 5. Remove from thigh and massage area. 6. Carefully cover the needle with carrying case. 	<p>TEVA (Epinephrine)</p> <ol style="list-style-type: none"> 1. Twist off yellow or green cap 2. Grasp with orange tip facing down. 3. Pull off blue safety release. 4. Swing and push firmly until it "clicks" and hold for 3 seconds. 5. Remove from thigh and massage for 10 seconds 

RESCUE INHALER DIRECTIONS

						
INHALER + SPACER 1. Remove the cap from the mouthpiece. Shake inhaler well.	2. Prime the inhaler if it hasn't been used in: <i>14 days - Albuterol</i> <i>3 days - Xopenex</i> Shake and spray 1 puff into air. Repeat for a total of 4 times.	3. Remove mouthpiece cap from spacer	4. Insert inhaler into the back of the chamber.	4. Insert mouthpiece into mouth or apply facemask to face.	5. Press the inhaler, breathe in closely and hold your breath for 5-10 seconds or breathe 2-3 times (For mask- count for 5-6 breaths). Then remove chamber from mouth and breathe normally.	6. Wait 1 minute. Shake inhaler and repeat steps 4-5 as directed.
						
INHALER 1. Remove the cap from the mouthpiece. Shake inhaler well.	2. Prime the inhaler if it hasn't been used in: <i>14 days - Albuterol</i> <i>3 days - Xopenex</i> Shake and spray 1 puff into air. REPEAT for a total of 4 times.	3. Hold the inhaler with the mouthpiece down.	4. Breathe out through your mouth and push as much air from your lungs as you can.	5. Put the mouthpiece into your mouth and close lips around it. Push the top of the metal canister firmly while you breathe in deeply and slowly through your mouth. Remove from mouth and hold your breath for 10 seconds. Breathe out slowly.	6. Wait 1 minute. Shake and repeat steps 3-5 as directed.	
						
DRY POWDER INHALER 1. Hold the inhaler upright and open the red cap until you feel/hear a "click".	2. Exhale/breathe out through mouth and push as much air from your lungs as possible.	3. Put mouthpiece in mouth and close lips tightly around it. Breathe in quickly and deeply. Remove inhaler from mouth and hold breath for about 10 seconds. Breathe normally.	4. Close red cap firmly over mouthpiece. Repeat steps 1-3 as directed.	5. Wait 1 minute. Repeat steps 1-4 as directed.		

See also BSD Policy/Procedures: 2410 Student Health; 3411 Life Threatening Health Conditions; 3413 Student Immunization and Life-Threatening Health Conditions; 3416 Medication at School; 3418 Emergency Treatment; 3419 Self-Administration of Asthma and Anaphylaxis Medication; 3420 Anaphylaxis Prevention

Other resources: <https://www.aerochambervhc.com/instructions-for-use>; <https://www.lung.org/getmedia/077e6952-e7c3-41ea-8193-73256f5aed37/RespiClick-one-pager.pdf?ext=.pdf>; https://www.nhlbi.nih.gov/sites/default/files/publications/How-to-Use-a-Metered-Dose-Inhaler_21-HL-8165.pdf; <https://www.auvi-q.com/faqs>; <https://www.epipen.com/-/media/files/epipen/howtouseepipenautoinjector.pdf>; <https://epinephrineautoinject.com/what-is-the-epinephrine-injector/how-to-use-the-epinephrine-injector/>; <https://www.tevaepinephrine.com/howtouse>