

SEVERE ALLERGY Care

Child's picture
Face only

This packet MUST be completed, signed, and ATTACHED to an Allergy Medical Action Plan (MAP). Please download, complete the packet and return to the main office.

Student's Name: _____ School: _____
 Date of birth: _____ Age: _____
 Grade: _____ Teacher: _____

This MAP is validated with signatures and dates, by both the licensed health care provider (Doctor of Osteopathic Medicine, D.O., Medical Doctor, M.D., Nurse Practitioner, N.P., or Physician Assistant, P.A.), and a parent/legal guardian. Recommended orders for medical interventions within this treatment plan, will expire at the end of the 2024-2025 school year.

Your child's health care provider will choose to either use their own MAP template, OR the Allergy MAP template within this packet.

Call First:	Call Second:	Call Third:
Name:	Name:	Name:
Relationship:	Relationship:	Relationship:
Phone 1:	Phone 1:	Phone 1:
Phone 2:	Phone 2:	Phone 2:
Email:	Email:	Email:

- YES NO My child has a history of receiving epinephrine for an allergic reaction.
 YES NO My child has asthma (If yes, higher risk for a severe allergic reaction).
 YES NO **REQUEST NO PEANUT OR TREE NUT LUNCH TABLE**

List **ALL** allergies that require a PRESCRIPTION for epinephrine: _____

List all other allergies: _____

PARENT/GUARDIAN CONSENT

I, (parent/guardian), _____, request that my child, _____, receive the attached medical management at school, according to standard school policy. I authorize consent to the ordering licensed health care provider staff and school to share information, as needed, to clarify orders and to assist with my child's health care needs. I agree to have the information, in this entire plan, shared with individuals that need to know. Also, I give permission to use my child's picture on this plan (if I did not supply a photo).

YES NO If my child is to self-carry epinephrine, I will supply the school with a back-up auto-injector.

PARENT/GUARDIAN SIGNATURE: _____ Date: _____

Bus # _____
 Driver: _____
 Route # _____
 Transportation Office Use ONLY if needed
 Medical File _____

Auto-injector type: _____
 School Office Use ONLY
 Expiration date: _____

**PLACE
PICTURE
HERE**

Name: _____ D.O.B.: _____

Allergic to: _____

Weight: _____ lbs. Asthma: **Yes (higher risk for a severe reaction)** **No**

NOTE: Do not depend on antihistamines or inhalers (bronchodilators) to treat a severe reaction. USE EPINEPHRINE.

Extremely reactive to the following allergens: _____








THEREFORE:

If checked, give epinephrine immediately if the allergen was **LIKELY** eaten, for **ANY** symptoms.

If checked, give epinephrine immediately if the allergen was **DEFINITELY** eaten, even if no symptoms are apparent.

FOR ANY OF THE FOLLOWING:





SEVERE SYMPTOMS

 LUNG Shortness of breath, wheezing, repetitive cough	 HEART Pale or bluish skin, faintness, weak pulse, dizziness	 THROAT Tight or hoarse throat, trouble breathing or swallowing	 MOUTH Significant swelling of the tongue or lips
 SKIN Many hives over body, widespread redness	 GUT Repetitive vomiting, severe diarrhea	 OTHER Feeling something bad is about to happen, anxiety, confusion	OR A COMBINATION of symptoms from different body areas.

↓ ↓ ↓

1. **INJECT EPINEPHRINE IMMEDIATELY.**
2. **Call 911.** Tell emergency dispatcher the person is having anaphylaxis and may need epinephrine when emergency responders arrive.
 - Consider giving additional medications following epinephrine:
 - » Antihistamine
 - » Inhaler (bronchodilator) if wheezing
 - Lay the person flat, raise legs and keep warm. If breathing is difficult or they are vomiting, let them sit up or lie on their side.
 - If symptoms do not improve, or symptoms return, more doses of epinephrine can be given about 5 minutes or more after the last dose.
 - Alert emergency contacts.
 - Transport patient to ER, even if symptoms resolve. Patient should remain in ER for at least 4 hours because symptoms may return.

MILD SYMPTOMS

 NOSE Itchy or runny nose, sneezing	 MOUTH Itchy mouth	 SKIN A few hives, mild itch	 GUT Mild nausea or discomfort
--	--	--	--

FOR MILD SYMPTOMS FROM MORE THAN ONE SYSTEM AREA, GIVE EPINEPHRINE.

FOR MILD SYMPTOMS FROM A SINGLE SYSTEM AREA, FOLLOW THE DIRECTIONS BELOW:

1. Antihistamines may be given, if ordered by a healthcare provider.
2. Stay with the person; alert emergency contacts.
3. Watch closely for changes. If symptoms worsen, give epinephrine.

MEDICATIONS/DOSES

Epinephrine Brand or Generic: _____

Epinephrine Dose: 0.1 mg IM 0.15 mg IM 0.3 mg IM


Antihistamine Brand or Generic: _____

Antihistamine Dose: _____

Other (e.g., inhaler-bronchodilator if wheezing): _____


HOW TO USE AUVI-Q® (EPINEPHRINE INJECTION, USP), KALEO

1. Remove Auvi-Q from the outer case. Pull off red safety guard.
2. Place black end of Auvi-Q against the middle of the outer thigh.
3. Press firmly until you hear a click and hiss sound, and hold in place for 2 seconds.
4. Call 911 and get emergency medical help right away.




HOW TO USE EPIPEN®, EPIPEN JR® (EPINEPHRINE) AUTO-INJECTOR AND EPINEPHRINE INJECTION (AUTHORIZED GENERIC OF EPIPEN®), USP AUTO-INJECTOR, MYLAN AUTO-INJECTOR, MYLAN

1. Remove the EpiPen® or EpiPen Jr® Auto-Injector from the clear carrier tube.
2. Grasp the auto-injector in your fist with the orange tip (needle end) pointing downward. With your other hand, remove the blue safety release by pulling straight up.
3. Swing and push the auto-injector firmly into the middle of the outer thigh until it 'clicks'. Hold firmly in place for 3 seconds (count slowly 1, 2, 3).
4. Remove and massage the injection area for 10 seconds. Call 911 and get emergency medical help right away.



HOW TO USE IMPAX EPINEPHRINE INJECTION (AUTHORIZED GENERIC OF ADRENALICK®), USP AUTO-INJECTOR, AMNEAL PHARMACEUTICALS

1. Remove epinephrine auto-injector from its protective carrying case.
2. Pull off both blue end caps: you will now see a red tip. Grasp the auto-injector in your fist with the red tip pointing downward.
3. Put the red tip against the middle of the outer thigh at a 90-degree angle, perpendicular to the thigh. Press down hard and hold firmly against the thigh for approximately 10 seconds.
4. Remove and massage the area for 10 seconds. Call 911 and get emergency medical help right away.



HOW TO USE TEVA'S GENERIC EPIPEN® (EPINEPHRINE INJECTION, USP) AUTO-INJECTOR, TEVA PHARMACEUTICAL INDUSTRIES

1. Quickly twist the yellow or green cap off of the auto-injector in the direction of the "twist arrow" to remove it.
2. Grasp the auto-injector in your fist with the orange tip (needle end) pointing downward. With your other hand, pull off the blue safety release.
3. Place the orange tip against the middle of the outer thigh at a right angle to the thigh.
4. Swing and push the auto-injector firmly into the middle of the outer thigh until it 'clicks'. Hold firmly in place for 3 seconds (count slowly 1, 2, 3).
5. Remove and massage the injection area for 10 seconds. Call 911 and get emergency medical help right away.



HOW TO USE SYMJEPi™ (EPINEPHRINE INJECTION, USP)

1. When ready to inject, pull off cap to expose needle. Do not put finger on top of the device.
2. Hold SYMJEPi by finger grips only and slowly insert the needle into the thigh. SYMJEPi can be injected through clothing if necessary.
3. After needle is in thigh, push the plunger all the way down until it clicks and hold for 2 seconds.
4. Remove the syringe and massage the injection area for 10 seconds. Call 911 and get emergency medical help right away.
5. Once the injection has been administered, using one hand with fingers behind the needle slide safety guard over needle.



ADMINISTRATION AND SAFETY INFORMATION FOR ALL AUTO-INJECTORS:

1. Do not put your thumb, fingers or hand over the tip of the auto-injector or inject into any body part other than mid-outer thigh. In case of accidental injection, go immediately to the nearest emergency room.
2. If administering to a young child, hold their leg firmly in place before and during injection to prevent injuries.
3. Epinephrine can be injected through clothing if needed.
4. Call 911 immediately after injection.

OTHER DIRECTIONS/INFORMATION (may self-carry epinephrine, may self-administer epinephrine, etc.):

Treat the person before calling emergency contacts. The first signs of a reaction can be mild, but symptoms can worsen quickly.

EMERGENCY CONTACTS — CALL 911

RESCUE SQUAD: _____

DOCTOR: _____ PHONE: _____

PARENT/GUARDIAN: _____ PHONE: _____

OTHER EMERGENCY CONTACTS

NAME/RELATIONSHIP: _____ PHONE: _____

NAME/RELATIONSHIP: _____ PHONE: _____

NAME/RELATIONSHIP: _____ PHONE: _____



ROCHESTER COMMUNITY SCHOOLS
 Authorization for Medication Administration
 School Year: 2024-2025

Student name: _____ Date of birth: _____ Grade: _____

To be completed by the Physician or Authorized Prescriber: ONE MEDICATION PER FORM

(Michigan law and district policy require written authorization for a student to take any medication during the school day).

Name of medication: _____ Reason for medication: _____

Dose (*please do not give a range*): _____ MG MG/ML ML MCG UNITS OTHER: _____

Route: Oral Injection Inhalation Intra-nasal Rectal Topical Transdermal (Patch) Other: _____

Routine time(s) to be given: _____ AM _____ PM Other: _____

Frequency: Daily Other (*please be specific*): _____

As needed (PRN), (*absent clear and objective criteria, medication cannot be administered during the school day*):

Special instructions or side effects: _____

Student is both capable and responsible for self-administering this medication (*applicable ONLY to high school students*):

No Yes- supervised Yes- unsupervised

Student may self-carry an inhaler (*applicable to all students*). Yes No Not applicable

Student may self-carry an Epi-Pen (*applicable to all students*). Yes No Not applicable

START: Date from received Other date/duration (please be specific): _____

STOP: End of school year Other date/duration (please be specific): _____

For episodic/emergency events only

Prescriber Name: _____ Signature: _____ Date: _____

Clinic/Hospital Name: _____ Address: _____

Phone number: _____ Fax number: _____

To be completed by Parent/Legal Guardian

I understand and agree that all medication must be in the original container, clearly marked with the student's name, name of medication, and prescribed dosage. I acknowledge that I am required to immediately inform the District of any changes to the healthcare provider's administration instructions. Authorization also includes permission for school personnel and health care provider to contact each other, if needed. I request and authorize the following (*check appropriate direction below*):

School personnel store and administer medication to the above-named student, as authorized by prescriber.

School personnel store medication only. The above-named student shall be responsible for self-administering medication.

Printed Name: _____ Signature: _____ Date: _____



ROCHESTER COMMUNITY SCHOOLS

Medication Procedures (as per standard school policy)

- Medication authorization is for the current school year only and will expire at the end of the school year.
- Only one medication per form. A separate form is required for each medication, each school year.
- Written authorization with medication order completed, signed by the student's authorized healthcare provider and a parent/guardian, is required before any medication can be given at school. Medications include prescription, and non-prescription over-the-counter, including but not limited to: homeopathic, herbal, vitamin, mineral preparation, topical creams or ointments, eye or ear drops, transdermal patches, nasal sprays or mists.
- Medication administration during school hours will be permitted only when failure to do so will jeopardize the health of a student, or the student would not be able to attend school if the medication or treatment were not available during school hours. Parents/legal guardians are urged to administer medication at home and on a schedule, other than school hours, if possible.
- Medication must be brought to school by the student's parent/legal guardian, unless the student has been authorized to self-carry the medication. The district reserves the right to determine that a student may not self-carry for any reason.
- Medication must be administered by an adult in the presence of a second adult, unless the medication is administered by a licensed registered professional nurse or there is an emergency that threatens the student's life or health.
- Parent/legal guardian will ensure that an adequate amount of medication is on hand at the school for the duration of the student's need to take medication, and responsible for checking the need for refills, including expired medications, and replenishing medication to the school in a timely manner.
- All medication must be in a container as prepared by a pharmacy, authorized healthcare provider, or pharmaceutical company, and clearly marked with the student's name, the name of the medication, the prescribed dose, time and frequency of medication administration and special instructions, if any.
- All controlled substance medication will be counted and recorded in the presence of the parent/legal guardian when brought to school.
- Changes in dosage, frequency, or time of administration cannot be made without written instruction from an authorized healthcare provider.
- Designated staff will be responsible for storage, administering medication and notifying parent/legal guardian, in the event that a student refuses medication.
- Medication left over at the end of the school year, or after a student has left the district shall be picked up by the parent/legal guardian. Any medication not retrieved by the parent/legal guardian will be properly disposed of within 7 days of the last student day of school and documented by the individual who is responsible for administering medication.