


## Hyperglycemia (High Blood Glucose)

**Causes:** Too much food, too little insulin or diabetes pills, illness, or stress.

**Onset:** Often starts slowly.



HIGH BLOOD GLUCOSE MAY LEAD TO A MEDICAL EMERGENCY IF NOT TREATED.

## What Can You Do?



**CHECK BLOOD GLUCOSE** 

If your blood glucose levels are higher than your goal for three days and you don't know why,

CALL YOUR
HEALTHCARE PROVIDER



For more information, call the Novo Nordisk Tip Line at 1-800-260-3730 or visit us online at ChangingDiabetes-us.com.

Novo Nordisk Inc. grants permission to reproduce this piece for non-profit educational purposes only on condition that the piece is maintained in its original format and that the copyright notice is displayed. Novo Nordisk Inc. reserves the right to revoke this permission at any time.

Concept developed by Rhonda Rogers, RN, BSN, CDE





## Hypoglycemia (Low Blood Glucose)

Some Symptoms: Causes: Too little food or skipping a meal; too much insulin

or diabetes pills; more active than usual.

Onset: Often sudden.



IF LOW BLOOD GLUCOSE IS LEFT UNTREATED, YOU MAY PASS OUT AND NEED MEDICAL HELP.

## What Can You Do?



**CHECK** your blood glucose, right away. If you can't check, treat anyway.



**TREAT** by eating 3 to 4 glucose tablets or 3 to 5 hard candies you can chew quickly (such as peppermints), or by drinking 4-ounces of fruit juice, or 1/2 can of regular soda pop.



**CHECK** your blood glucose again after 15 minutes. If it is still low, treat again. If symptoms don't stop, call your healthcare provider.

For more information, call the Novo Nordisk Tip Line at 1-800-260-3730 or visit us online at ChangingDiabetes-us.com.

Novo Nordisk Inc. grants permission to reproduce this piece for non-profit educational purposes only on condition that the piece is maintained in its original format and that the copyright notice is displayed. Novo Nordisk Inc. reserves the right to revoke this permission at any time.

Concept developed by Rhonda Rogers, RN, BSN, CDE

