Student Medical Exemption Certificate for Required Immunizations

The information below must be entered in the Department of Public Health Medical Exemptions portal. More

information is available here. Name of Primary Care Provider granting exemption: Please check one (practitioner granting exemption must be licensed as one of the following): ☐ Physician (MD or DO) ☐ Physician Assistant ☐ APRN CT License number: NPI: Phone number: Email: **Directions: Part 1.** Please complete the demographics section on the patient/student. Part 2. Please mark the contraindications/precautions that apply to this patient/student (indicate all that apply). Part 3. If no contraindications or precautions apply in part 2, write a brief explanation of the reason the patient/student requires the exemption. Part 4. Sign the Statement of Clinical Opinion and date the form. **Attach** a copy of the patient/student's most current immunization record. **Part 1.** Patient/Student Information: First name (in full) Middle initial Last name Date of Birth_____ Mailing Address_____City_____ State______Zip____ Parent/Guardian: First Name______Last name_____ Primary phone number _____ School name______ School address City_____ State Zip Current or Grade student is entering ______

Part 2. Please mark the vaccine(s), exemption duration, and all contraindications/precautions that apply to this patient/student for each vaccine.

Medical contraindications and precautions for immunizations are based upon the Advisory Committee on Immunization Practices (ACIP) <u>Comprehensive General Recommendations and Guidelines</u>, published by the Centers for Disease Control and Prevention.

A **contraindication** is a condition in a recipient that increases the risk for a serious vaccine adverse event (VAE) or compromises the ability of the vaccine to produce immunity.

A **precaution** is a condition in a recipient that might increase the risk for a serious VAE or that might compromise the ability of the vaccine to produce immunity. Under normal conditions, vaccinations are deferred when a precaution is self-limiting, but can be administered if the precaution condition improves.

CDC Recognized Contraindications and Precautions

Vaccine	Exemption Duration	ACIP Contraindications and Precautions (Check all that apply)		
☐ Diphtheria- Tetanus-and acellular	☐ Temporary through:	Contraindications		
		☐ Severe allergic reaction (e.g., anaphylaxis) after a previous dose or to a vaccine component		
Pertussis	mm/ yyyy	☐ Encephalopathy within seven days after receipt of previous dose		
(DTaP)	☐ Permanent	of DTP or DTaP		
		Precautions		
		☐ Progressive neurologic disorder, including infantile spasms, uncontrolled epilepsy, progressive encephalopathy: deferDTaP until neurologic status clarified and stabilized		
		☐ GBS <6 weeks after previous dose of tetanus-toxoid—containing vaccine		
		☐ Fever greater than 40.5°C (104.9°F) <48 hours after vaccination of previous dose of DTP or DTaP		
		☐ History of Arthus-type hypersensitivity reactions after a previous dose of diphtheria-toxoid—containing or tetanus-toxoid—containing vaccine; defer vaccination until at least 10 years have elapsed since the last tetanus-toxoid—containing vaccine		
		☐ Moderate or acute illness with or without fever		
☐ Hepatitis A	☐ Temporary through:	Contraindications		
		☐ Severe allergic reaction (e.g., anaphylaxis) after a previous dose		
		or to a vaccine component		
	mm/ yyyy	Precautions		
	☐ Permanent	☐ Moderate or severe acute illness with or without fever		

☐ Hepatitis B ☐ Haemophilus influenzae type b (HiB)	☐ Temporary through: /	 Contraindications □ Severe allergic reaction (e.g., anaphylaxis) after a previous dose or to a vaccine component □ Hypersensitivity to yeast Precautions □ Moderate or severe acute illness with or without fever Contraindications □ Severe allergic reaction (e.g., anaphylaxis) after a previous dose or to a vaccine component
	mm/ yyyy	☐ Age <6 weeks Precautions
		☐ Moderate or severe acute illness with or without fever
☐ Inactivated Influenza Virus (IIV)	☐ Temporary through: / mm/ yyyy ☐ Permanent	Contraindications ☐ Severe allergic reaction (e.g., anaphylaxis) after previous dose of influenza vaccine or to vaccine component Precautions ☐ GBS <6 weeks after a previous dose of influenza vaccine ☐ Moderate or severe acute illness with or without fever ☐ Egg allergy other than hives, e.g., angioedema, respiratory distress, lightheadedness, recurrent emesis; or required epinephrine or another emergency medical intervention (IIV may be administered in an inpatient or outpatient medical setting and under the supervision of a health care provider who is able to recognize and manage severe allergic conditions).
☐ Inactivated Polio Vaccine (IPV)	☐ Temporary through: / mm/ yyyy ☐ Permanent	Contraindications ☐ Severe allergic reaction (e.g., anaphylaxis) after a previous dose or to a vaccine component Precautions ☐ Pregnancy ☐ Moderate or acute illness with or without fever

г

☐ Live	☐ Temporary through:	Contraindications		
Attenuated Influenza Virus (LAIV)		☐ Severe allergic reaction (e.g., anaphylaxis) after a previous dose or to a vaccine component		
(LAIV)	mm/ yyyy ☐ Permanent	☐ Concomitant use of aspirin or aspirin-containing medication in children and adolescents		
		☐ LAIV4 should not be administered to persons who have taken oseltamivir or zanamivir within the previous 48 hours, peramivir within the previous 5 days, or baloxavir within the previous 17 days.(e)		
		☐ Pregnancy		
		☐ Children aged 2 through 4 years who have received a diagnosis of asthma or whose parents or caregivers report that a health care provider has told them during the preceding 12 months that their child had wheezing or asthma or whose medical record indicates a wheezing episode has occurred during the preceding 12 months.		
		☐ Persons with active cerebrospinal fluid/oropharyngeal communications/leaks.		
		☐ Close contacts and caregivers of severely immunosuppressed persons who require a protected environment.		
		☐ Persons with cochlear implants (due to the potential for CSF leak, which might exist for some period of time after implantation. Providers might consider consultation with a specialist concerning risk of persistent CSF leak if an age-appropriate inactivated or recombinant vaccine cannot be used).		
		☐ Altered Immunocompetence		
		☐ Anatomic or functional asplenia (e.g. sickle cell disease)		
		Precautions		
		☐ GBS <6 weeks after a previous dose of influenza vaccine		
		☐ Asthma in persons aged 5 years old or older		
		☐ Medical conditions which might predispose to higher risk of complications attributable to influenza(d)		
		☐ Moderate or severe acute illness with or without fever		
☐ Meningo- coccal conjugate vaccines	☐ Temporary through: / mm/ yyyy	Contraindications ☐ Severe allergic reaction (e.g., anaphylaxis) after a previous dose or to a vaccine component, including yeast Precautions		
(MenACWY)	☐ Permanent	☐ Moderate or severe acute illness with or without fever		

☐ Measles-	☐ Temporary	Contraindications		
Mumps-Rubella (MMR)	through:	Severe allergic reaction (e.g., anaphylaxis) after a previous dose or		
(,	mm/ yyyy	to a vaccine component □ Pregnancy		
		 ☐ Known severe immunodeficiency (e.g., from hematologic and solid tumors, receipt of chemotherapy, congenital immunodeficiency, long-term immunosuppressive therapy (i) or patients with HIV infection who are severely immunocompromised) ☐ Family history of altered immunocompetence (i) 		
		Precautions		
		☐ Recent (≤11 months) receipt of antibody-containing blood product (specific interval depends on product)		
		☐ History of thrombocytopenia or thrombocytopenic purpura		
		☐ Need for tuberculin skin testing or interferon-gamma release assay (IGRA) testing (k)		
		☐ Moderate or severe acute illness with or without fever		
☐ Pneumo-coccal (PCV13)	☐ Temporary through: / mm/ yyyy ☐ Permanent	Contraindications ☐ Severe allergic reaction (e.g., anaphylaxis) after a previous dose of PCV13 or any diphtheria-toxoid—containing vaccine or to a component of a vaccine (PCV13 or any diphtheria-toxoid—containing vaccine), including yeast Precautions ☐ Moderate or acute illness with or without fever		
☐ Tdap	☐ Temporary	Contraindications		
·	through:/_ mm/ yyyy □ Permanent	 □ Severe allergic reaction (e.g., anaphylaxis) after a previous dose or to a vaccine component □ Encephalopathy (e.g., coma, decreased level of consciousness, prolonged seizures), not attributable to another identifiable cause, within 7 days of administration of previous dose of DTP, 		
		DTaP, or Tdap Precautions		
		☐ GBS <6 weeks after a previous dose of tetanus-toxoid—containing		
		vaccine		
		☐ Progressive or unstable neurological disorder, uncontrolled seizures, or progressive encephalopathy until a treatment regimen has been established and the condition has stabilized		
		☐ History of Arthus-type hypersensitivity reactions after a previous		

		dose of diphtheria-toxoid—containing or tetanus-toxoid—containing vaccine; defer vaccination until at least 10 years have elapsed since the last tetanus-toxoid—containing vaccine		
		☐ Moderate or severe acute illness with or without fever		
☐ Varicella	☐ Temporary through:/mm/ yyyy ☐ Permanent	Contraindications ☐ Severe allergic reaction (e.g., anaphylaxis) after a previous dose or to a vaccine component ☐ Known severe immunodeficiency (e.g., from hematologic and solid tumors, receipt of chemotherapy, congenital immunodeficiency, long-term immunosuppressive therapy (i) or patients with HIV infection who are severely immunocompromised) (g) ☐ Pregnancy ☐ Family history of altered immunocompetence (i) Precautions ☐ Recent (≤11 months) receipt of antibody-containing blood product (specific interval depends on product) ☐ Moderate or acute illness with or without fever		
Complete this sec	tion if claiming a n CIP criteria for a co	ther Type of Medical Condition nedical exemption for a vaccine based on a condition that does NOT ontraindication or precaution listed in part 2.		
		t the allergic or other reaction for which medical exemption is being he following that apply:		
☐ This patient h	, ,	of an autoimmune disorder		
•		of a reaction to a vaccination sposition to a reaction to a vaccination as determined through genetic		
☐ This patient h	•	umented reaction that is correlated to a vaccination sted above (must specify):		
Please provide ar	n explanation of th	ne reaction/condition listed above:		

Part 4. Statement of Clinical Opinion

In accord with the legal requirements of Public Act 21-6, the vaccine(s) indicated above is/are in my clinical
opinion medically contraindicated for this patient/student due to the physical condition as explained above

Clinician's Signature		
_		
Date		

A person may be placed into quarantine or isolation when there are "reasonable grounds to believe [a person] to be infected with, or exposed to, a communicable disease or to be contaminated or exposed to contamination or at reasonable risk of having a communicable disease or being contaminated or passing such communicable disease or contamination to other persons if the commissioner determines that such individual or individuals pose a significant threat to the public health and that quarantine or isolation is necessary and the least restrictive alternative to protect or preserve the public health." Conn. Gen. Stat. § 19a-131b(a).