FARE Food Allergy Research & Education

FOOD ALLERGY & ANAPHYLAXIS EMERGENCY CARE PLAN

Name: _____ D.O.B.: _____

Allergic to: _____

Weight: _____ lbs. Asthma: ☐ **Yes (higher risk for a severe reaction)** ☐ **No**

PLACE PICTURE HERE

NOTE: Do not depend on antihistamines or inhalers (bronchodilators) to treat a severe reaction. USE EPINEPHRINE.

Extremely reactive to the following allergens:_____

THEREFORE:

☐ If checked, give epinephrine immediately if the allergen was LIKELY eaten, for ANY symptoms.

☐ If checked, give epinephrine immediately if the allergen was DEFINITELY eaten, even if no symptoms are apparent.

FOR **ANY** OF THE FOLLOWING:

SEVERE SYMPTOMS



LUNG
Shortness of breath, wheezing, repetitive cough

HEART
Pale or bluish skin, faintness, weak pulse, dizziness

THROAT
Tight or hoarse throat, trouble breathing or swallowing

MOUTH
Significant swelling of the tongue or lips

SKIN
Many hives over body, widespread redness

GUT
Repetitive vomiting, severe diarrhea

OTHER
Feeling something bad is about to happen, anxiety, confusion

OR A COMBINATION of symptoms from different body areas.

1. **ADMINISTER EPINEPHRINE IMMEDIATELY.**
2. **Call 911.** Tell emergency dispatcher the person is having anaphylaxis and may need epinephrine when emergency responders arrive.

- Consider giving additional medications following epinephrine:
  - » Antihistamine
  - » Inhaler (bronchodilator) if wheezing
- Lay the person flat, raise legs and keep warm. If breathing is difficult or they are vomiting, let them sit up or lie on their side.
- If symptoms do not improve, or symptoms return, more doses of epinephrine can be given about 5 minutes or more after the last dose.
- Alert emergency contacts.
- Transport patient to ER, even if symptoms resolve. Patient should remain in ER for at least 4 hours because symptoms may return.

MILD SYMPTOMS



NOSE
Itchy or runny nose, sneezing

MOUTH
Itchy mouth

SKIN
A few hives, mild itch

GUT
Mild nausea or discomfort

FOR **MILD SYMPTOMS** FROM **MORE THAN ONE** SYSTEM AREA, GIVE EPINEPHRINE.

FOR **MILD SYMPTOMS** FROM **A SINGLE SYSTEM** AREA, FOLLOW THE DIRECTIONS BELOW:

1. Antihistamines may be given, if ordered by a healthcare provider.
2. Stay with the person; alert emergency contacts.
3. Watch closely for changes. If symptoms worsen, give epinephrine.

MEDICATIONS/DOSES

Epinephrine Brand or Generic: _____

Epinephrine Dose: ☐ 0.1 mg IM (intramuscular) ☐ 0.15 mg IM ☐ 0.3 mg IM ☐ 1 mg IN (intranasal) ☐ 2 mg IN

Antihistamine Brand or Generic: _____

Antihistamine Dose: _____

Other (e.g., inhaler-bronchodilator if wheezing): _____

_____

☐ Patient may self-carry ☐ Patient may self-administer

PATIENT OR PARENT/GUARDIAN AUTHORIZATION SIGNATURE DATE PHYSICIAN/HCP AUTHORIZATION SIGNATURE DATE

FORM PROVIDED COURTESY OF FOOD ALLERGY RESEARCH & EDUCATION (FARE) (FOODALLERGY.ORG) 4/2025

FOOD ALLERGY & ANAPHYLAXIS EMERGENCY CARE PLAN

FARE
Food Allergy Research & Education

HOW TO USE AUVI-Q® (EPINEPHRINE INJECTION, USP), KALEO



1. Remove Auvi-Q from the outer case. Pull off red safety guard.
2. Place black end of Auvi-Q against the middle of the outer thigh.
3. Press firmly until you hear a click and hiss sound, and hold in place for 2 seconds.
4. Call 911 and get emergency medical help right away.

HOW TO USE EPIPEN®, EPIPEN JR® (EPINEPHRINE) AUTO-INJECTOR AND EPINEPHRINE INJECTION (AUTHORIZED GENERIC OF EPIPEN®), USP AUTO-INJECTOR, VIATRIS AUTO-INJECTOR, VIATRIS



1. Remove the EpiPen® or EpiPen Jr® Auto-Injector from the clear carrier tube.
2. Grasp the auto-injector in your fist with the orange tip (needle end) pointing downward. With your other hand, remove the blue safety release by pulling straight up.
3. Swing and push the auto-injector firmly into the middle of the outer thigh until it 'clicks'. Hold firmly in place for 3 seconds (count slowly 1, 2, 3).
4. Remove and massage the injection area for 10 seconds. Call 911 and get emergency medical help right away.

HOW TO USE IMPAX EPINEPHRINE INJECTION (AUTHORIZED GENERIC OF ADRENACLICK®), USP AUTO-INJECTOR, AMNEAL PHARMACEUTICALS



1. Remove epinephrine auto-injector from its protective carrying case.
2. Pull off both blue end caps: you will now see a red tip. Grasp the auto-injector in your fist with the red tip pointing downward.
3. Put the red tip against the middle of the outer thigh at a 90-degree angle, perpendicular to the thigh. Press down hard and hold firmly against the thigh for approximately 10 seconds.
4. Remove and massage the area for 10 seconds. Call 911 and get emergency medical help right away.

HOW TO USE TEVA'S GENERIC EPIPEN® (EPINEPHRINE INJECTION, USP) AUTO-INJECTOR, TEVA PHARMACEUTICAL INDUSTRIES

1. Quickly twist the yellow or green cap off of the auto-injector in the direction of the "twist arrow" to remove it.
2. Grasp the auto-injector in your fist with the orange tip (needle end) pointing downward. With your other hand, pull off the blue safety release.
3. Place the orange tip against the middle of the outer thigh at a right angle to the thigh.
4. Swing and push the auto-injector firmly into the middle of the outer thigh until it 'clicks'. Hold firmly in place for 3 seconds (count slowly 1, 2, 3).
5. Remove and massage the injection area for 10 seconds. Call 911 and get emergency medical help right away.

ADMINISTRATION AND SAFETY INFORMATION FOR ALL AUTO-INJECTORS:

1. Do not put your thumb, fingers or hand over the tip of the auto-injector or inject into any body part other than mid-outer thigh. In case of accidental injection, go immediately to the nearest emergency room.
2. If administering to a young child, hold their leg firmly in place before and during injection to prevent injuries.
3. Epinephrine can be injected through clothing if needed.
4. Call 911 immediately after injection.

HOW TO USE NEFFY® (EPINEPHRINE NASAL SPRAY)

1. Remove neffy from packaging. Pull open the packaging to remove the neffy nasal spray device.
2. Hold device as shown. Hold the device with your thumb on the bottom of the plunger and a finger on either side of the nozzle. Do not pull or push on the plunger. Do not test or prime (pre-spray). Each device has only 1 spray.
3. Insert the nozzle into a nostril until your fingers touch your nose. Keep the nozzle straight into the nose pointed toward your forehead. Do not point (angle) the nozzle to the nasal septum (wall between your 2 nostrils) or outer wall of the nose.
4. Press plunger up firmly until it snaps up and sprays liquid into the nostril. Do not sniff during or after the dose is given. If any liquid drips out of the nose, you may need to give a second dose of neffy after checking for symptoms.
5. If symptoms don't improve or worsen within 5 minutes of initial dose, administer a second dose into the same nostril with a new neffy device.

Treat the person before calling emergency contacts. The first signs of a reaction can be mild, but symptoms can worsen quickly.

EMERGENCY CONTACTS — CALL 911

RESCUE SQUAD: _____

DOCTOR: _____ PHONE: _____

PARENT/GUARDIAN: _____ PHONE: _____

OTHER EMERGENCY CONTACTS

NAME/RELATIONSHIP: _____ PHONE: _____

NAME/RELATIONSHIP: _____ PHONE: _____

NAME/RELATIONSHIP: _____ PHONE: _____

San Mateo Union High School District
Authorization for Medication(s) to be Taken During School Hours

Pursuant to Section 49423 and subdivision (b) of Section 49423.6 of the California Education Code, any pupil who is required to take, during the regular school day, prescribed medication may be assisted by a school nurse or other designated school personnel if both of the following conditions are met: (a) The pupil's authorized health care provider executes a written statement specifying, at a minimum, the medication the pupil is to take, the dosage, and the period of time during which the medication is to be take, as well as otherwise detailing (as may be necessary) the method, amount, and time schedule by which the medication is to be taken. (b) The pupil's parent or legal guardian provides a written statement initiating a request to have the medication administered to the pupil or to have the pupil otherwise assisted in the administration of the medication, in accordance with the authorized health care provider's written statement.
With the approval of the pupil's authorized health care provider and the approval of the pupil's parent or legal guardian, a local education agency may allow a pupil to carry medication and to self-administer the medication.

THE FOLLOWING SECTION IS TO BE COMPLETED BY THE PARENT: School Name_____

Student Name _____Gender_____ Date of Birth_____
Last First

_____(_____)_____
Physician/Health Care Provider's Name Address Telephone

In regards to the medication authorized below by her/his physician/health care provider:

I request that my student be assisted in taking the medicine(s) at school by authorized persons: Yes_____ No_____

I request that my student be permitted to carry medication & self-medicate her/himself: Yes_____ No_____

I understand that the medication must be in the original pharmacy container, labeled with name of student, prescribing health care provider, and medication; date of the original prescription; strength and dose of medication; and directions for use. If medication is kept at school in the health office, it will be destroyed unless picked up within one week after the end of the school year or end of the medical order. I have read and signed the attached consent (reverse side) to allow designated school personnel to consult with my student's health care provider regarding medication questions. I understand that the medication may be discontinued with written parental request. As parent/guardian of the above-named student, I hereby indemnify and hold harmless from any demands, actions, suits, or liability of any nature or kind, any and all personnel, employees, and agents of the San Mateo Union High School District who may act pursuant to the instruction of my student's health care provider.

_____ _____ (_____)_____(_____)_____
Date Signature of Parent/Guardian Home Phone Emergency

THE FOLLOWING SECTION IS TO BE COMPLETED BY THE PHYSICIAN:

Diagnosis for which medication is given:_____

Name of medication:_____

Form: _____Dose & route: _____

If medicine is to be given DAILY, at what time(s):_____

If medicine is to be given WHEN NEEDED, describe indications: _____

How soon can it be repeated?: _____ Length of time this treatment is recommended: _____

List significant side effects of medication: _____

In my opinion, this student shows the capability to carry and self-medicate the above medication: Yes_____ No_____

If necessary, this medication may be safely and appropriately administered by trained unlicensed school personnel: Yes_____ No_____ N/A_____

Date: _____ **Signature of Authorized Health Care Provider:** _____

Health Care Provider Address Stamp (required):

SAN MATEO UNION HIGH SCHOOL DISTRICT
AUTHORIZATION FOR USE OR DISCLOSURE OF HEALTH INFORMATION

Completion of this document authorizes the disclosure and/or use of individually identifiable health information, as set forth below, consistent with California and Federal law concerning the privacy of such information. Failure to provide all information requested may invalidate this authorization. I have a right to receive a copy of this Authorization. Signing this Authorization may be required in order for this student to obtain appropriate services at school.

USE AND DISCLOSURE INFORMATION:

Patient/Student Name: _____
Last First MI Date of Birth

I, the undersigned, do hereby authorize (name of agency and/or health care providers):

(1) _____ (2) _____

to provide health information from the above-named student's medical record to and from:

San Mateo Union High School District	**650 North Delaware St., San Mateo, CA 94401**
School District to which disclosure is made	Address/City and State/Zip Code
Sara Devaney, Health Services Manager	**650-558-2222 (Confidential Fax 650-762-0250)**
Contact person at School District	Area Code and Telephone Number

The disclosure of health information is required for the following purpose:

_____

Requested information shall be limited to the following: □ All health information; or □ Disease-specific information as described:

_____

DURATION: This authorization shall become effective immediately and shall remain in effect until _____ (enter date) or for one year from the date of signature, if no date entered.

RESTRICTIONS: California law prohibits the School District from making further disclosure of my health information unless the School District obtains another authorization form from me or unless such disclosure is specifically required or permitted by law. I understand that the School District will protect this information as prescribed by the Family Educational Rights Privacy Act (FERPA) and state law and that the information becomes part of the student's education record. The information will be shared with individuals working at or with the School District for the purpose of providing safe, appropriate, and least restrictive educational settings and school health services and programs. *If you move to another School District, records will be transferred automatically to that School District.*

YOUR RIGHTS: I understand that I have the following rights with respect to this Authorization: *I may revoke this Authorization at any time. My revocation must be in writing, signed by me or on my behalf, and delivered to the health care agencies/persons listed above. My revocation will be effective upon receipt, but will not be effective to the extent that the School District or others have acted in reliance to this Authorization.*

APPROVAL: _____ _____ _____
Printed Name Signature Date

_____ _____
Relationship to Patient/Student Area Code and Telephone Number