

# Table 1 Recommended Child and Adolescent Immunization Schedule for Ages 18 Years or Younger, United States, 2025

These recommendations must be read with the notes that follow. For those who fall behind or start late, provide catch-up vaccination at the earliest opportunity as indicated by the green bars. To determine minimum intervals between doses, see the catch-up schedule (Table 2).

| Vaccine and other immunizing agents                      | Birth   | 1 mo         | 2 mos     | 4 mos        | 6 mos                                   | 9 mos                           | 12 mos                    | 15 mos | 18 mos   | 19–23 mos | 2–3 yrs         | 4–6 yrs | 7–10 yrs  | 11–12 yrs  | 13–15 yrs | 16 yrs   | 17–18 yrs |  |  |
|--|---|--------------|-----------|--------------|---|---------------------------------|---------------------------|--------|----------|-----------|-----------------|---------|-----------|--|-----------|----------|-----------|--|--|
| Respiratory syncytial virus (RSV-mAb [Nirsevimab])       | 1 dose depending on maternal RSV vaccination status (See Notes) |              |           |              | 1 dose (8 through 19 months), See Notes |                                 |                           |        |          |           |                 |         |           |  |           |          |           |  |  |
| Hepatitis B (HepB)                                       | 1st dose  | ← 2nd dose → |           | ← 3rd dose → |   |                                 |                           |        |          |           |                 |         |           |  |           |          |           |  |  |
| Rotavirus (RV): RV1 (2-dose series), RV5 (3-dose series) |   |              | 1st dose  | 2nd dose     | See Notes                               |                                 |                           |        |          |           |                 |         |           |  |           |          |           |  |  |
| Diphtheria, tetanus, acellular pertussis (DTaP <7 yrs)   |   |              | 1st dose  | 2nd dose     | 3rd dose                                | ← 4th dose →                    |                           |        | 5th dose |           |                 |         |           |  |           |          |           |  |  |
| Haemophilus influenzae type b (Hib)                      |   |              | 1st dose  | 2nd dose     | See Notes                               | ← 3rd or 4th dose (See Notes) → |                           |        |          |           |                 |         |           |  |           |          |           |  |  |
| Pneumococcal conjugate (PCV15, PCV20)                    |   |              | 1st dose  | 2nd dose     | 3rd dose                                | ← 4th dose →                    |                           |        |          |           |                 |         |           |  |           |          |           |  |  |
| Inactivated poliovirus (IPV)                             |   |              | 1st dose  | 2nd dose     | ← 3rd dose →                            |                                 |                           |        | 4th dose |           |                 |         |           |  |           |          |           |  |  |
| COVID-19 (1vCOV-mRNA, 1vCOV-aPS)                         | 1 or more doses of 2024–2025 vaccine (See Notes)                |              |           |              |   |                                 |                           |        |          |           |                 |         |           |  |           |          |           |  |  |
| Influenza (IIV3, cclIV3)                                 | 1 or 2 doses annually   |              |           |              |   |                                 |                           |        |          |           | 1 dose annually |         |           |  |           |          |           |  |  |
| Influenza (LAIV3)  | 1 or 2 doses annually   |              |           |              |   |                                 |                           |        |          |           | 1 dose annually |         |           |  |           |          |           |  |  |
| Measles, mumps, rubella (MMR)                            |   |              |           |              | See Notes                               |                                 | ← 1st dose →              |        | 2nd dose |           |                 |         |           |  |           |          |           |  |  |
| Varicella (VAR)  |   |              |           |              |   |                                 | ← 1st dose →              |        | 2nd dose |           |                 |         |           |  |           |          |           |  |  |
| Hepatitis A (HepA)                                       |   |              |           |              | See Notes                               |                                 | 2-dose series (See Notes) |        |          |           |                 |         |           |  |           |          |           |  |  |
| Tetanus, diphtheria, acellular pertussis (Tdap ≥7 yrs)   |   |              |           |              |   |                                 |                           |        |          |           |                 |         | 1 dose    |  |           |          |           |  |  |
| Human papillomavirus (HPV)                               |   |              |           |              |   |                                 |                           |        |          |           |                 |         | See Notes |  |           |          |           |  |  |
| Meningococcal (MenACWY-CRM ≥2 mos, MenACWY-TT ≥2years)   |   |              | See Notes |              |   |                                 |                           |        |          |           |                 |         |           |  | 1st dose  | 2nd dose |           |  |  |
| Meningococcal B (MenB-4C, MenB-FHbp)                     |   |              |           |              |   |                                 |                           |        |          |           |                 |         |           | See Notes  |           |          |           |  |  |
| Respiratory syncytial virus vaccine (RSV [Abrysvo])      |   |              |           |              |   |                                 |                           |        |          |           |                 |         |           | Seasonal administration during pregnancy (See Notes) |           |          |           |  |  |
| Dengue (DEN4CYD: 9–16 yrs)                               |   |              |           |              |   |                                 |                           |        |          |           |                 |         |           | Seropositive in endemic dengue areas (See Notes)     |           |          |           |  |  |
| Mpox   |   |              |           |              |   |                                 |                           |        |          |           |                 |         |           |  |           |          |           |  |  |

Range of recommended ages for all children
Range of recommended ages for catch-up vaccination
Range of recommended ages for certain high-risk groups or populations
Recommended vaccination can begin in this age group
Recommended vaccination based on shared clinical decision-making
No Guidance/Not Applicable