


PARENT & ATHLETE CONCUSSION INFORMATION SHEET



WHAT IS A CONCUSSION?

A concussion is a type of traumatic brain injury that changes the way the brain normally works. A concussion is caused by a bump, blow, or jolt to the head or body that causes the head and brain to move quickly back and forth. Even a "ding," "getting your bell rung," or what seems to be a mild bump or blow to the head can be serious.

WHAT ARE THE SIGNS AND SYMPTOMS OF CONCUSSION?

Signs and symptoms of concussion can show up right after the injury or may not appear or be noticed until days or weeks after the injury.

If an athlete reports one or more symptoms of concussion after a bump, blow, or jolt to the head or body, s/he should be kept out of play the day of the injury. The athlete should only return to play with permission from a health care professional experienced in evaluating for concussion.

SYMPTOMS REPORTED BY ATHLETE:

- Headache or "pressure" in head
- Nausea or vomiting
- Balance problems or dizziness
- Double or blurry vision
- Sensitivity to light
- Sensitivity to noise
- Feeling sluggish, hazy, foggy, or groggy
- Concentration or memory problems
- Confusion
- Just not "feeling right" or is "feeling down"

DID YOU KNOW?

- Most concussions occur without loss of consciousness.
- Athletes who have, at any point in their lives, had a concussion have an increased risk for another concussion.
- Young children and teens are more likely to get a concussion and take longer to recover than adults.

SIGNS OBSERVED BY COACHING STAFF:

- Appears dazed or stunned
- Is confused about assignment or position
- Forgets an instruction
- Is unsure of game, score, or opponent
- Moves clumsily
- Answers questions slowly
- Loses consciousness (even briefly)
- Shows mood, behavior, or personality changes
- Can't recall events prior to hit or fall
- Can't recall events after hit or fall

Michigan Department
of Community Health



Rick Snyder, Governor
James K. Haveman, Director

► **"IT'S BETTER TO MISS ONE GAME
THAN THE WHOLE SEASON"**

CONCUSSION DANGER SIGNS

In rare cases, a dangerous blood clot may form on the brain in a person with a concussion and crowd the brain against the skull. An athlete should receive immediate medical attention if after a bump, blow, or jolt to the head or body s/he exhibits any of the following danger signs:

- One pupil larger than the other
- Is drowsy or cannot be awakened
- A headache that gets worse
- Weakness, numbness, or decreased coordination
- Repeated vomiting or nausea
- Slurred speech
- Convulsions or seizures
- Cannot recognize people or places
- Becomes increasingly confused, restless, or agitated
- Has unusual behavior
- Loses consciousness (even a brief loss of consciousness should be taken seriously)

WHAT SHOULD YOU DO IF YOU THINK YOUR ATHLETE HAS A CONCUSSION?

1. If you suspect that an athlete has a concussion, remove the athlete from play and seek medical attention. Do not try to judge the severity of the injury yourself. Keep the athlete out of play the day of the injury and until a health care professional, experienced in evaluating for concussion, says s/he is symptom-free and it's OK to return to play.
2. Rest is key to helping an athlete recover from a concussion. Exercising or activities that involve a lot of concentration, such as studying, working on the computer, and playing video games, may cause concussion symptoms to reappear or get worse. After a concussion, returning to sports and school is a gradual process that should be carefully managed and monitored by a health care professional.
3. Remember: Concussions affect people differently. While most athletes with a concussion recover quickly and fully, some will have symptoms that last for days, or even weeks. A more serious concussion can last for months or longer.

WHY SHOULD AN ATHLETE REPORT THEIR SYMPTOMS?

If an athlete has a concussion, his/her brain needs time to heal. While an athlete's brain is still healing, s/he is much more likely to have another concussion. Repeat concussions can increase the time it takes to recover. In rare cases, repeat concussions in young athletes can result in brain swelling or permanent damage to their brain. They can even be fatal.

STUDENT-ATHLETE NAME PRINTED

STUDENT-ATHLETE NAME SIGNED

DATE

PARENT OR GUARDIAN NAME PRINTED

PARENT OR GUARDIAN NAME SIGNED

DATE

JOIN THE CONVERSATION  www.facebook.com/CDCHeadsUp

TO LEARN MORE GO TO

Educational Material for Parents and Students (Content from MDHHS Requirements)

Sources: Michigan Dept. of Health and Human Services. Created through a grant to the CDC Foundation from NOCSAE.

UNDERSTANDING CONCUSSION

Some Common Symptoms

Headache	Balance Problems	Sensitive to Noise	Poor Concentration	Not "Feeling Right"
Pressure in the Head	Double Vision	Sluggishness	Memory Problems	Feeling Irritable
Nausea/Vomiting	Blurry Vision	Haziness	Confusion	Slow Reaction Time
Dizziness	Sensitive to Light	Fogginess	"Feeling Down"	Sleep Problems
		Grogginess	Lost Consciousness	

WHAT IS A CONCUSSION?

A **concussion** is a **type of traumatic brain injury** that changes the way the brain normally works. A concussion is caused by a fall, bump, blow, or jolt to the head or body that causes the head and brain to move quickly back and forth. A concussion can be caused by a shaking, spinning or a sudden stopping and starting of the head. Even a "ding," "getting your bell rung," or what seems to be a mild bump or blow to the head can be serious. A concussion can happen even if you haven't been knocked out.

You can't see a concussion. Signs and symptoms of concussions can show up right after the injury or may not appear or be noticed until days or weeks after the injury. If the student reports any symptoms of a concussion, or if you notice symptoms yourself, seek medical attention right away. A student who may have had a concussion should not return to play on the day of the injury and until a health care professional says they are okay to return to play.

IF YOU SUSPECT A CONCUSSION:

- 1. SEEK MEDICAL ATTENTION RIGHT AWAY – DON'T HIDE IT, REPORT IT.** Playing or practicing with concussion symptoms is dangerous and can lead to a longer recovery. A health care professional will be able to decide how serious the concussion is and when it is safe for the student to return to regular activities, including sports. Ignoring symptoms and trying to "tough it out" often makes it worse.
- 2. KEEP YOUR STUDENT OUT OF PLAY –** Concussions take time to heal. Don't let the student return to play the day of injury and until a health care professional says it's okay. A student, who returns to play too soon, while the brain is still healing, risks a greater chance of having a second concussion. Young children and teens are more likely to get a concussion and take longer to recover than adults. Repeat or second concussions increase the time it takes to recover and can be very serious. They can cause permanent brain damage, affecting the student for a lifetime. They can be fatal. It is better to miss one game than the whole season.
- 3. TELL THE SCHOOL ABOUT ANY PREVIOUS CONCUSSION –** Schools should know if a student had a previous concussion. A student's school may not know about a concussion received in another sport or activity unless you notify them.

SIGNS OBSERVED BY PARENTS:

- Appears dazed or stunned
- Is confused about assignment or position
- Forgets an instruction
- Can't recall events prior to or after a hit or fall
- Is unsure of game, score, or opponent
- Moves clumsily
- Answers questions slowly
- Loses consciousness (even briefly)
- Shows mood, behavior, or personality changes

CONCUSSION DANGER SIGNS:

In rare cases, a dangerous blood clot may form on the brain in a person with a concussion and crowd the brain against the skull. A student should receive immediate medical attention if after a bump, blow, or jolt to the head or body s/he exhibits any of the following danger signs:

- One pupil larger than the other
- Is drowsy or cannot be awakened
- A headache that gets worse
- Weakness, numbness, or decreased coordination
- Repeated vomiting or nausea
- Slurred speech
- Convulsions or seizures
- Cannot recognize people/places
- Becomes increasingly confused, restless or agitated
- Has unusual behavior
- Loses consciousness (even a brief loss of consciousness should be taken seriously.)

HOW TO RESPOND TO A REPORT OF A CONCUSSION:

If a student reports one or more symptoms of a concussion after a bump, blow, or jolt to the head or body, s/he must be kept out of athletic activity the day of the injury. The student shall only return to activity (practice, scrimmage or competition) with written unconditional permission from an MD, DO, Physician's Assistant or Nurse Practitioner. During recovery, rest is key. Exercising or activities that involve a lot of concentration (such as studying, working on the computer, or playing video games) may cause concussion symptoms to reappear or get worse. Students who return to school after a concussion may need to spend fewer hours at school, take rests breaks, be given extra help and time, spend less time reading, writing or on a computer. After a concussion, returning to sports and school is a gradual process that should be monitored by a health care professional.

Remember: Concussion affects people differently. While most students with a concussion recover quickly and fully, some will have symptoms that last for days, or even weeks. A more serious concussion can last for months or longer.

To learn more, go to www.cdc.gov/concussion.