

Action Care Plan: Prader-Willis

Overview: Prader-Willi Syndrome is a genetic disorder usually caused by a deletion of part of chromosome 15. Symptoms may include but not limited to, intellectual disability, behavioral concerns, delayed growth, poor muscle tone, exhaustion, excessive hunger/thirst, difficulty with temperature regulation, and high pain tolerance.

| Signs & Symptoms | Teacher's Action Plan |
|--|--|
| <ul style="list-style-type: none"> Refusing to drink Consuming food/drink excessively Overheated to include flushed face, sweating, confusion, fatigue, headache, fainting Vomiting or distended abdomen Skin easily turns red if touched too long. | <ul style="list-style-type: none"> If any of these symptoms are noted, call parents immediately. If any symptoms of distended abdomen, vomiting, extreme fatigue, overexertion, or dehydration, notify parents AND 911. All food and drink recommendations will be provided by parents for safety. ABSOLUTELY NO OUTSIDE FOOD AND DRINK without parent consent. If red mark on skin does not resolve quickly after pressure of position or object on skin, notify parents |

WHEN TO CALL 911

If the student presents with vomiting, abdominal distention, extreme fatigue, overexertion, or dehydration, notify health assistant and call EMS/9-1-1.

[illegible]