

Eastern Lebanon County School District Standing Order for Medications

Emergency Medications

Oxygen: To be used when an individual is displaying signs of lack of oxygen.

Call 911

Call parent/guardian

Call family physician/school physician if parent cannot be reached

Epinephrine: For anaphylactic reaction to bee stings, insect bites, medication, food allergies. Dose to appropriate age/weight. (Epi-Pen or Epi-Pen Jr based on weight)

Call 911

Call parent/guardian

Call family physician/school physician if parent cannot be reached

Aspirin, ASA oral tablets: For adult use only due to a cardiac event. Tablets should be chewed. Dose one 325 mg tablet or 4 baby ASA.

Call 911

Call emergency contact

Naloxone: To be used in treating opioid drug overdoses. Dose: Injectable solution 0.4mg/mL; IM/SC auto-injector 0.4mg/0.4mL.

Call 911

Call parent/guardian or emergency contact if being administered to an adult

Call family physician/school physician if parent cannot be reached

Diphenhydramine Hydrochloride (Benadryl): To be used for localized swelling, redness, itching associated with reaction to medication, latex, food, insect bites or stings, or chemicals. Dose to appropriate age/weight.

Non-emergency Medications

Parental permission required via the "Annual OTC Medication Permission Form" prior to administration

Acetaminophen: Confirm permission to administer. Check temperature. If temperature is less than 100° F, acetaminophen may be administered for complaints from colds, headaches, general aches and pains, menses discomfort. Temperatures of 100.0° F or higher, parent or guardian should be called. Dose appropriate to age/weight. Not to exceed a single dose in a school day. Provide written documentation to parent of administration.

Ibuprofen: Confirm permission to administer. Check temperature. If temperature is 100° F or less, ibuprofen may be administered for complaints from colds, headaches, general aches and pains, menses discomfort. Temperatures of 100.0° F or higher, parent or guardian should be called. Dose appropriate to age/weight with food. Not to exceed a single dose in a school day. Provide written documentation to parent of administration.

These OTC medications or generic equivalents, commonly requested for episodic symptom relief, may be administered by the school/building nurse at his/her discretion, as directed by the package dosage guidelines. Tums, Ibuprofen, and Acetaminophen not to exceed a single dose in one school day. Confirm permission to administer. Provide written documentation to parent of administration.

Topical: Bacitracin, Calamine/Caladryl Lotion, Hydrocortisone cream 1%

Mouth and Throat: Anbesol, Throat Lozenges

Antacid: Tums (12 years old and older)

Artificial Tears: Refresh

Anesthetic Swabs: Sting Kill



Steven Shay, D.O.

6/17/25
Date

Epinephrine Standing Order Protocol

I, the undersigned Physician, for the purpose of facilitating the use of epinephrine in the case of anaphylaxis, a life-threatening allergic reaction, in individuals and in compliance with all applicable state laws and regulations, issue this epinephrine standing order Protocol ("Protocol") on the following terms:

Physician License: I represent that I: (a) am licensed to prescribe legend drugs in this state as set forth below; (b) am qualified to practice medicine in this state; and (c) am in good standing with the appropriate professional licensing board.

Epinephrine: This Protocol constitutes my standing order for the treatment of anaphylaxis and the use of epinephrine in emergency situations as further described below in a school setting.

Delegation: I, the undersigned Physician, delegate authority to all appropriate medical and school personnel employed by or acting on behalf of the below described school system.

Issued to: Eastern Lebanon County School District
Name of School/District
180 ELCO Drive
Street Address
Myerstown PA 17067
City, Zip Code


Standing Order: All appropriate medical and school personnel (including, but not limited to, any Registered Nurse) employed by or acting on behalf of the school system may administer epinephrine via an undesignated epinephrine auto-injector to an individual using professional judgment if an individual is experiencing a potentially life-threatening allergic reaction (anaphylaxis).

Emergency Treatment Procedures: The following treatment Protocol will be utilized to manage anaphylactic reactions. Anaphylaxis is a life-threatening allergic reaction that is rapid in onset.

1. **Dosage:** If conditions of anaphylaxis are developing or present themselves, administer epinephrine USP as epinephrine auto-injector, EpiPen® (epinephrine injection, USP) or EpiPen Jr® (epinephrine injection, USP) Auto-Injector, or the authorized generic for EpiPen® or EpiPen Jr® Auto-Injector, intramuscularly into the anterolateral aspect of the thigh (through clothing if necessary). Selection of the appropriate dosage strength (EpiPen® 0.3 mg or EpiPen Jr® 0.15 mg, or the authorized generics of these strengths) is determined according to patient body weight, as discussed in the product labeling.
 - a. For individuals 33 to 66 pounds, use one EpiPen Jr® (0.15 mL epinephrine injection, USP) Auto-Injector or the authorized generic for EpiPen Jr® Auto-Injector to deliver 0.15 mg of epinephrine injection, USP.
 - b. For individuals approximately 66 pounds and greater, use one EpiPen® (0.3 mL epinephrine injection, USP) Auto-Injector or the authorized generic for EpiPen® Auto-Injector to deliver 0.3 mg of epinephrine injection, USP.
2. **Frequency:** Up to 20% of individuals who receive epinephrine will require more than one dose before symptoms are alleviated. More than two sequential doses of epinephrine for the same episode should be administered only under direct medical supervision.
3. **Referral:** The individual must be referred to a physician for medical evaluation, even if symptoms resolve completely. Symptoms may recur after the epinephrine wears off, as much as 24 hours later.
4. **Documentation and Notification:** Document the details of the incident and notify the individual's parent, guardian, or caretaker and primary care physician in accordance with school policy.

In every case, emergency services must be contacted as soon as possible by calling 911 or local emergency medical services.

Please review the attached prescription:

Effective Date: 6/17/25
Physician Signature: 

Physician Name (printed): Steven Shay, DO
Physician Contact Number: 717-949-2777 / 717-949-6841
Physician Address: 2496 Striegel Pike, Schaefferstown, PA 17088
Physician State of License: PA
Physician State License Number: 05-012088

Strength	Quantity Requested*
0.3mg EpiPen 2-Pak®	
0.15mg EpiPen Jr 2-Pak®	

*Please note there are two auto-injectors in each EpiPen 2-Pak® or EpiPen Jr 2-Pak®. Example: If you wish to order 100 EpiPen® Auto-Injectors and 80 EpiPen Jr® Auto-Injectors, put the number 50 in the quantity requested box next to the 0.3mg EpiPen 2-Pak® and 40 in the quantity requested box next to the 0.15mg EpiPen Jr 2-Pak®. **Please note that schools may receive the authorized generic versions of EpiPen® and EpiPen Jr®.**

Indications

EpiPen® (epinephrine injection, USP) 0.3 mg and EpiPen Jr® (epinephrine injection, USP) 0.15 mg Auto-Injectors are indicated in the emergency treatment of Type I allergic reactions, including anaphylaxis, to allergens, idiopathic and exercise-induced anaphylaxis, and in patients with a history or increased risk of anaphylactic reactions. Selection of the appropriate dosage strength is determined according to body weight.

Important Safety Information

EpiPen® and EpiPen Jr® Auto-Injectors are intended for immediate administration as emergency supportive therapy only and are not intended as a substitute for immediate medical or hospital care. **In conjunction with the administration of epinephrine, the patient should seek immediate medical or hospital care.** More than two sequential doses of epinephrine should only be administered under direct medical supervision.

EpiPen® and EpiPen Jr® should **only** be injected into the anterolateral aspect of the thigh. **Do not inject intravenously, into buttock, or into digits, hands, or feet.** Instruct caregivers to hold the leg of young children firmly in place and limit movement prior to and during injection to minimize risk of injection-related injury.

Rare cases of serious skin and soft tissue infections have been reported following epinephrine injection. Advise patients to seek medical care if they develop symptoms of infection such as persistent redness, warmth, swelling, or tenderness at the injection site.

Epinephrine should be used with caution in patients with heart disease, and in patients who are on drugs that may sensitize the heart to arrhythmias, because it may precipitate or aggravate angina pectoris and produce ventricular arrhythmias. Arrhythmias, including fatal ventricular fibrillation, have been reported, particularly in patients with underlying cardiac disease or taking cardiac glycosides, diuretics, or anti-arrhythmics.

Patients with certain medical conditions or who take certain medications for allergies, depression, thyroid disorders, diabetes, and hypertension, may be at greater risk for adverse reactions. Common adverse reactions to epinephrine include anxiety, apprehensiveness, restlessness, tremor, weakness, dizziness, sweating, palpitations, pallor, nausea and vomiting, headache, and/or respiratory difficulties.

Please see the full [Prescribing Information](#) and [Patient Information](#).

For additional information please contact us at 800-395-3376.

You are encouraged to report negative side effects of prescription drugs to the FDA.

Visit www.fda.gov/medwatch or call 1-800-FDA-1088.

EpiPen®, EpiPen Jr®, EpiPen 2-Pak®, EpiPen Jr 2-Pak® and *EpiPen4Schools*® are registered trademarks owned by the Mylan companies. © 2019 Mylan Specialty L.P. All rights reserved.