

**Epinephrine Auto Injector Self-Administration
Authorization Packet for Anaphylaxis**
(A new packet must be completed yearly)

Packet Contents:

1. Anaphylaxis Medication Self-Administration Form
(requires physician and parent/guardian signature)
2. Anaphylaxis Student Skills Checklist
3. Severe Allergy Action Plan (requires physician and
parent/guardian signature)
4. Missouri Revised Statutes – August 2016

EPINEPHRINE AUTO INJECTOR SELF-ADMINISTRATION FORM

Student Name: _____ Grade: _____ School Year: _____

The Missouri Safe Schools Act of 1996 provides for students to carry and self-administer life-saving medications when the following criteria are met:

- (1) A licensed physician prescribed or ordered the medication for use by the child and instructed such child in the correct and responsible use of the medication.*
- (2) The child has demonstrated to the child's licensed physician or the licensed physician's designee, and the school nurse, if available, the skill level necessary to use the medication and any device necessary to administer such medication prescribed or ordered.*
- (3) The child's physician has approved and signed a written treatment plan for managing asthma or anaphylaxis episodes of the child and for medication for use by the child. Such plan shall include a statement that the child is capable of self-administering the medication under the treatment plan.*
- (4) The child's parent or guardian has completed and submitted to the school any written documentation required by the school, including the treatment plan required in (3) above and the liability statement required in (5) below.*
- (5) The child's parent or guardian has signed a statement acknowledging that the school district and its employees or agents shall incur no liability as a result of any injury arising from the self-administration of medication by the child or the administration of such medication by school staff. Such statement shall not be construed to release the school district and its employees or agents from liability for negligence. (Missouri Revised Statute; Chapter 167; Pupils and Special Services; Section 167.627.1; 08-28-2016).*

MEDICATION NAME _____ Dose _____ Time or Interval _____
Route/Inhalation device _____ Instructions _____

MEDICATION NAME _____ Dose _____ Time or Interval _____
Route/Inhalation device _____ Instructions _____

ALLERGIES: List known allergies to medications, foods, or air-borne substances _____

I, THE PARENT OR LEGAL GUARDIAN OF THE STUDENT LISTED ABOVE, GIVE PERMISSION FOR THIS CHILD TO CARRY AND SELF-ADMINISTER THE ABOVE LISTED MEDICATIONS. I HAVE INSTRUCTED MY CHILD TO NOTIFY THE SCHOOL STAFF ANYTIME THIS DEVICE IS USED. I UNDERSTAND THAT, ABSENT ANY NEGLIGENCE, THE SCHOOL SHALL INCUR NO LIABILITY AS A RESULT OF ANY INJURY ARISING FROM THE SELF-ADMINISTRATION OF MEDICATION BY MY CHILD.

SIGNATURE OF PARENT OR LEGAL
GUARDIAN _____

DATE _____

PARENT/GUARDIAN:

NAME: _____ HOME PHONE: _____

ADDRESS: _____ WORK AND CELL PHONES: _____

NAME: _____ HOME PHONE: _____

ADDRESS: _____ WORK AND CELL PHONES: _____

EMERGENCY CONTACT:

NAME: _____ PHONE #'S: _____

I, A LICENSED PHYSICIAN OR NURSE PRACTITIONER, CERTIFY THAT THIS CHILD HAS A MEDICAL HISTORY OF SEVERE ALLERGIC REACTIONS, HAS BEEN TRAINED IN THE USE OF THE LISTED MEDICATION, AND IS JUDGED TO BE CAPABLE OF CARRYING AND SELF-ADMINISTERING THE LISTED MEDICATION(S). THE CHILD SHOULD NOTIFY SCHOOL STAFF ANYTIME THE MEDICATION/INJECTOR IS USED. THIS CHILD UNDERSTANDS THE HAZARDS OF SHARING MEDICATIONS WITH OTHERS AND HAS AGREED TO REFRAIN FROM THIS PRACTICE.

SIGNATURE OF HEALTH CARE PROVIDER _____ DATE _____

NAME OF HEALTH CARE PROVIDER _____ PHONE: _____ FAX: _____

ADDRESS: _____ CITY: _____ ZIP: _____



FARE
Food Allergy Research & Education

FOOD ALLERGY & ANAPHYLAXIS EMERGENCY CARE PLAN

Name: _____ D.O.B.: _____

Allergy to: _____

Weight: _____ lbs. Asthma: [] Yes (higher risk for a severe reaction) [] No

PLACE
PICTURE
HERE

NOTE: Do not depend on antihistamines or inhalers (bronchodilators) to treat a severe reaction. USE EPINEPHRINE.

Extremely reactive to the following allergens: _____

THEREFORE:

- [] If checked, give epinephrine immediately if the allergen was **LIKELY** eaten, for **ANY** symptoms.
- [] If checked, give epinephrine immediately if the allergen was **DEFINITELY** eaten, even if no symptoms are apparent.

FOR ANY OF THE FOLLOWING: **SEVERE SYMPTOMS**



LUNG

Shortness of breath, wheezing, repetitive cough



HEART

Pale or bluish skin, faintness, weak pulse, dizziness



THROAT

Tight or hoarse throat, trouble breathing or swallowing



MOUTH

Significant swelling of the tongue or lips



SKIN

Many hives over body, widespread redness



GUT

Repetitive vomiting, severe diarrhea



OTHER

Feeling something bad is about to happen, anxiety, confusion

**OR A
COMBINATION**
of symptoms
from different
body areas.

1. **INJECT EPINEPHRINE IMMEDIATELY.**
2. **Call 911.** Tell emergency dispatcher the person is having anaphylaxis and may need epinephrine when emergency responders arrive.
- Consider giving additional medications following epinephrine:
 - » Antihistamine
 - » Inhaler (bronchodilator) if wheezing
- Lay the person flat, raise legs and keep warm. If breathing is difficult or they are vomiting, let them sit up or lie on their side.
- If symptoms do not improve, or symptoms return, more doses of epinephrine can be given about 5 minutes or more after the last dose.
- Alert emergency contacts.
- Transport patient to ER, even if symptoms resolve. Patient should remain in ER for at least 4 hours because symptoms may return.

MILD SYMPTOMS



NOSE

Itchy or runny nose, sneezing



MOUTH

Itchy mouth



SKIN

A few hives, mild itch



GUT

Mild nausea or discomfort

**FOR MILD SYMPTOMS FROM MORE THAN ONE
SYSTEM AREA, GIVE EPINEPHRINE.**

**FOR MILD SYMPTOMS FROM A SINGLE SYSTEM
AREA, FOLLOW THE DIRECTIONS BELOW:**

1. Antihistamines may be given, if ordered by a healthcare provider.
2. Stay with the person; alert emergency contacts.
3. Watch closely for changes. If symptoms worsen, give epinephrine.

MEDICATIONS/DOSES

Epinephrine Brand or Generic: _____

Epinephrine Dose: [] 0.15 mg IM [] 0.3 mg IM

Antihistamine Brand or Generic: _____

Antihistamine Dose: _____

Other (e.g., inhaler-bronchodilator if wheezing): _____

PATIENT OR PARENT/GUARDIAN AUTHORIZATION SIGNATURE

DATE

PHYSICIAN/HCP AUTHORIZATION SIGNATURE


DATE

**FARE**


FOOD ALLERGY RESEARCH & EDUCATION

FOOD ALLERGY & ANAPHYLAXIS EMERGENCY CARE PLAN**HOW TO USE AUVI-Q® (EPINEPHRINE INJECTION, USP), KALEO**


1. Remove Auvi-Q from the outer case.
2. Pull off red safety guard.
3. Place black end of Auvi-Q against the middle of the outer thigh.
4. Press firmly, and hold in place for 5 seconds.
5. Call 911 and get emergency medical help right away.

**HOW TO USE EPIPEN® AND EPIPEN JR® (EPINEPHRINE) AUTO-INJECTOR, MYLAN**


1. Remove the EpiPen® or EpiPen Jr® Auto-Injector from the clear carrier tube.
2. Grasp the auto-injector in your fist with the orange tip (needle end) pointing downward.
3. With your other hand, remove the blue safety release by pulling straight up.
4. Swing and push the auto-injector firmly into the middle of the outer thigh until it 'clicks'.
5. Hold firmly in place for 3 seconds (count slowly 1, 2, 3).
6. Remove and massage the injection area for 10 seconds.
7. Call 911 and get emergency medical help right away.

**HOW TO USE EPINEPHRINE INJECTION (AUTHORIZED GENERIC OF EPIPEN®), USP AUTO-INJECTOR, MYLAN**

1. Remove the epinephrine auto-injector from the clear carrier tube.
2. Grasp the auto-injector in your fist with the orange tip (needle end) pointing downward.
3. With your other hand, remove the blue safety release by pulling straight up.
4. Swing and push the auto-injector firmly into the middle of the outer thigh until it 'clicks'.
5. Hold firmly in place for 3 seconds (count slowly 1, 2, 3).
6. Remove and massage the injection area for 10 seconds.
7. Call 911 and get emergency medical help right away.

**HOW TO USE IMPAX EPINEPHRINE INJECTION (AUTHORIZED GENERIC OF ADRENALCLICK®), USP AUTO-INJECTOR, IMPAX LABORATORIES**

1. Remove epinephrine auto-injector from its protective carrying case.
2. Pull off both blue end caps; you will now see a red tip.
3. Grasp the auto-injector in your fist with the red tip pointing downward.
4. Put the red tip against the middle of the outer thigh at a 90-degree angle, perpendicular to the thigh.
5. Press down hard and hold firmly against the thigh for approximately 10 seconds.
6. Remove and massage the area for 10 seconds.
7. Call 911 and get emergency medical help right away.

**ADMINISTRATION AND SAFETY INFORMATION FOR ALL AUTO-INJECTORS:**

1. Do not put your thumb, fingers or hand over the tip of the auto-injector or inject into any body part other than mid-outer thigh. In case of accidental injection, go immediately to the nearest emergency room.
2. If administering to a young child, hold their leg firmly in place before and during injection to prevent injuries.
3. Epinephrine can be injected through clothing if needed.
4. Call 911 immediately after injection.

OTHER DIRECTIONS/INFORMATION (may self-carry epinephrine, may self-administer epinephrine, etc.):

Treat the person before calling emergency contacts. The first signs of a reaction can be mild, but symptoms can worsen quickly.

EMERGENCY CONTACTS — CALL 911

RESCUE SQUAD: _____

DOCTOR: _____ PHONE: _____

PARENT/GUARDIAN: _____ PHONE: _____

OTHER EMERGENCY CONTACTS

NAME/RELATIONSHIP: _____

PHONE: _____

NAME/RELATIONSHIP: _____

PHONE: _____

Epi-Pen/Adrenaclick/Auvi-Q Student Skills Checklist

Student _____ Age _____ Grade/Team _____

School Nurse Signature _____ Date _____

EpiPen and EpiPen Jr Trainer Skills Checklist:

1. Remove the epipen or epipen jr auto-injector from the clear carrier tube.
2. Grasp the auto-injector in your fist with the orange tip (needle end) pointing downward.
3. With your other hand, remove the blue safety release by pulling straight up.
4. Swing and push the auto-injector firmly into the middle of the outer thigh until it 'clicks'.
5. hold firmly in place for 3 seconds.
6. Remove and massage the injection area for 10 seconds.
7. Verbalize that you will tell the school nurse whenever you use the EpiPen. If the school nurse is unavailable, you will tell the principal or appropriate district personnel.

_____ Requires Supervision

_____ Performs Independently

Generic Epinephrine Trainer Skills Checklist:

1. Remove the epinephrine auto-injector from the clear carrier tube.
2. Grasp the auto-injector in your fist with the orange tip (needle end) pointing downward.
3. With your other hand, remove the blue safety release by pulling straight up.
4. Swing and push the auto-injector firmly into the middle of the outer thigh until it 'clicks'.
5. Hold firmly in place for 3 seconds; then remove.
6. Remove and massage the injection area for 10 seconds.
7. Verbalize that you will tell the school nurse whenever you use the EpiPen. If the school nurse is unavailable, you will tell the principal or appropriate district personnel.

_____ Requires Supervision

_____ Performs independently

IMPAX Epinephrine (generic Adrenaclick) Trainer Skills Checklist:

1. Remove the epinephrine auto-injector from the clear carrier tube.
2. Pull off both blue end caps; you will now see a red tip.
3. Grasp the auto-injector in your fist with the red tip pointing downward.
4. Put Red tip against the middle of the outer thigh at a 90-degree angle, perpendicular to the thigh.
5. Press down hard and hold firmly against the thigh for 10 seconds, then remove.
6. Verbalize that you will tell the school nurse whenever you use the impax. If the school nurse is unavailable, you will tell the principal or appropriate district personnel.

_____ Requires Supervision

_____ Performs Independently

Auvi-Q Trainer Skills Checklist:

1. Remove auvi-q from the outer case. This will automatically activate the voice instructions.
2. Pull off red safety guard.
3. Place black end of auvi-q against the middle of the outer thigh.
4. Press firmly and hold for 5 seconds.
5. Remove from thigh.
6. Verbalize that you will tell the school nurse whenever you use Auvi-Q. If the school nurse is unavailable, you will tell the principal or appropriate district personnel.

_____ Requires Supervision

_____ Performs Independently